

# NONHOLONOMIC SYSTEMS AND THE HAMILTONIZATION PROBLEM.

**Paula Balseiro**

Instituto de Matemática e Estatística  
Universidade Federal Fluminense  
P.O. Box 24020-140  
Niteroi, Rio de Janeiro, Brazil  
pbalseiro@vm.uff.br

## **Abstract**

A nonholonomic system is a mechanical system with constraints in the velocities and thus the equations of motion are not hamiltonian. This fact has consequences in the dynamical behaviour as well as in the geometry underlying the mechanical system. In this talk we will study geometric features of nonholonomic systems and their behaviour after a reduction by a group of symmetries. In particular, we are interested in how Poisson geometry may help in understanding how far these systems are from being hamiltonian.

## REFERENCES

- [1] P. Balseiro, The Jacobiator of nonholonomic systems and the geometry of reduced non-holonomic brackets. Preprint, 2013. arXiv:1301.1091.
- [2] P. Balseiro, L. Garcia-Naranjo, Gauge Transformations, Twisted Poisson Brackets and Hamiltonization of Nonholonomic Systems, *Archive for Rational Mechanics and Analysis*, **205**, 1 (2012), 267-310.

# ON THE LOCATION OF SATURN F RING: A SIMPLE MODEL

**Luis Benet**

Instituto de Ciencias Físicas  
Universidad Nacional Autónoma de México  
Cuernavaca, Mor, Mexico  
benet@fis.unam.mx

## Abstract

The confinement of planetary narrow rings is understood in terms of the shepherd theory [1]. This theory proposes the existence of two shepherd moons that orbit around the ring, and involves angular momentum transfer mechanisms between them and the ring particles, self-gravity, viscous damping, and assumes the existence of lower-order resonance at the ring boundaries in order to confine the ring. Saturn's F ring is a fascinating example of a narrow eccentric ring that displays a rich dynamical structure: besides its non-zero eccentricity and sharp edges, it has multiple components entangled in a complicated way which shows a variety of short-time features [2]. The two shepherd moons, Prometheus and Pandora, influence importantly many of the short-time dynamical features observed. Yet, the shepherd theory does not apply to this ring since there are no mean-motion resonances that confine the ring and the mass of the shepherd moons is too small [3]. The confinement of Saturn's F-ring thus remains unexplained. In this talk, I shall describe our scattering approach to narrow rings [4], illustrating it on a simple billiard system, where the relevant physics can be easily understood. Within this approach, I shall present numerical results based of accurate long-time integrations on a realistic 5-body model for the occurrence and location of Saturn's F ring. Test particles that remain trapped and display some stability properties form a narrow elliptic ring displaying sharp edges. Comparison of our results with the observations shall be provided. This work is in collaboration with Àngel Jorba (U. Barcelona).

## REFERENCES

- [1] P. Goldreich, S.D. Tremaine, *Nature* **277**, 1979, 97–99.
- [2] J.E. Colwell, et al., in *Saturn from Cassini-Huygens*, Dougherty, M.K., Esposito, L.W., Krimigis, S.M. (eds.), Springer, 2009, pp 375-412.
- [3] L.W. Esposito, *Planetary rings*, Cambridge University Press, 2006.
- [4] O. Merlo and L. Benet, *Celest. Mech. Dyn. Astr.* **97**, 2007, 49-72.

# VORTEX DYNAMICS ON SURFACES OF REVOLUTION

Stefanella Boatto<sup>1,2</sup>, David Dritschel<sup>3</sup> and Carles Simó<sup>4</sup>

<sup>1</sup>Depto de Matematica Aplicada, Instituto de Matemática  
and

<sup>2</sup>Posgraduação Instituto de Matemática  
Universidade de Rio de Janeiro  
Av. Athos da Silveira Ramos 149  
Edifício do centro de Tecnologia, Bloco C  
Caixa Postal 68530 - Ilha do Fundo - CEP 21941-909  
Rio de Janeiro - RJ - Brazil  
boattostefanella@gmail.com

<sup>3</sup>Department of Applied Mathematics  
University of St Andrews  
North Haugh  
St Andrews, KY 16 9SS, Scotland, UK  
dgd@mcs.st-and.ac.uk

<sup>4</sup>Dept. de Matematica Aplicada i Analisi  
Universitat de Barcelona  
Gran Via de les Cortes Catalanas 585,  
Barcelona 08007, Spain  
carles@maia.ub.es

## Abstract

Many key features of the motion of satellites, planets, stars, even galaxies can be captured by point mass dynamics. Likewise, many key features of fluid motion such as atmospheric storms, ocean eddies, super fluid vortices, and early stages of mass aggregation in gravitational systems can be captured by point vortex dynamics. Yet, serious mathematical challenges remain. Systems consisting of more than a few points are non-integrable, and complexity increases dramatically with the number of particles. Furthermore, the underlying geometry has a profound influence on the particle motion, as has only just begun to be investigated. Indeed one of today's challenges is a formulation of the  $N$ -vortex dynamics on Riemann surfaces. There are formulations over surfaces with constant Gaussian curvature [1, 4, 5, 6, 9, 10], and lately, for surfaces with not constant Gaussian curvature, conform to the sphere [2, 7]. We present some results about vortex dynamics on surfaces of revolution (among others the sphere and the ellipsoid of rotation) and we show the importance of the curvature in the stability of the relative equilibria

[1, 2]. The stabilization by curvature appears also in the classical  $N$ -body problem [5, 8]

#### REFERENCES

- [1] S. Boatto , Curvature perturbations and stability of a ring of vortices. *Discrete Contin. Dyn. Syst. Ser. B*, 10, no. 2-3, 349375 (2008).
- [2] D. Dritschel and S. Boatto, Vortex dynamics on surface of revolutions, in preparation.
- [3] S. Boatto, R. Schaefer, D. Dritschel, The  $N$ -body, the  $N$ -vortices and the  $N$ -charges dynamics on surfaces : a common view point, in preparation
- [4] S. Boatto and C. Simó, Thomson's Heptagon : A case of bifurcation at infinity, *Physica D*, **237**, 2051-2055 (2008)
- [5] S. Boatto and C. Simó, Existence and stability of rings of vortices on a sphere, in preparation.
- [6] Y. Kimura , Vortex motion pn surfaces with constant curvature. *R. Soc. Lond. Proc. Ser. A Math Phys. Eng. Sci.* (1999), vol. 455, pp 245-259.
- [7] J. Koiller and S. Boatto, Vortices on closed surfaces, preprint (2009)
- [8] R. Martínez, C. Simó, On the stability of the Lagrangian homographic solutions in a curved three-body problem on  $S^2$ , *Discrete and Continuous Dynamical Systems A*, 33, (2013), 1157–1175.
- [9] S. Pekarsky, and J.E. Marsden , Point-vortices on a sphere: Stability of relative equilibria. *J. Math. Phys.* 39, 58945907 (1998).
- [10] L.M. Polvani and D.G. Dritschel, Wave and vortex dynamics on the surface of the sphere. *J. Fluid Mech.*, Vol. 255, pp 35-64 (1993).

# STABILITY RESULTS OF SOME MODELS IN CELESTIAL MECHANICS

Alessandra Celletti<sup>1</sup>

<sup>1</sup>Department of Mathematics  
University of Roma Tor Vergata  
Via della Ricerca Scientifica, 1  
00133 - Roma (Italy)  
celletti@mat.uniroma2.it

## Abstract

I will present some stability results based on KAM theory. I will consider both conservative and dissipative (i.e., conformally symplectic) systems (either maps and flows). The proof is constructive and it provides efficient algorithms to evaluate the breakdown threshold of invariant tori ([2]). For low-dimensional degrees of freedom, KAM theory allows to get stability results. In this context, I will discuss the stability of the restricted 3-body problem (with particular reference to the asteroid 12 Victoria) and the stability of the Moon–Earth synchronous resonance under tidal effects. I will show that computer-assisted proofs may be devised in order to show the existence of invariant tori for realistic values of the parameters ([1], [3], [4]).

## REFERENCES

- [1] R. Calleja and A. Celletti, Breakdown of invariant attractors for the dissipative standard map, *CHAOS*, **20**, issue 1, 013121 (2010).
- [2] R. Calleja, A. Celletti and R. de la Llave, *A KAM theory for conformally symplectic systems: efficient algorithms and their validation*, Preprint (2012).
- [3] A. Celletti and L. Chierchia, KAM Stability and Celestial Mechanics, *Memoirs Amer. Math. Soc.* **187**, 878 (2007).
- [4] A. Celletti and L. Chierchia, Quasi-periodic attractors in Celestial Mechanics, *Archive for Rational Mechanics and Analysis* **191**, 2, 311–345 (2009).

# PERIODIC BRAKE ORBITS IN THE PLANAR ISOSCELES THREE-BODY PROBLEM

**Nai-Chia Chen**

School of Mathematics  
University of Minnesota  
Minneapolis, Minnesota, USA  
chen1945@umn.edu

## Abstract

The isosceles three body problem is a special case of the three body problem whose triangular configurations describe an isosceles triangle at every instant. A brake orbit is an orbit that starts with zero initial velocity. The purpose of this talk is to find periodic brake orbits in the isosceles three-body problem. We use various topological shooting arguments; we follow a curve of brake initial conditions under the flow until it reaches a suitable surface, and show that the image curve includes a point that corresponds to a periodic orbit. As a result, we find six types of periodic brake orbits.

## REFERENCES

- [1] N.-C. Chen, Periodic Brake Orbits in the Planar Isosceles Three-Body Problem, *submitted* (2013).
- [2] R. Devaney, Triple Collision in the Planar Isosceles Three Body Problem, *Invent. Math.* **60**, 3 (1980), 249-267.
- [3] R. Moeckel, R. Montgomery, and A. Venturelli, From brake to syzygy, *Arch. Ration. Mech.* **204**, 3 (2012), 1009-1060.
- [4] C. Simó and R. Martinez, Qualitative study of the planar isosceles three-body problem, *Celest. Mech. Dyn. Astr.* **41**, 1-4 (1987), 179-251.

# CLOSED TRIPLE COLLISION ORBITS IN PHOTOIONIZATION OF THE HELIUM ATOM

Nark Nyul Choi<sup>1</sup>, Min-Ho Lee<sup>1</sup>, Chang Woo Byun<sup>1</sup> and Gregor Tanner<sup>2</sup>

<sup>1</sup>School of Liberal Arts and Teacher Training, Kumoh National Institute of Technology  
Gumi, Gyeongbuk 730-701, South Korea

and

<sup>2</sup>School of Mathematical Sciences, University of Nottingham  
University Park, Nottingham NG7 2RD, UK  
nnchoi2001@gmail.com, minho.lee.kr@gmail.com, cwbyun@gmail.com and  
gregor.tanner@nottingham.ac.uk

## Abstract

The cross section for single-electron photoionization in the helium atom shows fluctuations which decrease in amplitude as the total energy of the atom approaches the double-ionization threshold [1]. Treating the helium atom as a three-body Coulomb problem, we predicted that the decay of the fluctuations can be characterized in terms of a threshold law with an exponent obtained as a combination of Siegel's stability exponents of the triple-collision singularity; the details of the fluctuations were predicted to be linked to a set of infinitely unstable classical orbits starting and ending at the nonregularizable triple collision - so called CTCOs (Closed Triple Collision Orbits) [2]. The validity of the prediction was confirmed by numerical calculations for the *collinear* helium model [3], which was presented at the workshop 'Few-Body Dynamics in Atoms, Molecules, and Planetary Systems' held in 2010. We extend our calculational method to planar helium and analyze the results to confirm the validity of the prediction for the *planar* helium model atom. In addition, it is shown that the CTCOs play the same role in the partial cross sections as well as in the total cross section.

## REFERENCES

- [1] R. Püttner, B. Grémaude, D. Delande, M. Domke, M. Martins, A.S. Schlachter, and G. Kaindl, Statistical Properties of Inter-Series Mixing in Helium: From Integrability to Chaos, *Phys. Rev. Lett.* **86**, 17 (2001), 3747–3750.
- [2] C. W. Byun, N. N. Choi, M.-H. Lee, and G. Tanner, Scaling Laws for the Photoionization Cross Section of Two-Electron Atoms, *Phys. Rev. Lett.* **98**, 11 (2007), 113001.
- [3] M.-H. Lee, C. W. Byun, N. N. Choi, and G. Tanner, Photoionization of two-electron atoms via highly doubly excited states: Numerical and semiclassical results, *Phys. Rev. A.* **81**, 4 (2010), 043419.

# GENERALIZED HIP–HOP SOLUTIONS OF THE $(N + N)$ –BODY PROBLEM

**Josep M. Cors**

Departament de Matemàtica Aplicada III  
Escola Politècnica Superior d'Enginyeria de Manresa  
Universitat Politècnica de Catalunya  
Avda. Bases de Manresa, 61-73. 08242 Manresa  
Catalonia  
cors@epsem.upc.edu

## Abstract

Hip–hop solutions of the equal–mass  $2N$ –body problem are periodic solutions in which the bodies lie on the vertices of a regular antiprism for all time. A regular antiprism is a polyhedron formed by two congruent regular  $N$ –gons lying on parallel planes and such that the orthogonal projection on each other is a regular  $2N$ –gon.

In [1] and [2] it was shown that there exist families of Hip–hop solutions in the neighbourhood of planar regular  $2N$ –gon relative equilibria solution and of planar highly eccentric elliptic homographic solutions, the planes containing each group of bodies performing small oscillations along the line joining the centres of the  $N$ –gons.

Consider now two regular  $N$ –gons, with masses equal within each group, lying on two planes perpendicular to the line joining their centres. It is easily seen that, if the initial velocities within each group are invariant under a  $2\pi/N$  rotation around that line, then the bodies remain on such a configuration for all time, provided no collision occurs. A periodic solution of this type with the same mean angular velocity for each group will be called a *Generalized Hip–Hop* solution of the  $(N + N)$ –body problem.

We show the existence of Generalized Hip–Hop solutions in the case of almost equal mass. These solutions are close to the planar regular  $2N$ –gon relative equilibria solutions with small vertical oscillations described above.

## REFERENCES

- [1] Esther Barrabés, Josep Maria Cors, Conxita Pinyol, and Jaume Soler. Hip–hop solutions of the  $2N$ –body problem. *Celestial Mechanics and Dynamical Astronomy*, 95(1-4):55–66, August 2006.
- [2] Esther Barrabés, Josep M. Cors, Conxita Pinyol, and Jaume Soler. Highly eccentric Hip-hop solutions of the  $2N$ –body problem. *Physica D: Nonlinear Phenomena*, 239(3-4):214–219, February 2010.

# RELATIVE EQUILIBRIA OF 3 BODIES IN DIMENSION $d$

Holger R. Dullin<sup>1</sup>

<sup>1</sup>School of Mathematics and Statistics  
The University of Sydney  
Sydney, NSW 2006, Australia  
holger.dullin@sydney.edu.au

## Abstract

The 3-body problem can be effectively symmetry reduced using invariants [1]. This leads to a Lie-Poisson structure and we analyse the equilibria of this 10-dimensional reduced system with two Casimirs. The Casimirs are the length of the angular momentum vector, and the Gram determinant of four difference vectors. When the spatial dimension  $d$  is equal to three then the Gram determinant is zero and the classical relative equilibria of Euler and Lagrange are found. However, when  $d > 3$  the Gram determinant may be non-zero, and this allows for additional families of relative equilibria, with shapes that are distinct from the collinear and the equilateral families. These additional relative equilibria were described by Albouy and Chenciner [2] who called them balanced configurations [3]. We will give a detailed description of such relative equilibria and their linear stability.

## REFERENCES

- [1] H.R. Dullin, The Lie-Poisson structure of the reduced n-body problem, arXiv:1207.5883
- [2] A. Albouy, A. Chenciner, Le Problème des N corps et les distances mutuelles, *Inventiones mathematicae* **131** (1998), 151-184.
- [3] A. Chenciner, The Lagrange reduction of the N-body problem, a survey, arXiv:1111.1334

# SCATTERING MONODROMY IN THE TWO CENTER PROBLEM

**Holger R. Dullin**<sup>1</sup> and **Holger Waalkens**<sup>2</sup>

<sup>1</sup> School of Mathematics and Statistics  
University of Sydney  
New South Wales 2006  
Australia

<sup>2</sup> Johann Bernoulli Institute for Mathematics and Computer Science  
University of Groningen  
9747 AG Groningen  
The Netherlands

Holger.Dullin@Sydney.edu.au and H.Waalkens@rug.nl

## Abstract

The Liouville-Arnold theorem states that the compact preimage of regular value of the energy momentum map of a Liouville integrable system is an  $n$ -dimensional torus where  $n$  is the number of degrees of freedom, and in the neighbourhood of this torus one can construct action-angle variables. Isolated singular values of the energy momentum map can cause a twist in the torus bundle over the regular image of an energy momentum map. As a result action-angle variables cannot be defined globally. This obstruction to the global construction of action-angle variables has been coined Hamiltonian monodromy, and many prominent examples of Liouville-integrable systems exhibiting this obstruction have been studied in recent years. Through the Bohr-Sommerfeld quantization of action variables monodromy carries over to quantum mechanics where it causes an obstruction to the global existence of quantum numbers. In this talk we discuss the classical and quantum mechanical implications of monodromy for scattering systems, i.e. for Liouville integrable systems where the pre images of the energy momentum map are not compact. As examples we study planar scattering at a repulsive central potential [1] and Euler's problem of two fixed centers [2, 3].

## REFERENCES

- [1] H. R. Dullin and H. Waalkens, Nonuniqueness of the phase shift in central scattering due to monodromy, *Phys. Rev. Lett.* **101** (2008), 070405.
- [2] H. Waalkens, A. Junge and H. R. Dullin, Quantum monodromy in the two center-problem, *J. Phys. A* **36** (2003), L307-L314.
- [3] H. Waalkens, H. R. Dullin, and P. H. Richter, The problem of two fixed centers: bifurcations, actions, monodromy, *Physica D* **196** (2004) 265-310.

# PHASE SPACE STRUCTURE OF THE WANNIER RIDGE OF DRIVEN HELIUM

Markus Heinrich, Alberto Rodriguez and Andreas Buchleitner

Institute of Physics  
Department of Mathematics and Physics  
Albert-Ludwigs-Universität Freiburg  
Freiburg, Germany

## Abstract

The three-body physics of helium in a strong driving field has caught a lot of attention since the famous “helium knee” showed the importance of electronic correlation in the double ionisation process [1]. Nowadays, there still exist many open questions on the role of electron-electron interaction in non-sequential ionisation [2, 3]. We study the Wannier ridge sub-manifold of the classical helium phase space for vanishing total angular momentum (i.e. planar configurations). The Wannier ridge corresponds to a completely symmetric and therefore correlated motion of the electrons. We prove that this configuration is invariant in the presence of periodic driving, if and only if the field is linearly polarised along the symmetry axis. Thus, correlated field-induced double ionisation can occur through the Wannier ridge, at least at the classical level. We use a time-frequency representation of orbits given by wavelet analysis to study the 5-dimensional phase space. Although the field mostly destroys the regular dynamics, we observe that it is also able to stabilise some regions of phase space for certain field frequencies and intensities [4].

## REFERENCES

- [1] B. Walker *et al.*, Precision Measurement of Strong Field Double Ionization of Helium, *Phys. Rev. Lett.* **73**, 9 (1994), 1227–1230.
- [2] G. Tanner, K. Richter and J.-M. Rost, The theory of two-electron atoms: Between ground state and complete fragmentation, *Rev. Mod. Phys.*, **72** (2000), 497–544.
- [3] W. Becker *et al.*, Theories of photoelectron correlation in laser-driven multiple atomic ionization, *Rev. Mod. Phys.*, **84** (2012), 1011-1043.
- [4] M. Heinrich, Phase space structure and classical chaos on the Wannier ridge of driven helium, B.Sc. thesis, Universität Freiburg (2013).

# RECOLLIDING PERIODIC ORBITS IN ATTOSECOND PHYSICS

Adam Kamor<sup>1,2</sup>, Francois Mauger<sup>2</sup>, Cristel Chandre<sup>2</sup>, and Turgay Uzer<sup>1</sup>

<sup>1</sup>School of Physics, Georgia Institute of Technology, Atlanta, Ga. USA

and

<sup>2</sup> Centre de Physique Théorique - UMR 7332, CNRS – Aix-Marseille University, Marseille, France

turgay.uzer@physics.gatech.edu

## Abstract

Can an ionized electron be driven back to the core by an ultrastrong laser pulse? This is a high-stakes issue in attosecond physics since the returning electron, by carrying back the energy it has absorbed from the laser, can act as the agent of many key processes in intense laser physics, including the ultrafast imaging of macromolecules and the design of new light sources through generation of ultra-high harmonics [1, 2]. When the laser is linearly polarized, the “recollision” (or “rescattering”) model [3, 4] has been immensely successful in both interpreting current experiments and devising new ones. However the picture is much less clear for all other polarizations [5, 6]. The very same recollision model predicts that returns must be suppressed in circularly polarized laser pulses because ionized electrons tend to spiral away from the core. In contrast, we present evidence that electrons return to the core along specific families of periodic orbits which shuttle electrons towards and away from the core. We show that these special periodic orbits are the key agents of attosecond processes in circularly polarized fields.

## REFERENCES

- [1] P.B. Corkum, Recollision Physics, *Physics Today* **64**, (2011).
- [2] W. Becker and X. Liu and P.J. Ho and J.H. Eberly, Theories of photoelectron correlation in laser-driven multiple atomic ionization, *Rev. Mod. Phys.* **84**, (2012).
- [3] P.B. Corkum, Plasma perspective on strong field multi-photon ionization, *Phys. Rev. Lett.* **71**, (1993).
- [4] K.J. Schafer and B. Yang and L.F. DiMauro and K.C. Kulander, Above threshold ionization beyond the high harmonic cutoff, *Phys. Rev. Lett* **70**, (1993).
- [5] F. Mauger and C. Chandre and T. Uzer, Recollision and correlated double ionization with circularly polarized light, *Phys. Rev. Lett.* **105**, (2010).
- [6] X. Wang and J.H. Eberly, Elliptical trajectories in non-sequential double ionization., *New. J. Phys.* **12**, (2010).

# TRAIN CORRESPONDENCES FOR CELESTIAL BODIES

Andreas Knauf<sup>1</sup>

<sup>1</sup>Department of Mathematics, Friedrich-Alexander-University Erlangen-Nuremberg,  
Cauerstr. 11, D-91058 Erlangen, Germany, [knauf@mi.uni-erlangen.de](mailto:knauf@mi.uni-erlangen.de)

## Abstract

Joint with Richard Montgomery (UC Santa Cruz); to appear.

By kinematics of scattering we mean motion of  $n$  particles on line segments of  $\mathbb{R}^d$ , with two-body collisions at the end points, preserving momentum and energy. It is known (see [1]) that the number of collisions is finite.

We analyze the kinematics of scattering. Then, adapting techniques from [2], we show that every transversal kinematical scattering process can be approximated by the dynamics of particles with Coulomb interaction (attractive or repulsive), in the limit of large energy or large spatial extensiveness.

## REFERENCES

- [1] D. Burago, S. Ferleger, S. Kononenko: Uniform estimates on the number of collisions in semi-dispersing billiards. *Annals of Mathematics* **147**, 695–708 (1998)
- [2] A. Knauf: The  $n$ -centre problem of celestial mechanics for large energies. *Journal of the European Mathematical Society* **4**, 1–114 (2002)

# INVARIANT TORI IN THE SPATIAL 3-BODY PROBLEM BY AVERAGING AND REDUCTION

**Jesús Palacián, Flora Sayas and Patricia Yanguas**

Departamento de Ingeniería Matemática e Informática  
Universidad Pública de Navarra  
Campus de Arrosadía, 31006 Pamplona (Navarra) Spain  
palacian@unavarra.es, flora.sayas@unavarra.es and yanguas@unavarra.es

## Abstract

We study the spatial three-body problem, which is a system of nine degrees of freedom. In order to apply perturbation theory we need to establish the possible regimes where the Hamiltonian of the three-body problem can be split into two parts: the unperturbed Hamiltonian composed of two Keplerian terms and the perturbation, which is supposed to be small with respect to the principal part.

Our approach is based on a combination of averaging techniques with reduction theory with the aim of building a reduced Hamiltonian and a reduced phase space as simple as possible. The reduction process takes into account all possible continuous symmetries of the problem, including the symmetry generated by the two approximate integrals obtained after performing the normalisation with respect to the two fast angles and truncating the higher-order terms.

Once the reduction process is completed, the averaged Hamiltonian defines a system of one degree of freedom in the fully reduced space whose dimension is two and which is embedded in  $\mathbb{R}^3$ . This phase space is a surface that depends on three parameters, it is parametrised using Deprit's elements [1] and may have zero, one, two or three singular points. Our approach is global in the sense that we deal with the flow of the fully-reduced system in the whole fully-reduced space.

The next step is the discussion of the occurrence of the different relative equilibria of the reduced Hamiltonian system. This is done in terms of the invariants and the fundamental constraint that define the fully-reduced phase space. There are two basic parameters to perform the analysis, namely the modulus of the total angular momentum vector and the modulus of the angular momentum vector of the outer ellipse. They generate the plane of parameters which is divided into six different regions and presents five bifurcation lines. Each region has a different number of relative equilibria, ranging from two to six. The number and stability of the equilibria change when crossing the different lines [3].

We consider every elliptic point in the reduced phase space and reconstruct it step by step to compute the corresponding KAM invariant tori that persist in the original space  $\mathbb{R}^{12}$ . In the cases where the equilibria are placed on the singular points of the two-dimensional reduced phase space we need to make a local analysis around the singularities, passing to a surface with no singular points.

In the process of reconstruction we define intermediate spaces and sets of variables depending on the type of motions under consideration. Due to the scale in the perturbation, the Hamiltonian is too degenerate to apply classical results of KAM theory. Thus, we use a specific result by Han, Li and Yi [2] for this kind of systems.

We find Cantor families of five-dimensional KAM tori for the spatial three-body problem.

#### REFERENCES

- [1] A. Deprit, Elimination of the nodes in problems of  $n$  bodies, *Celest. Mech.* **30**, 181-195 (1983).
- [2] Y. Han, Y. Li and Y. Yi, Invariant tori in Hamiltonian systems with high order proper degeneracy, *Ann. Henri Poincaré* **10**, 1419-1436 (2010).
- [3] J.F. Palacián, F. Sayas and P. Yanguas, Regular and singular reductions in the spatial three-body problem, to appear in *Qual. Theory Dyn. Syst.* (2013).

# QUANTUM AND CLASSICAL MOTION OF TWO ELECTRONS IN ATOMS AND LASER FIELDS

**Thomas Pfeifer**

Max-Planck Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany  
Center for Quantum Dynamics, Ruprecht-Karls-Universität Heidelberg, 69120 Heidelberg,  
Germany

## Abstract

In atomic physics, the few-body problem arises already for the second element in the periodic table, helium (He), exhibiting two electrons "orbiting" inside the  $1/r$  Coulomb potential around the doubly-charged nucleus. Any atom or ion besides hydrogen (H) or H-like systems is thus in general not analytically solved and theoretical description relies on approximation or numerics that need to be tested.

Here, experiments are presented on measurements and the control of two-electron dynamics in atoms by applying laser/electric fields, on time scales of femtoseconds to attoseconds. Quantum-mechanical correlated wave packets of two excited electrons in helium can be observed and controlled [1]. In stronger laser fields, quasi-classical recollision occurs, where a single electron is freed, accelerated in the laser field and returns to its parent ion where it can further excite other electrons. Experiments show that a two-electron (doubly-) excited state can also be formed in this process [2]. Comparing measurements with classical trajectory simulations, we find that two electrons are then removed from the atom sequentially, but within a fraction of a laser-optical cycle ( $\sim 200$  as) even when initially bound to a fully symmetric orbit.

## REFERENCES

- [1] C. Ott *et al.*, Quantum Interferometry and Correlated Two-Electron Wave-Packet Observation in Helium, arXiv:1205.0519.
- [2] N. Camus *et al.*, Attosecond Correlated Dynamics of Two Electrons Passing through a Transition State, *Phys. Rev. Lett.* **108** (2012), 073003.

# BIFURCATION ANALYSIS OF SYMMETRIC HAMILTONIAN RESONANCES

Giuseppe Pucacco<sup>1,2</sup>

<sup>1</sup>Dipartimento di Fisica

and

<sup>2</sup>INFN – Sezione di Roma II

Università di Roma “Tor Vergata”

Via della Ricerca Scientifica, 1 - 00133 Roma

pucacco@roma2.infn.it

## Abstract

We present a general analysis of the bifurcation sequences of periodic orbits in general position of resonant Hamiltonian normal forms invariant under  $\mathbb{Z}_2 \times \mathbb{Z}_2$  symmetry. The low-order cases 2:2 and 2:4 show rich structure [1, 2, 3] investigated both with geometric methods and a singularity theory approach.

## REFERENCES

- [1] Marchesiello, A. Pucacco, G. 2011, *Relevance of the 1:1 resonance in galactic dynamics*, European Physical Journal Plus, **126**, 104.
- [2] Marchesiello, A. & Pucacco, G. 2013, *The symmetric 1:2 resonance*, European Physical Journal Plus, **128**, 21.
- [3] Marchesiello, A. Pucacco, G. 2013, *Resonances and bifurcations in systems with elliptical equipotentials*, Monthly Notices of the Royal Astronomical Society, **428**, 2029.

# STABILITY OF RELATIVE EQUILIBRIA IN THE PLANAR N-VORTEX PROBLEM: A TOPOLOGICAL APPROACH

**Gareth E. Roberts**

Department of Mathematics and Computer Science  
College of the Holy Cross  
1 College Street  
Worcester, MA 01604, USA  
groberts@holycross.edu

## Abstract

In the weather research and forecasting models of certain hurricanes [1], vortex crystals are found within a polygonal-shaped eyewall. These special configurations can be interpreted as relative equilibria (rigidly rotating solutions) of the point vortex problem introduced by Helmholtz. Their stability is thus of considerable importance. Adapting the approach of Moeckel [2] for the companion problem in celestial mechanics, we present some theory and results on the linear and nonlinear stability of relative equilibria in the planar N-vortex problem [3]. A topological approach is taken to show that for the case of positive circulations, a relative equilibrium is linearly stable if and only if it is a nondegenerate minimum of the Hamiltonian restricted to a level surface of the angular impulse (moment of inertia). Using a criterion of Dirichlet's, this implies that any linearly stable relative equilibrium with positive vorticities is also nonlinearly stable. Two symmetric families, the rhombus and the isosceles trapezoid, are analyzed, with stable solutions found in each case.

## REFERENCES

- [1] Corbosiero, K., Advanced Research WRF High Resolution Simulations of the Inner Core Structure of Hurricanes Katrina, Rita and Wilma (2005), <http://www.atmos.albany.edu/facstaff/kristen/wrf/wrf.html>.
- [2] Moeckel, R., Linear stability analysis of some symmetrical classes of relative equilibria, *Hamiltonian dynamical systems (Cincinnati, OH, 1992)*, IMA Vol. Math. Appl., 63, (1995), Springer, 291–317.
- [3] Roberts, G. E., Stability of Relative Equilibria in the Planar N-Vortex Problem, arXiv:1301.6194, *SIAM Journal on Applied Dynamical Systems* (to appear).

# REGULARIZATION OF THE KEPLER PROBLEM ON THE SPHERE

**Manuele Santoprete<sup>1</sup>**

<sup>1</sup>Department of Mathematics  
Wilfrid Laurier University  
75 University Avenue West  
Waterloo, ON  
msantopr@wlu.ca

## Abstract

In this talk I will present some results concerning the Kepler problem on the three-sphere  $S^3$ , that I obtained in collaboration with Shengda Hu [1]. I will show how to perform a Moser-type regularization and how to adapt the Ligon-Schaaf regularization to the problem under consideration. Then, I will explain the relationship between these regularizations and the corresponding regularizations for the Kepler problem in  $\mathbb{R}^3$ . This will be done by showing that the Moser regularization and the Ligon-Schaaf map we obtained can be understood as the composition of the corresponding maps for the Kepler problem in Euclidean space, and the gnomonic transformation.

## REFERENCES

- [1] S. Hu, M. Santoprete, Regularization of the Kepler problem on the Sphere, *Canadian Journal of Mathematics*, <http://dx.doi.org/10.4153/CJM-2012-039-9>, (to appear in print)

# NORMAL FORMS FOR ROTATIONALLY-INVARIANT SYSTEMS

**Cristina Stoica**

Department of Mathematics and Statistics  
Wilfrid Laurier University  
Waterloo, ON, Canada, N2L 3C5  
cstoica@wlu.ca

## Abstract

Rotationally invariant cotangent-bundle systems and the particular the case of  $n$ -mass point systems are studied using a recently-obtained parametrisation of the phase space. This parametrisation, in which the Marsden-Weinstein symplectic reduced space becomes a vector space with a canonical symplectic form, is beneficial in the study of dynamics near relative equilibria. In particular, it can be shown that Nekhoroshev's estimates near an elliptic equilibrium point in the reduced space induce similar estimates in the full phase-space.

## REFERENCES

- [1] V. I. Arnol'd, *Mathematical Methods in Classical Mechanics*, Springer, 1989.
- [2] U. Çiftçi and H. Waalkens, *Phase space structures governing reaction dynamics in rotating molecules*, *Nonlinearity* 25, 2012.
- [3] J. Montaldi and M. Roberts, *Relative Equilibria of Molecules*, *J. Nonlinear Science* 9, 1999.
- [4] J.E. Marsden, *Lectures in Mechanics*, Cambridge Univ. Press, 1992.
- [5] T. Schmah and C. Stoica, *Normal forms for  $SO(3)$ -invariant cotangent bundle systems*, in *Geometry, Mechanics and Dynamics: the Legacy of Jerry Marsden* volume, Fields Institute Communications Series, Springer (to appear).

# INVARIANT TORI IN THE SPATIAL 3-BODY PROBLEM BY AVERAGING AND REDUCTION

**Jesús Palacián, Flora Sayas and Patricia Yanguas**

Departamento de Ingeniería Matemática e Informática  
Universidad Pública de Navarra  
Campus de Arrosadía, 31006 Pamplona (Navarra) Spain  
palacian@unavarra.es, flora.sayas@unavarra.es and yanguas@unavarra.es

## **Abstract**

We study the spatial three-body problem, which is a system of nine degrees of freedom. In order to apply perturbation theory we need to establish the possible regimes where the Hamiltonian of the three-body problem can be split into two parts: the unperturbed Hamiltonian composed of two Keplerian terms and the perturbation, which is supposed to be small with respect to the principal part.

Our approach is based on a combination of averaging techniques with reduction theory with the aim of building a reduced Hamiltonian and a reduced phase space as simple as possible. The reduction process takes into account all possible continuous symmetries of the problem, including the symmetry generated by the two approximate integrals obtained after performing the normalisation with respect to the two fast angles and truncating the higher-order terms.

Once the reduction process is completed, the averaged Hamiltonian defines a system of one degree of freedom in the fully reduced space whose dimension is two and which is embedded in  $\mathbb{R}^3$ . This phase space is a surface that depends on three parameters, it is parametrised using Deprit's elements [1] and may have zero, one, two or three singular points. Our approach is global in the sense that we deal with the flow of the fully-reduced system in the whole fully-reduced space.

The next step is the discussion of the occurrence of the different relative equilibria of the reduced Hamiltonian system. This is done in terms of the invariants and the fundamental constraint that define the fully-reduced phase space. There are two basic parameters to perform the analysis, namely the modulus of the total angular momentum vector and the modulus of the angular momentum vector of the outer ellipse. They generate the plane of parameters which is divided into six different regions and presents five bifurcation lines. Each region has a different number of relative equilibria, ranging from two to six. The number and stability of the equilibria change when crossing the different lines [3].

We consider every elliptic point in the reduced phase space and reconstruct it step by step to compute the corresponding KAM invariant tori that persist in the original space  $\mathbb{R}^{12}$ . In the cases where the equilibria are placed on the singular points of the two-dimensional reduced phase space we need to make a local analysis around the singularities, passing to a surface with no singular points.

In the process of reconstruction we define intermediate spaces and sets of variables depending on the type of motions under consideration. Due to the scale in the perturbation, the Hamiltonian is too degenerate to apply classical results of KAM theory. Thus, we use a specific result by Han, Li and Yi [2] for this kind of systems.

We find Cantor families of five-dimensional KAM tori for the spatial three-body problem.

#### REFERENCES

- [1] A. Deprit, Elimination of the nodes in problems of  $n$  bodies, *Celest. Mech.* **30**, 181-195 (1983).
- [2] Y. Han, Y. Li and Y. Yi, Invariant tori in Hamiltonian systems with high order proper degeneracy, *Ann. Henri Poincaré* **10**, 1419-1436 (2010).
- [3] J.F. Palacián, F. Sayas and P. Yanguas, Regular and singular reductions in the spatial three-body problem, to appear in *Qual. Theory Dyn. Syst.* (2013).

# RECOLLIDING PERIODIC ORBITS IN ATTOSECOND PHYSICS

Adam Kamor<sup>1,2</sup>, Francois Mauger<sup>2</sup>, Cristel Chandre<sup>2</sup>, and Turgay Uzer<sup>1</sup>

<sup>1</sup>School of Physics, Georgia Institute of Technology, Atlanta, Ga. USA

and

<sup>2</sup> Centre de Physique Théorique - UMR 7332, CNRS – Aix-Marseille University, Marseille, France

turgay.uzer@physics.gatech.edu

## Abstract

Can an ionized electron be driven back to the core by an ultrastrong laser pulse? This is a high-stakes issue in attosecond physics since the returning electron, by carrying back the energy it has absorbed from the laser, can act as the agent of many key processes in intense laser physics, including the ultrafast imaging of macromolecules and the design of new light sources through generation of ultra-high harmonics [1, 2]. When the laser is linearly polarized, the “recollision” (or “rescattering”) model [3, 4] has been immensely successful in both interpreting current experiments and devising new ones. However the picture is much less clear for all other polarizations [5, 6]. The very same recollision model predicts that returns must be suppressed in circularly polarized laser pulses because ionized electrons tend to spiral away from the core. In contrast, we present evidence that electrons return to the core along specific families of periodic orbits which shuttle electrons towards and away from the core. We show that these special periodic orbits are the key agents of attosecond processes in circularly polarized fields.

## REFERENCES

- [1] P.B. Corkum, Recollision Physics, *Physics Today* **64**, (2011).
- [2] W. Becker and X. Liu and P.J. Ho and J.H. Eberly, Theories of photoelectron correlation in laser-driven multiple atomic ionization, *Rev. Mod. Phys.* **84**, (2012).
- [3] P.B. Corkum, Plasma perspective on strong field multi-photon ionization, *Phys. Rev. Lett.* **71**, (1993).
- [4] K.J. Schafer and B. Yang and L.F. DiMauro and K.C. Kulander, Above threshold ionization beyond the high harmonic cutoff, *Phys. Rev. Lett* **70**, (1993).
- [5] F. Mauger and C. Chandre and T. Uzer, Recollision and correlated double ionization with circularly polarized light, *Phys. Rev. Lett.* **105**, (2010).
- [6] X. Wang and J.H. Eberly, Elliptical trajectories in non-sequential double ionization., *New. J. Phys.* **12**, (2010).

# A SIMPLE METHOD TO CONSTRUCT LOCAL EQUILIBRIUM FUNCTION

P. WANG<sup>1</sup> and S. Q. ZHANG<sup>1</sup>

<sup>1</sup>Yangtze Center of Mathematics and College of Mathematics Sichuan University  
Chengdu 610064, People's Republic of China.  
7790326@qq.com and zhangshiqing@msn.com

## Abstract

We have developed a simple method to construct local equilibrium function for lattice Boltzmann method (LBM). This new method can make LBM model satisfy compressible flow with a flexible specific-heat ratio. Test cases, including the one dimensional Sod flow, one dimensional Lax flow and thermal Couette flow are presented. Good results obtained using proposed new method, indicate that the proposed method is potentially capable of constructing of the local equilibrium function for LBM.

## REFERENCES

- [1] F. J. Alexander, S. Chen and J. D. Sterling, *Phys. Rev. E* **47**, 2249 (1993).
- [2] Y. Chen, H. Ohashi and M. Akiyama, *Phys. Rev. E* **50**, 2776 (1994).
- [3] M. Watari and M. Tsutahara, *Phys. Rev. E* **67**, 036306 (2003).
- [4] K. Qu, C. Shu, and Y. T. Chew, *Phys. Rev. E* **75**, 036706 (2007).
- [5] Q. Li, Y. L. He and Y. J. Gao, *Int. J. Mod. Phys. C* **19**, 1581 (2008).
- [6] H. X. Zhang, *Acta Aerodyn. Sin.* **6**, 143 (1988).
- [7] C. Cercignani, (Springer-Verlag, Berlin, 1988).
- [8] P. L. Bhatnagar, E. P. Gross and M. Krook, *Phys. Rev.* **94**, 511 (1954).
- [9] C. H. Sun, *Phys. Rev. E* **58**, 7283 (1998).
- [10] Y.L. He, Y. Wang and Q. Li, (Science Publishing House, Beijing, 2008).
- [11] Z. L. Guo and T. S. Zhao, *Phys. Rev. E* **67**, 066709 (2003).
- [12] Y. Wang, Y. L. He, T. S. Zhao, G. H. Tang and W. Q. Tao, *Int. J. Mod. Phys. C* **18**, 1961 (2007).
- [13] C. H. Sun, *Phys. Rev. E* **61**, 2645 (2000).
- [14] Z. L. Guo, C. G. Zheng and B. C. Shi *Chinese Phys.* **11**, 366 (2002).

# THE RELATIONSHIPS BETWEEN REGULAR POLYGONAL CENTRAL CONFIGURATIONS AND MASSES FOR NEWTONIAN N-BODY PROBLEMS

Wang Zhiqiang

Department of Mathematics  
Sichuan University  
Chengdu, China  
wangzhiqiang0213@gmail.com

## Abstract

This is a joint work with Li Wei.

In 1985, Perko and Walter ([5]) proved that the  $N(N \geq 4)$  point masses that locate at the vertices of a regular  $N$ -polygon form a central configuration if and only if all the masses are equal. In this talk, we will give a simpler proof of their theorem and show that if the  $N$  point masses locate at the vertices of a regular  $N+1$ -polygon, they can't form a central configuration for any masses.

## REFERENCES

- [1] A. Chenciner, *Are there perverse choreographies*, Proceedings of the HAMSYS conference, held in Guanajuato, March 2001
- [2] F. Diacu, *Polygonal homographic orbits of the curved  $n$ -body problem*, Trans. Amer. Math. Soc. 364 , 2783-2802, 2012
- [3] M. Marcus and H. Minc, *A survey of matrix theory and matrix inequalities*, Allyn and Bacon 1964, Reprinted by Dover, 1992
- [4] R. Moeckel, *On central configurations*, Math.Z. 205, 499-517, 1990
- [5] L.M. Perko and E.L. Walter, *Regular polygon solutions of the  $n$ -body problem*, Proceedings of the AMS, 1985
- [6] D. Saari, *Central configurations—A problem for the 21st Century*, in <http://math.uci.edu/~dsaari/BAMA-pap.pdf>
- [7] A. Wintner, *The analytical foundations of celestial mechanics*, Princeton Math. Vol. 5, Princeton University Press, Princeton, 1947
- [8] Z.F. Xie and S.Q. Zhang, *A simpler proof of regular polygon solution of  $N$ -body problem*, Physics Letters A, vol.277, 156-158, 2000

# LONG-LIVED ORBITS WITH NON-VANISHING ANGULAR MOMENTUM IN DOUBLY-EXCITED PLANAR HELIUM

**Klaus Zimmermann<sup>1,5</sup> and Pierre Lugan<sup>2</sup> and Celsus Bouri<sup>3</sup> and Javier  
Madronero<sup>4</sup> and Andreas Buchleitner<sup>1</sup>**

<sup>1</sup> Physikalisches Institut, Albert-Ludwigs-Universität Freiburg,  
Hermann-Herder-Str. 3, D-79104 Freiburg, Germany

<sup>2</sup> Laboratory of Theoretical Physics of Nanosystems,  
École Polytechnique Fédérale de Lausanne

PH H2 425 (Bâtiment PH), Station 3, CH-1015 Lausanne, Swiss

<sup>3</sup> Centre of Atomic Molecular Physics and Quantum Optics (CEPAMOQ), Faculty of Science,  
University of Douala, Douala, P.O. Box 8580, Cameroon

<sup>4</sup>Physik Department, Technische Universität München, D-85747 Garching, Germany

<sup>5</sup> klaus.zimmermann@physik.uni-freiburg.de

## Abstract

Helium as a microscopic embodiment of the 3-body problem is known to be chaotic. Early studies ([1, 2]) have shown that there are (quasi-)stable submanifolds, which support, e.g., the frozen planet configuration. We present some new long-lived classical orbits that exhibit surprisingly long life times and unusual shapes. These orbits were inferred from a spectral analysis, including (auto-)ionization rates, of the quantum problem.

## REFERENCES

- [1] K. Richter and D. Wintgen, Phys. Rev. Lett. **65**, 1965 (1990).
- [2] P. Schlagheck and A. Buchleitner, Eur. Phys. J. D **22**, 401 (2003).