

PIMS DISTINGUISHED CHAIR LECTURE SERIES CRG-CLIMATE MODELLING

UNIVERSITY OF VICTORIA
SEOS/CCCMA SPECIAL TAO SEMINAR

Applied & Theoretical Challenges for Multi-Scale Hyperbolic PDE's in the Tropics

Distinguished Visitor:
Dr. Andrew Majda, NYU
Courant institute of Mathematical Sciences

Thursday,
February 14, 2008
3:30-4:30pm ECS 125

Geophysical flows are a rich source of novel problems for applied mathematics and the contemporary theory of partial differential equations. The reason for this is that many physically important geophysical flows involve complex nonlinear interaction over multi-scales in both time and space so developing simplified reduced models which are simpler yet capture key physical phenomena is of central importance. In mid-latitudes, the fact that the rotational Coriolis terms are bounded away from zero leads to a strict temporal frequency scale separation between slow potential vorticity dynamics and fast gravity waves: this physical fact leads to new theorems justifying the quasi-geostrophic midlatitude dynamics even with general unbalanced initial data for both rapidly rotating shallow water equations and completely stratified flows.

At the equator, the tangential projection of the Coriolis force from rotation vanishes identically so that there is no longer a time scale separation between potential vortical flows and gravity waves. This has profound consequences physically that allow the tropics to behave as a waveguide with extremely warm surface temperatures. The resulting behavior profoundly influences longer term mid-latitude weather prediction and climate change through hurricanes, monsoons, El Nino, and global teleconnections with the mid-latitude atmosphere. How this happens through detailed physical mechanisms is one of the most important contemporary problems in the atmosphere-ocean science community with a central role played by nonlinear interactive heating involving the interaction of clouds, moisture, and convection. The variable coefficient degeneracy of the Coriolis term at the equator alluded to earlier leads to both important new physical effects as well as fascinating new mathematical phenomena and PDE's. In this equatorial context, new multi-scale reduced dynamical PDE models are even relatively recent in origin.



Gaspard Gustave de Coriolis
1792-1843



www.pims.math.ca

