

Distance Spectrum of Graphs - Some Recent Studies

Ambat Vijayakumar

Department of Mathematics
Cochin University of Science and Technology
India.
(vambat@gmail.com)

Abstract

The distance matrix of a connected graph G is an $n \times n$ matrix $D(G) = [d_{ij}]$, where d_{ij} is the distance between the vertices v_i and v_j . The distance spectrum of G is $\{\lambda_1, \dots, \lambda_n\}$, where λ_i 's are the eigenvalues of $D(G)$. The distance energy of G is defined by $E_D(G) = \sum_{i=1}^n |\lambda_i|$. The distance matrix, distance eigenvalue, and distance energy of a connected graph have been studied intensively in literature, see [1, 7, 8, 9, 12]. We discuss a new problem of how the distance energy changes when an edge is deleted. Similar problem for adjacency energy of a graph was studied by Day and So in [3, 4]. It turns out that the results for distance energy change and adjacency energy change are quite different. From an observation in [12], it follows that, for any connected graph with a unique positive eigenvalue, the deletion of any edge increases the distance energy provided that the resulting graph is still connected.

We prove that the distance energy of a complete bipartite graph is always increased when an edge is deleted even though it has two positive distance eigenvalues. Also, we give a set of examples of connected graph whose distance energy decreases when a specific edge is deleted. This is a joint work with G. Indulal and A. Varghese.

References

- [1] M. Aouchiche and P. Hansen, Distance spectra of graphs: a survey, *Linear Algebra and its Applications* 458 (2014) 301–386.
- [2] R. Bapat, S. Kirkland and M. Neumann, On distance matrices and Laplacians, *Linear Algebra and its Applications* 401 (2005) 193–209.

- [3] J. Day and W. So, Singular value inequality and graph energy change, *Electron. J. Linear Algebra* 16 (2007) 291–299.
- [4] J. Day and W. So, Graph energy change due to edge deletion, *Linear Algebra and its Applications* 428 (2008) 2070–2078.
- [5] B. Durgi, P. Hampiholi, S. Mekkalike, Distance spectra and distance energy of some cluster graphs, *Mathematica Aeterna* 4 (2014) 817 – 825.
- [6] R. Horn and C. Johnson, *Matrix Analysis*, Cambridge University press, 1985.
- [7] A. Ilic, Distance spectra and distance energy of integral circulant graphs, *Linear Algebra and its Applications* 433 (2010) 1005–1014.
- [8] G. Indulal, Distance spectrum of graph compositions, *Ars Mathematica Contemporanea* 2 (2009) 93–100.
- [9] G. Indulal, I. Gutman and A. Vijayakumar, On distance energy of graphs, *MATCH Commun. Math. Comput. Chem.* 60 (2008) 461–472.
- [10] W. So, A shorter proof of the distance energy of complete multipartite graphs, *Special Matrices* 5 (2017) 61–63.
- [11] D. Stevanovic and G. Indulal, The distance spectrum and energy of the composition of regular graphs, *Appl. Math. Lett.* 22 (2009) 1136–1140.
- [12] B. Zhou and A. Ilic, On distance spectral radius and distance energy of graphs, *MATCH Commun. Math. Comput. Chem.* 64 (2010) 261–280.
- [13] A. Varghese, W. So, A. Vijayakumar, Distance energy change of Complete-bipartite graph due to edge deletion, *Linear Algebra and Applications* 553 (2018) 211–222.