

# A DNA Copy Number Co-deletion Network Identifies Genetic Interactions

Xianlong Wang<sup>\*,1</sup>, Dongqing Huang<sup>†,1</sup>, Chenwei Lin<sup>1</sup>, Brian D. Piening<sup>1</sup>, Amanda G. Paulovich<sup>1</sup>, Pei Wang<sup>1</sup>

<sup>1</sup>Fred Hutchinson Cancer Research Center, Seattle, WA, USA

\*Correspondence: contact email, [xwan2@fhcrc.org](mailto:xwan2@fhcrc.org)

†Correspondence: contact email, [dhuang@fhcrc.org](mailto:dhuang@fhcrc.org)

## Abstract

The complex interplay among multiple genes in breast cancer etiology remains an important yet largely unexplored phenomenon. Therefore, understanding gene-gene interactions will shed new light on the genetic architecture of breast cancer. Since genome stability, partially reflected by copy number variation, plays a key role in breast cancer initiation and progression, copy number co-deletion events may help us reveal important disease-related genetic interactions. Such co-deletion events may be quantified on a genome-wide level using comparative genome hybridization (CGH) arrays. In this work, a bioinformatics pipeline was implemented to detect and prioritize disease-related genetic interactions using a copy number co-deletion network coupled with association scores between genomics signatures and patient clinical outcome. We applied the pipeline on a collection of genomics arrays of breast cancer and identified a list of candidate disease-related genetic interactions. As a proof of principle, we successfully validated a subset of candidate genetic interactions using yeast as a model organism from the perspectives of DNA repair and growth. Further experiments indicate that the growth defect may result from genome instability.

**Keywords:** gene-gene interaction; copy number co-deletion network; gene prioritization; integrative genomics analysis; data integration