

Semisimple Symplectic Characters of Finite Unitary Groups

Ryan Vinroot
joint with Bhama Srinivasan

Department of Mathematics
The College of William & Mary

June 4, 2012 / CMS Summer Meeting, Regina, Canada

Frobenius-Schur Indicators

- Let G be a finite group, and (π, V) an irreducible complex representation of G with character χ .
- The *Frobenius-Schur indicator* of π , or of χ , is
$$\varepsilon(\chi) = \frac{1}{|G|} \sum_{g \in G} \chi(g^2).$$
- Then $\varepsilon(\chi) = 1, -1$, or 0 when π is an \mathbb{R} -representation, χ is real-valued but π is not an \mathbb{R} -representation, or χ is not real-valued, respectively.
- If $\varepsilon(\chi) = -1$, χ is *symplectic*. If $\varepsilon(\chi) = 1$, χ is *orthogonal*.

Frobenius-Schur Indicators

- Let G be a finite group, and (π, V) an irreducible complex representation of G with character χ .
- The *Frobenius-Schur indicator* of π , or of χ , is
$$\varepsilon(\chi) = \frac{1}{|G|} \sum_{g \in G} \chi(g^2).$$
- Then $\varepsilon(\chi) = 1, -1$, or 0 when π is an \mathbb{R} -representation, χ is real-valued but π is not an \mathbb{R} -representation, or χ is not real-valued, respectively.
- If $\varepsilon(\chi) = -1$, χ is *symplectic*. If $\varepsilon(\chi) = 1$, χ is *orthogonal*.

Frobenius-Schur Indicators

- Let G be a finite group, and (π, V) an irreducible complex representation of G with character χ .
- The *Frobenius-Schur indicator* of π , or of χ , is
$$\varepsilon(\chi) = \frac{1}{|G|} \sum_{g \in G} \chi(g^2).$$
- Then $\varepsilon(\chi) = 1, -1$, or 0 when π is an \mathbb{R} -representation, χ is real-valued but π is not an \mathbb{R} -representation, or χ is not real-valued, respectively.
- If $\varepsilon(\chi) = -1$, χ is *symplectic*. If $\varepsilon(\chi) = 1$, χ is *orthogonal*.

Frobenius-Schur Indicators

- Let G be a finite group, and (π, V) an irreducible complex representation of G with character χ .
- The *Frobenius-Schur indicator* of π , or of χ , is
$$\varepsilon(\chi) = \frac{1}{|G|} \sum_{g \in G} \chi(g^2).$$
- Then $\varepsilon(\chi) = 1, -1$, or 0 when π is an \mathbb{R} -representation, χ is real-valued but π is not an \mathbb{R} -representation, or χ is not real-valued, respectively.
- If $\varepsilon(\chi) = -1$, χ is *symplectic*. If $\varepsilon(\chi) = 1$, χ is *orthogonal*.

Basic Goal

- Given a finite group of Lie type G , determine the Frobenius-Schur indicator $\varepsilon(\chi)$ for each complex irreducible character χ of G . This determines the classification of the irreducible representations of G over \mathbb{R} .
- Much work has been done on this question (along with the related questions regarding rationality) by Geck, Gow, Janusz, Lusztig, Malle, Marberg, Ohmori, Prasad, Przygocki, Tiep, Turull, Zalesski,...

Basic Goal

- Given a finite group of Lie type G , determine the Frobenius-Schur indicator $\varepsilon(\chi)$ for each complex irreducible character χ of G . This determines the classification of the irreducible representations of G over \mathbb{R} .
- Much work has been done on this question (along with the related questions regarding rationality) by Geck, Gow, Janusz, Lusztig, Malle, Marberg, Ohmori, Prasad, Przygocki, Tiep, Turull, Zalesski,...

However, this goal is not yet completely accomplished for the finite unitary group, $G = \mathrm{U}(n, \mathbb{F}_q)$.

Here, $\mathrm{U}(n, \mathbb{F}_q)$ is defined as \mathbf{G}^F , where $\mathbf{G} = \mathrm{GL}(n, \bar{\mathbb{F}}_q)$ and

$$F((a_{ij})) = {}^T(a_{ij}^q)^{-1}.$$

Note that on $\mathrm{GL}(1, \bar{\mathbb{F}}_q)$, which we may identify with $\bar{\mathbb{F}}_q^\times$, F acts as $F(\alpha) = \alpha^{-q}$.

Semisimple characters

Let \mathbf{G} be a connected reductive group with connected center, defined over \mathbb{F}_q by Frobenius F , and let $G = \mathbf{G}^F$. Suppose $p = \text{char}(\mathbb{F}_q)$ is a *good* prime for \mathbf{G} . Then $\chi \in \text{Irr}(G)$ is a *semisimple* character of G when $\chi(1)$ is prime to p .

Main Result

Theorem (Srinivasan-V)

Let $G = \mathrm{U}(2m, \mathbb{F}_q)$, with q odd. Then the number of semisimple symplectic characters of G is q^{m-1} , and these are in one-to-one correspondence with self-dual polynomials in $\mathbb{F}_q[t]$ of degree $2m$ with constant term -1 .

A polynomial $f(t) \in \mathbb{F}_q[t]$ is self-dual when it is monic, and $\alpha \in \bar{\mathbb{F}}_q$ is a root of $f(t)$ with multiplicity k if and only if α^{-1} is.

Proposition (Ohmori, V)

Let $G = \mathrm{U}(n, \mathbb{F}_q)$, and $\chi \in \mathrm{Irr}(G)$ a semisimple real-valued character. Then:

- ❶ *If n is odd or q is even, then $\varepsilon(\chi) = 1$.*
- ❷ *If n is even and q is odd, then $\varepsilon(\chi) = \omega_\chi(z)$, where ω_χ is the central character of the representation affording χ , and z is a generator for $Z(G)$.*

That is, G can only have semisimple symplectic characters if $n = 2m$ and q is odd.

Remark. Ohmori calculated $\varepsilon(\chi)$ when χ is a unipotent character of $G = \mathrm{U}(n, \mathbb{F}_q)$, q odd. The result implies there can be no $t \in Z(G)$ such that $\varepsilon(\chi) = \omega_\chi(t)$ for all $\chi \in \mathrm{Irr}(G)$.

Real-valued semisimple characters

- If G is a connected reductive group with connected center, defined over \mathbb{F}_q by Frobenius F , let G^* be a group dual to G , with dual Frobenius F^* . Let $G = G^F$ and $G^* = G^{*F^*}$.
- The Deligne-Lusztig theory of characters for G^F gives a bijection $(s) \leftrightarrow \chi_{(s)}$ between semisimple classes of G^* and semisimple characters of G .
- The conjugacy class (s) is *real* if s is conjugate to s^{-1} .

Real-valued semisimple characters

- If G is a connected reductive group with connected center, defined over \mathbb{F}_q by Frobenius F , let G^* be a group dual to G , with dual Frobenius F^* . Let $G = G^F$ and $G^* = G^{*F^*}$.
- The Deligne-Lusztig theory of characters for G^F gives a bijection $(s) \leftrightarrow \chi_{(s)}$ between semisimple classes of G^* and semisimple characters of G .
- The conjugacy class (s) is *real* if s is conjugate to s^{-1} .

Real-valued semisimple characters

- If G is a connected reductive group with connected center, defined over \mathbb{F}_q by Frobenius F , let G^* be a group dual to G , with dual Frobenius F^* . Let $G = G^F$ and $G^* = G^{*F^*}$.
- The Deligne-Lusztig theory of characters for G^F gives a bijection $(s) \leftrightarrow \chi_{(s)}$ between semisimple classes of G^* and semisimple characters of G .
- The conjugacy class (s) is *real* if s is conjugate to s^{-1} .

Real-valued semisimple characters

Lemma

In the bijection $(s) \leftrightarrow \chi_{(s)}$, (s) is a real semisimple class if and only if $\chi_{(s)}$ is a real-valued character.

Navarro-Tiep: One may replace "real" with "rational" in the above, for a more general result.

Take G in duality to G^* as before, with respect to the F -stable maximal torus T of G , in an F -stable Borel, and an F^* -stable maximal torus T^* of G^* in an F^* -stable Borel of G^* . There is an isomorphism

$$(T^*)^{F^*} \cong \text{Irr}(T^F),$$

which gives rise to a homomorphism $z \mapsto \hat{z}$ from $Z(G^*)^{F^*}$ to linear characters of G^F .

This map, in turn, allows us to define a map $s \mapsto \hat{s}$ from semisimple elements of G^* (an F^* -stable element of the center of $C_{G^*}(s)$) to linear characters of $C_G(s)^F$.

Lemma (Fong-Srinivasan)

Let $G^F = \mathrm{U}(n, \mathbb{F}_q)$, $\chi_{(s)}$ be a semisimple character of G^F , and $z \in Z(G^F)$. Then $\omega_{\chi_{(s)}}(z) = \hat{s}(z)$.

Lemma

$$\hat{s}(z) = \hat{z}(s).$$

So, for $G = \mathrm{U}(2m, \mathbb{F}_q)$, q odd, we have $\varepsilon(\chi_{(s)}) = \omega_{\chi_{(s)}}(z) = \hat{z}(s)$ for real-valued $\chi_{(s)}$, where z is a generator for $Z(G)$.

Lemma

When $G = \mathrm{U}(n, \mathbb{F}_q)$, q is odd, and z is a generator of $Z(G)$, then for any semisimple s of G , $\hat{z}(s) = (-1)^k$, where k is the multiplicity of -1 as an eigenvalue of s .

To obtain the main result, we now analyze the real semisimple classes of $\mathrm{U}(n, \mathbb{F}_q)$.

- When $G^F = \mathrm{U}(n, \mathbb{F}_q)$, $F(\alpha) = \alpha^{-q}$ for $\alpha \in \bar{\mathbb{F}}_q^\times$.
- A semisimple class of G^F consists of a choice of F -orbits $[\alpha]_F$ with multiplicity, with total size n .
- The semisimple class is real whenever the orbit $[\alpha]_F$ always has the same multiplicity as $[\alpha^{-1}]_F$.
- If $\tilde{F}(\alpha) = \alpha^q$, then $[\alpha]_{\tilde{F}} \cup [\alpha^{-1}]_{\tilde{F}} = [\alpha]_F \cup [\alpha^{-1}]_F$.

- When $\mathbf{G}^F = \mathrm{U}(n, \mathbb{F}_q)$, $F(\alpha) = \alpha^{-q}$ for $\alpha \in \bar{\mathbb{F}}_q^\times$.
- A semisimple class of \mathbf{G}^F consists of a choice of F -orbits $[\alpha]_F$ with multiplicity, with total size n .
- The semisimple class is real whenever the orbit $[\alpha]_F$ always has the same multiplicity as $[\alpha^{-1}]_F$.
- If $\tilde{F}(\alpha) = \alpha^q$, then $[\alpha]_{\tilde{F}} \cup [\alpha^{-1}]_{\tilde{F}} = [\alpha]_F \cup [\alpha^{-1}]_F$.

- When $\mathbf{G}^F = \mathrm{U}(n, \mathbb{F}_q)$, $F(\alpha) = \alpha^{-q}$ for $\alpha \in \bar{\mathbb{F}}_q^\times$.
- A semisimple class of \mathbf{G}^F consists of a choice of F -orbits $[\alpha]_F$ with multiplicity, with total size n .
- The semisimple class is real whenever the orbit $[\alpha]_F$ always has the same multiplicity as $[\alpha^{-1}]_F$.
- If $\tilde{F}(\alpha) = \alpha^q$, then $[\alpha]_{\tilde{F}} \cup [\alpha^{-1}]_{\tilde{F}} = [\alpha]_F \cup [\alpha^{-1}]_F$.

- When $\mathbf{G}^F = \mathrm{U}(n, \mathbb{F}_q)$, $F(\alpha) = \alpha^{-q}$ for $\alpha \in \bar{\mathbb{F}}_q^\times$.
- A semisimple class of \mathbf{G}^F consists of a choice of F -orbits $[\alpha]_F$ with multiplicity, with total size n .
- The semisimple class is real whenever the orbit $[\alpha]_F$ always has the same multiplicity as $[\alpha^{-1}]_F$.
- If $\tilde{F}(\alpha) = \alpha^q$, then $[\alpha]_{\tilde{F}} \cup [\alpha^{-1}]_{\tilde{F}} = [\alpha]_F \cup [\alpha^{-1}]_F$.

It follows that real semisimple classes of $U(n, \mathbb{F}_q)$ correspond to self-dual polynomials in $\mathbb{F}_q[t]$ of degree n .

When $n = 2m$, q odd, the constant term of the polynomial corresponding to real (s) is exactly $(-1)^k$, where k is the multiplicity of -1 as an eigenvalue of (s) .

The main result follows, since, for (s) a real semisimple class, $\chi_{(s)}$ is real-valued, and

$$\varepsilon(\chi_{(s)}) = \hat{z}(s) = (-1)^k,$$

while $(-1)^k$ is precisely the constant term of the self-dual polynomial of degree $2m$ in $\mathbb{F}_q[t]$ corresponding to (s) .

One can quickly enumerate the number of such self-dual polynomials with k odd to be q^{m-1} , corresponding to semisimple symplectic characters, and those with k even to be q^m , corresponding to semisimple orthogonal characters.

- As mentioned, when $G = \mathrm{U}(n, \mathbb{F}_q)$, q odd, Ohmori determined $\varepsilon(\psi)$ for ψ unipotent. We also know $\varepsilon(\chi_{(s)})$ when $\chi_{(s)}$ is semisimple.
- If one can understand the behavior of the F-S indicator under Deligne-Lusztig induction R_L^G , we could take advantage of the Jordan decomposition of characters, and then the basic goal of computing $\varepsilon(\chi)$ would be reduced to that of unipotent characters.

- As mentioned, when $G = \mathrm{U}(n, \mathbb{F}_q)$, q odd, Ohmori determined $\varepsilon(\psi)$ for ψ unipotent. We also know $\varepsilon(\chi_{(s)})$ when $\chi_{(s)}$ is semisimple.
- If one can understand the behavior of the F-S indicator under Deligne-Lusztig induction R_L^G , we could take advantage of the Jordan decomposition of characters, and then the basic goal of computing $\varepsilon(\chi)$ would be reduced to that of unipotent characters.

- More generally, if $G = \mathbf{G}^F$ is a finite reductive group, and χ is a complex irreducible character of G with Jordan decomposition (s, ψ) , so s is a semisimple element of G^* , and ψ is a unipotent character of $C_{G^*}(s)$, precisely what conditions of (s, ψ) are equivalent to χ being real-valued? It seems we need s is a real element of G^* , and if $ws w^{-1} = s^{-1}$, then we also need ${}^w\psi = \bar{\psi}$.
- Thank you!

- More generally, if $G = \mathbf{G}^F$ is a finite reductive group, and χ is a complex irreducible character of G with Jordan decomposition (s, ψ) , so s is a semisimple element of G^* , and ψ is a unipotent character of $C_{G^*}(s)$, precisely what conditions of (s, ψ) are equivalent to χ being real-valued? It seems we need s is a real element of G^* , and if $ws w^{-1} = s^{-1}$, then we also need ${}^w\psi = \bar{\psi}$.
- Thank you!