

Counting characters in blocks of solvable groups with abelian defect group

Mark L. Lewis

Kent State University

June 4, 2012

Joint work with James P. Cossey
2012 CMS Summer Meeting - Regina
Special Session on Representations of Groups,
Lie Algebras, and Hopf Algebras

Introduction

Throughout this note G is a finite group and p is a fixed prime.

Introduction

Throughout this note G is a finite group and p is a fixed prime.

We write $\text{Irr}(G)$ for the irreducible characters of G .

Introduction

Throughout this note G is a finite group and p is a fixed prime.

We write $\text{Irr}(G)$ for the irreducible characters of G .

We write $\text{IBr}_p(G)$ for the irreducible Brauer characters of G .

Introduction

Throughout this note G is a finite group and p is a fixed prime.

We write $\text{Irr}(G)$ for the irreducible characters of G .

We write $\text{IBr}_p(G)$ for the irreducible Brauer characters of G .

These are complex-valued class functions on the set of p -regular elements of G that in some sense describe the irreducible representations of G over an algebraically closed field of characteristic p .

Introduction

Throughout this note G is a finite group and p is a fixed prime.

We write $\text{Irr}(G)$ for the irreducible characters of G .

We write $\text{IBr}_p(G)$ for the irreducible Brauer characters of G .

These are complex-valued class functions on the set of p -regular elements of G that in some sense describe the irreducible representations of G over an algebraically closed field of characteristic p .

Let α° denote the restriction of the complex valued class function α to the p -regular elements of G .

Introduction

It has long been known that if $\chi \in \text{Irr}(G)$, we have

$$\chi^o = \sum_{\varphi \in \text{IBr}_p(G)} d_{\chi\varphi} \varphi,$$

where the $\{d_{\chi\varphi}\}$ are uniquely defined non-negative integers, called the decomposition numbers of χ .

Introduction

It has long been known that if $\chi \in \text{Irr}(G)$, we have

$$\chi^o = \sum_{\varphi \in \text{IBr}_p(G)} d_{\chi\varphi} \varphi,$$

where the $\{d_{\chi\varphi}\}$ are uniquely defined non-negative integers, called the decomposition numbers of χ .

If G is p -solvable, there is more that can be said.

Introduction

It has long been known that if $\chi \in \text{Irr}(G)$, we have

$$\chi^o = \sum_{\varphi \in \text{IBr}_p(G)} d_{\chi\varphi} \varphi,$$

where the $\{d_{\chi\varphi}\}$ are uniquely defined non-negative integers, called the decomposition numbers of χ .

If G is p -solvable, there is more that can be said.

The Fong-Swan theorem says that if $\varphi \in \text{IBr}_p(G)$, then there is a character $\chi \in \text{Irr}(G)$ such that $\chi^o = \varphi$.

Introduction

It has long been known that if $\chi \in \text{Irr}(G)$, we have

$$\chi^o = \sum_{\varphi \in \text{IBr}_p(G)} d_{\chi\varphi} \varphi,$$

where the $\{d_{\chi\varphi}\}$ are uniquely defined non-negative integers, called the decomposition numbers of χ .

If G is p -solvable, there is more that can be said.

The Fong-Swan theorem says that if $\varphi \in \text{IBr}_p(G)$, then there is a character $\chi \in \text{Irr}(G)$ such that $\chi^o = \varphi$.

In this case, we say that χ is a lift of φ .

Introduction

We define $L_\varphi = \{\chi \in \text{Irr}(G) \mid \chi^\circ = \varphi\}$ to be the set of lifts of φ .

Introduction

We define $L_\varphi = \{\chi \in \text{Irr}(G) \mid \chi^\circ = \varphi\}$ to be the set of lifts of φ .

If G is p -solvable and B is a block of G with abelian defect group D , it is known that if $\varphi \in \text{IBr}_p(B)$, then $|L_\varphi| \leq |D|$.

Introduction

We define $L_\varphi = \{\chi \in \text{Irr}(G) \mid \chi^\circ = \varphi\}$ to be the set of lifts of φ .

If G is p -solvable and B is a block of G with abelian defect group D , it is known that if $\varphi \in \text{IBr}_p(B)$, then $|L_\varphi| \leq |D|$.

We examine necessary and sufficient conditions for a character $\varphi \in \text{IBr}_p(B)$ with abelian defect group to attain the upper bound of $|L_\varphi| = |D|$.

Introduction

We want to point out that the proofs of the main results are “character theoretic” in nature, in that the proofs only use the characters and not the underlying representations.

Introduction

We want to point out that the proofs of the main results are “character theoretic” in nature, in that the proofs only use the characters and not the underlying representations.

In particular this means that all of the results in this paper also hold in the setting of Isaacs’ “ π -partial” characters of π -separable groups and the proofs would be the same.

Theorems

Given a Brauer character φ of G , we let
$$\text{Irr}(G : \varphi) = \{\chi \in \text{Irr}(G) \mid d_{\chi\varphi} \neq 0\}.$$

Theorems

Given a Brauer character φ of G , we let
 $\text{Irr}(G : \varphi) = \{\chi \in \text{Irr}(G) \mid d_{\chi\varphi} \neq 0\}.$

Theorem

Let G be a p -solvable group and B a block of G with abelian defect group D and let $\varphi \in \text{IBr}_p(B)$. Then the following conditions are equivalent:

- ① $|L_\varphi| = |D|.$
- ② $\text{Irr}(G : \varphi) = L_\varphi.$
- ③ $|\text{Irr}(G : \varphi)| = |D|.$

Theorems

The condition $\text{Irr}(G : \varphi) = L_\varphi$ says that any irreducible character whose restriction contains φ is a lift of φ .

Theorems

The condition $\text{Irr}(G : \varphi) = L_\varphi$ says that any irreducible character whose restriction contains φ is a lift of φ .

The condition $|\text{Irr}(G : \varphi)| = |D|$ says that there are $|D|$ irreducible characters whose restriction contains φ . Notice that this condition does not involve the number of lifts at all.

Theorems

The condition $\text{Irr}(G : \varphi) = L_\varphi$ says that any irreducible character whose restriction contains φ is a lift of φ .

The condition $|\text{Irr}(G : \varphi)| = |D|$ says that there are $|D|$ irreducible characters whose restriction contains φ . Notice that this condition does not involve the number of lifts at all.

It is interesting to have a condition about lifts be equivalent to a “global” condition that does not involve lifts.

Theorems

The condition $\text{Irr}(G : \varphi) = L_\varphi$ says that any irreducible character whose restriction contains φ is a lift of φ .

The condition $|\text{Irr}(G : \varphi)| = |D|$ says that there are $|D|$ irreducible characters whose restriction contains φ . Notice that this condition does not involve the number of lifts at all.

It is interesting to have a condition about lifts be equivalent to a “global” condition that does not involve lifts.

It turns out that these global conditions are fairly easy to prove. It is perhaps more interesting to look at a local condition.

Theorems

Given any block B of a solvable group G with defect group D , we define (up to conjugacy) a certain canonical subgroup V_B containing (a conjugate of) D as a Sylow p -subgroup, called the Fong subgroup of B .

Theorems

Given any block B of a solvable group G with defect group D , we define (up to conjugacy) a certain canonical subgroup V_B containing (a conjugate of) D as a Sylow p -subgroup, called the Fong subgroup of B .

In our Main Theorem, we show that condition that $|L_\varphi| = |D|$ is equivalent to a condition about V_B .

Theorems

Given any block B of a solvable group G with defect group D , we define (up to conjugacy) a certain canonical subgroup V_B containing (a conjugate of) D as a Sylow p -subgroup, called the Fong subgroup of B .

In our Main Theorem, we show that condition that $|L_\varphi| = |D|$ is equivalent to a condition about V_B .

Theorem

Let G be a p -solvable group and B a block of G with abelian defect group, let V_B be a Fong subgroup for B , and let D be a defect group for B contained in V_B . Suppose $\varphi \in \text{IBr}_p(B)$. Then $|L_\varphi| = |D|$ if and only if $D \leq Z(N_{V_B}(D))$.

The subgroup V_B

Let B be a block of a p -solvable group G . We will define the Fong pair (V_B, B_V) and discuss some of the basic properties of this pair.

The subgroup V_B

Let B be a block of a p -solvable group G . We will define the Fong pair (V_B, B_V) and discuss some of the basic properties of this pair.

Our definition is similar to (but not identical) to the classic reduction by Fong.

The subgroup V_B

Let B be a block of a p -solvable group G . We will define the Fong pair (V_B, B_V) and discuss some of the basic properties of this pair.

Our definition is similar to (but not identical) to the classic reduction by Fong.

However, the classical approach developed there relies heavily on many of the module-theoretic aspects of representation theory that do not translate well (if at all) to the π -case.

The subgroup V_B

Let B be a block of a p -solvable group G . We will define the Fong pair (V_B, B_V) and discuss some of the basic properties of this pair.

Our definition is similar to (but not identical) to the classic reduction by Fong.

However, the classical approach developed there relies heavily on many of the module-theoretic aspects of representation theory that do not translate well (if at all) to the π -case.

Thus we develop a purely “character theoretic” approach.

The subgroup V_B

Let G be a solvable group and let B be a block of G .

The subgroup V_B

Let G be a solvable group and let B be a block of G .

Write $M = \mathbf{O}_{p'}(G)$, and let $\alpha \in \text{Irr}(M)$ be covered by B .

The subgroup V_B

Let G be a solvable group and let B be a block of G .

Write $M = \mathbf{O}_{p'}(G)$, and let $\alpha \in \text{Irr}(M)$ be covered by B .

If α is invariant in G , then we define the Fong pair (V_B, B_V) to be (G, B) .

The subgroup V_B

Let G be a solvable group and let B be a block of G .

Write $M = \mathbf{O}_{p'}(G)$, and let $\alpha \in \text{Irr}(M)$ be covered by B .

If α is invariant in G , then we define the Fong pair (V_B, B_V) to be (G, B) .

If α is not invariant in G , then we let T be the stabilizer of α in G .

The subgroup V_B

Let G be a solvable group and let B be a block of G .

Write $M = \mathbf{O}_{p'}(G)$, and let $\alpha \in \text{Irr}(M)$ be covered by B .

If α is invariant in G , then we define the Fong pair (V_B, B_V) to be (G, B) .

If α is not invariant in G , then we let T be the stabilizer of α in G .

By the Fong-Reynolds theorem, there exists a block B_1 of T covering α such that induction is a bijection from B_1 to G .

The subgroup V_B

Let G be a solvable group and let B be a block of G .

Write $M = \mathbf{O}_{p'}(G)$, and let $\alpha \in \text{Irr}(M)$ be covered by B .

If α is invariant in G , then we define the Fong pair (V_B, B_V) to be (G, B) .

If α is not invariant in G , then we let T be the stabilizer of α in G .

By the Fong-Reynolds theorem, there exists a block B_1 of T covering α such that induction is a bijection from B_1 to G .

In this case we define the Fong pair (V_B, B_V) for B to be a Fong pair for B_1 .

The subgroup V_B

It is perhaps worth mentioning in the above definition that in general one could have $\mathbf{O}_{p'}(T) > \mathbf{O}_{p'}(G)$, so you might have to repeat the process several times.

The subgroup V_B

It is perhaps worth mentioning in the above definition that in general one could have $\mathbf{O}_{p'}(T) > \mathbf{O}_{p'}(G)$, so you might have to repeat the process several times.

By the Fong-Reynolds theorem, any Sylow p -subgroup of V_B is a defect group for the block B .

The subgroup V_B

It is perhaps worth mentioning in the above definition that in general one could have $\mathbf{O}_{p'}(T) > \mathbf{O}_{p'}(G)$, so you might have to repeat the process several times.

By the Fong-Reynolds theorem, any Sylow p -subgroup of V_B is a defect group for the block B .

One could have chosen a different character α of M covered by B , but a different choice would have yielded a conjugate Fong pair.

The subgroup V_B

Induction preserves decomposition numbers. In other words, with the above notation, if $\chi \in \text{Irr}(B)$ and $\varphi \in \text{IBr}_p(B)$ are induced by $\chi_1 \in \text{Irr}(B_1)$ and $\varphi_1 \in \text{IBr}_p(B_1)$, respectively, then $d_{\chi\varphi} = d_{\chi_1\varphi_1}$.

The subgroup V_B

Induction preserves decomposition numbers. In other words, with the above notation, if $\chi \in \text{Irr}(B)$ and $\varphi \in \text{IBr}_p(B)$ are induced by $\chi_1 \in \text{Irr}(B_1)$ and $\varphi_1 \in \text{IBr}_p(B_1)$, respectively, then $d_{\chi\varphi} = d_{\chi_1\varphi_1}$.

This yields:

The subgroup V_B

Induction preserves decomposition numbers. In other words, with the above notation, if $\chi \in \text{Irr}(B)$ and $\varphi \in \text{IBr}_p(B)$ are induced by $\chi_1 \in \text{Irr}(B_1)$ and $\varphi_1 \in \text{IBr}_p(B_1)$, respectively, then $d_{\chi\varphi} = d_{\chi_1\varphi_1}$.

This yields:

Lemma

Let G be a p -solvable group and B a block of G with Fong pair (V_B, B_V) . Let $\varphi \in \text{IBr}_p(B)$ and let $\widehat{\varphi}$ denote the corresponding character of V_B . Then induction is a bijection from $L_{\widehat{\varphi}}$ to L_{φ} and from $\text{Irr}(V_B : \widehat{\varphi})$ to $\text{Irr}(G : \varphi)$.

The subgroup V_B

Induction preserves decomposition numbers. In other words, with the above notation, if $\chi \in \text{Irr}(B)$ and $\varphi \in \text{IBr}_p(B)$ are induced by $\chi_1 \in \text{Irr}(B_1)$ and $\varphi_1 \in \text{IBr}_p(B_1)$, respectively, then $d_{\chi\varphi} = d_{\chi_1\varphi_1}$.

This yields:

Lemma

Let G be a p -solvable group and B a block of G with Fong pair (V_B, B_V) . Let $\varphi \in \text{IBr}_p(B)$ and let $\widehat{\varphi}$ denote the corresponding character of V_B . Then induction is a bijection from $L_{\widehat{\varphi}}$ to L_{φ} and from $\text{Irr}(V_B : \widehat{\varphi})$ to $\text{Irr}(G : \varphi)$.

We note that this works whether or not D is abelian.

The subgroup V_B

We then use a result of Dade (whose proof was finally published by Turull) to obtain the following (this result uses the fact that D is abelian):

The subgroup V_B

We then use a result of Dade (whose proof was finally published by Turull) to obtain the following (this result uses the fact that D is abelian):

Lemma

Let G be a p -solvable group and B a block of G with Fong pair (V_B, B_V) and abelian defect group D contained in V_B . Then there exists a Brauer character $\tilde{\varphi} \in \text{IBr}_p(N_{V_B}(D))$ that satisfies $|\text{Irr}(N_{V_B}(D) : \tilde{\varphi})| = |\text{Irr}(G : \varphi)|$ and $|L_\varphi| = |L_{\tilde{\varphi}}|$.

The subgroup V_B

We then use a result of Dade (whose proof was finally published by Turull) to obtain the following (this result uses the fact that D is abelian):

Lemma

Let G be a p -solvable group and B a block of G with Fong pair (V_B, B_V) and abelian defect group D contained in V_B . Then there exists a Brauer character $\tilde{\varphi} \in \text{IBr}_p(N_{V_B}(D))$ that satisfies $|\text{Irr}(N_{V_B}(D) : \tilde{\varphi})| = |\text{Irr}(G : \varphi)|$ and $|L_\varphi| = |L_{\tilde{\varphi}}|$.

This gives a way of translating the hypotheses between G and $N_{V_B}(D)$.

The Local condition

We now give some idea of the proofs.

The Local condition

We now give some idea of the proofs.

We start by assuming that $D \leq Z(N_{V_B}(G))$. This next lemma can be viewed as having $G = N_{V_B}(G)$, and we prove the result in this case.

The Local condition

We now give some idea of the proofs.

We start by assuming that $D \leq Z(N_{V_B}(G))$. This next lemma can be viewed as having $G = N_{V_B}(G)$, and we prove the result in this case.

Lemma

Let p be a prime, let G be a p -solvable group, let $N = O_{p'}(G)$, and let $\varphi \in \text{IBr}_p(G)$ be contained in the block B with defect group D . Suppose $\alpha \in \text{Irr}(N)$ is an irreducible constituent of φ_N and is G -invariant. If $D \leq Z(G)$, then $|L_\varphi| = |D|$. Moreover, φ is the unique Brauer character of B .

The Local condition

Proof:

The Local condition

Proof:

D is a Sylow p -subgroup of G by a theorem of Fong.

The Local condition

Proof:

D is a Sylow p -subgroup of G by a theorem of Fong.

Let H be a Hall p -complement for G .

The Local condition

Proof:

D is a Sylow p -subgroup of G by a theorem of Fong.

Let H be a Hall p -complement for G .

Since D is a Sylow p -subgroup and central, we have that
 $G = H \times D$.

The Local condition

Proof:

D is a Sylow p -subgroup of G by a theorem of Fong.

Let H be a Hall p -complement for G .

Since D is a Sylow p -subgroup and central, we have that $G = H \times D$.

$\varphi_H = \theta \in \text{Irr}(H)$, and the set of lifts of φ is the set $\text{Irr}(G \mid \theta)$.

The Local condition

Proof:

D is a Sylow p -subgroup of G by a theorem of Fong.

Let H be a Hall p -complement for G .

Since D is a Sylow p -subgroup and central, we have that $G = H \times D$.

$\varphi_H = \theta \in \text{Irr}(H)$, and the set of lifts of φ is the set $\text{Irr}(G \mid \theta)$.

Since $|\text{Irr}(G \mid \theta)| = |D|$, this gives $|L_\varphi| = |D|$, as desired.

The Local condition

Proof:

D is a Sylow p -subgroup of G by a theorem of Fong.

Let H be a Hall p -complement for G .

Since D is a Sylow p -subgroup and central, we have that $G = H \times D$.

$\varphi_H = \theta \in \text{Irr}(H)$, and the set of lifts of φ is the set $\text{Irr}(G \mid \theta)$.

Since $|\text{Irr}(G \mid \theta)| = |D|$, this gives $|L_\varphi| = |D|$, as desired.

Finally, since G is the direct product of a p -group and a p' -group, then φ is the unique Brauer character in B .

The Local condition

Using the correspondence between G and $N_{V_B}(D)$, this yields the theorem:

The Local condition

Using the correspondence between G and $N_{V_B}(D)$, this yields the theorem:

Theorem

Let G be a p -solvable group and let B be a block of G with abelian defect group D , and suppose $\varphi \in \text{IBr}_p(B)$. Let V_B be a Fong subgroup associated to B . If D is central in $N_{V_B}(D)$, then $|L_\varphi| = |D|$. Moreover, in this case φ is the unique Brauer character in B .

$$|\mathrm{Irr}(G : \varphi)| = |D|$$

We next consider $|\mathrm{Irr}(G : \varphi)| = |D|$ when D is a normal, Sylow subgroup.

$$|\mathrm{Irr}(G : \varphi)| = |D|$$

We next consider $|\mathrm{Irr}(G : \varphi)| = |D|$ when D is a normal, Sylow subgroup.

Lemma

Let G be a p -solvable group, let D be a normal, abelian Sylow p -subgroup, and let $\varphi \in \mathrm{IBr}_p(G)$. Then the following are true:

- 1 $|\mathrm{Irr}(G : \varphi)| \leq |D|$.
- 2 If $|\mathrm{Irr}(G : \varphi)| = |D|$, then $\mathrm{Irr}(G : \varphi) = L_\varphi$.

$$|\mathrm{Irr}(G : \varphi)| = |D|$$

We next consider $|\mathrm{Irr}(G : \varphi)| = |D|$ when D is a normal, Sylow subgroup.

Lemma

Let G be a p -solvable group, let D be a normal, abelian Sylow p -subgroup, and let $\varphi \in \mathrm{IBr}_p(G)$. Then the following are true:

- 1 $|\mathrm{Irr}(G : \varphi)| \leq |D|$.
- 2 If $|\mathrm{Irr}(G : \varphi)| = |D|$, then $\mathrm{Irr}(G : \varphi) = L_\varphi$.

We note that the first conclusion is a consequence of the $k(B)$ -conjecture, but we have a simple proof.

$$|\mathrm{Irr}(G : \varphi)| = |D|$$

We use the correspondence to reduce this next theorem to the situation of the previous lemma, and applying that lemma, we obtain:

$$|\mathrm{Irr}(G : \varphi)| = |D|$$

We use the correspondence to reduce this next theorem to the situation of the previous lemma, and applying that lemma, we obtain:

Theorem

Let G be a p -solvable group, and let $\varphi \in \mathrm{IBr}_p(G)$ lie in a block B with abelian defect group D . If $|\mathrm{Irr}(G : \varphi)| = |D|$, then $\mathrm{Irr}(G : \varphi) = L_\varphi$. Moreover, $\mathrm{IBr}_p(B) = \{\varphi\}$ and $\mathrm{Irr}(B) = L_\varphi$.

$$|L_\varphi| = |D|$$

We now consider the hypothesis that $|L_\varphi| = |D|$.

$$|L_\varphi| = |D|$$

We now consider the hypothesis that $|L_\varphi| = |D|$.

We first consider the case where D is normal.

$$|L_\varphi| = |D|$$

We now consider the hypothesis that $|L_\varphi| = |D|$.

We first consider the case where D is normal.

For a character δ of the normal subgroup D of G , we will let G_δ denote the inertia subgroup of δ .

$$|L_\varphi| = |D|$$

We now consider the hypothesis that $|L_\varphi| = |D|$.

We first consider the case where D is normal.

For a character δ of the normal subgroup D of G , we will let G_δ denote the inertia subgroup of δ .

Lemma

Let G be a p -solvable group, and let $\varphi \in \text{IBr}_p(G)$ be in a block with a normal, abelian defect group D . Let $\delta \in \text{Irr}(D)$. If $|L_\varphi| = |D|$, then $|L_\varphi \cap \text{Irr}(G \mid \delta)| = |G : G_\delta|$.

$$|L_\varphi| = |D|$$

The following theorem includes the most difficult part of our argument:

$$|L_\varphi| = |D|$$

The following theorem includes the most difficult part of our argument:

Theorem

Let G be a p -solvable group, let D be a normal, abelian Sylow p -subgroup, and let $\varphi \in \text{IBr}_p(G)$. Assume that $\varphi_{\mathbf{O}_{p'}(G)}$ is homogeneous. If $|L_\varphi| = |D|$, then $D \leq Z(G)$.

$$|L_\varphi| = |D|$$

The following theorem includes the most difficult part of our argument:

Theorem

Let G be a p -solvable group, let D be a normal, abelian Sylow p -subgroup, and let $\varphi \in \text{IBr}_p(G)$. Assume that $\varphi_{\mathbf{O}_{p'}(G)}$ is homogeneous. If $|L_\varphi| = |D|$, then $D \leq Z(G)$.

In particular, this uses the following theorem of Dolfi:

$$|L_\varphi| = |D|$$

The following theorem includes the most difficult part of our argument:

Theorem

Let G be a p -solvable group, let D be a normal, abelian Sylow p -subgroup, and let $\varphi \in \text{IBr}_p(G)$. Assume that $\varphi_{\mathbf{O}_{p'}(G)}$ is homogeneous. If $|L_\varphi| = |D|$, then $D \leq Z(G)$.

In particular, this uses the following theorem of Dolfi:

Theorem (Dolfi)

Let G be a solvable group acting faithfully and coprimely on a group K . Then there exist $x, y \in K$ so that $C_G(x) \cap C_G(y) = 1$.

$$|L_\varphi| = |D|$$

This proves:

$$|L_\varphi| = |D|$$

This proves:

Corollary

Let G be a p -solvable group, and let $\varphi \in \text{IBr}_p(G)$ lie in a block B with abelian defect group D . Let V be a Fong subgroup for B containing D . If $|L_\varphi| = |D|$, then $D \leq Z(N_V(D))$.

$$\text{Irr}(G : \varphi) = L_\varphi$$

Interestingly, in the next lemma we conclude that the defect group is abelian if the defect group is a normal Sylow p -subgroup and every character in $\text{Irr}(G : \varphi)$ is a lift of φ .

$$\text{Irr}(G : \varphi) = L_\varphi$$

Interestingly, in the next lemma we conclude that the defect group is abelian if the defect group is a normal Sylow p -subgroup and every character in $\text{Irr}(G : \varphi)$ is a lift of φ .

Lemma

Let G be a p -solvable group, let D be a normal Sylow p -subgroup, and let $\varphi \in \text{IBr}_p(G)$. If $\text{Irr}(G : \varphi) = L_\varphi$, then $|L_\varphi| = |D|$ and D is abelian.

Example

This section contains a example showing that our Theorem is not true if we replace $N_V(D)$ with $N_G(D)$.

Example

This section contains an example showing that our Theorem is not true if we replace $N_V(D)$ with $N_G(D)$.

Let G be the semi-direct product of a group of order 3 acting on a group of order 91 as a fixed-point free automorphism, so G is a Frobenius group of order 273.

Example

This section contains an example showing that our Theorem is not true if we replace $N_V(D)$ with $N_G(D)$.

Let G be the semi-direct product of a group of order 3 acting on a group of order 91 as a fixed-point free automorphism, so G is a Frobenius group of order 273.

Take $p = 7$.

Example

This section contains an example showing that our Theorem is not true if we replace $N_V(D)$ with $N_G(D)$.

Let G be the semi-direct product of a group of order 3 acting on a group of order 91 as a fixed-point free automorphism, so G is a Frobenius group of order 273.

Take $p = 7$.

Let D be the subgroup of order 7, and notice that D is normal in G but not central.

Example

The group G has a Brauer character $\varphi \in \text{IBr}_7(G)$ whose defect group is D where $|L_\varphi| = 7 = |D|$.