

Lie racks of type D

Unipotent conjugacy classes in finite groups of Lie type

Gastón Andrés García

FaMAF, Universidad Nacional de Córdoba, Argentina
CIEM-CONICET

CMS Summer Meeting

June 2-4, 2012, Regina, Canada

Joint work with N. Andruskiewitsch and G. Carnovale.

Joint work with N. Andruskiewitsch and G. Carnovale.

Problem: Determine when a Lie rack is of type D.

Joint work with N. Andruskiewitsch and G. Carnovale.

Problem: Determine when a Lie rack is of type D.

In group-theoretical terms:

Joint work with N. Andruskiewitsch and G. Carnovale.

Problem: Determine when a Lie rack is of type D.

In group-theoretical terms:

Question

Joint work with N. Andruskiewitsch and G. Carnovale.

Problem: Determine when a Lie rack is of type D.

In group-theoretical terms:

Question

Let \mathcal{O} be a conjugacy class in a finite group G .

Joint work with N. Andruskiewitsch and G. Carnovale.

Problem: Determine when a Lie rack is of type D.

In group-theoretical terms:

Question

Let \mathcal{O} be a conjugacy class in a finite group G . Are there elements $r, s \in \mathcal{O}$ such that their conjugacy classes in the subgroup $\langle r, s \rangle$ of G are different and

Joint work with N. Andruskiewitsch and G. Carnovale.

Problem: Determine when a Lie rack is of type D.

In group-theoretical terms:

Question

Let \mathcal{O} be a conjugacy class in a finite group G . Are there elements $r, s \in \mathcal{O}$ such that their conjugacy classes in the subgroup $\langle r, s \rangle$ of G are different and $(rs)^2 \neq (sr)^2$?

Joint work with N. Andruskiewitsch and G. Carnovale.

Problem: Determine when a Lie rack is of type D.

In group-theoretical terms:

Question

Let \mathcal{O} be a conjugacy class in a finite group G . Are there elements $r, s \in \mathcal{O}$ such that their conjugacy classes in the subgroup $\langle r, s \rangle$ of G are different and $(rs)^2 \neq (sr)^2$?

We deal concretely with the case when G is a finite group of Lie type.

Definition

A rack is a non-empty set X

Definition

A rack is a non-empty set X endowed with an map $\triangleright : X \times X \rightarrow X$ satisfying

Definition

A rack is a non-empty set X endowed with an map $\triangleright : X \times X \rightarrow X$ satisfying

(a) $x \triangleright _$ is a bijection for any $x \in X$,

Definition

A rack is a non-empty set X endowed with an map $\triangleright : X \times X \rightarrow X$ satisfying

- (a) $x \triangleright _$ is a bijection for any $x \in X$,
- (b) $x \triangleright (y \triangleright z) = (x \triangleright y) \triangleright (x \triangleright z)$ for all $x, y, z \in X$.

Definition

A rack is a non-empty set X endowed with an map $\triangleright : X \times X \rightarrow X$ satisfying

- (a) $x \triangleright _$ is a bijection for any $x \in X$,
- (b) $x \triangleright (y \triangleright z) = (x \triangleright y) \triangleright (x \triangleright z)$ for all $x, y, z \in X$.

The archetypical example of a rack is a conjugacy class in a group.

Definition

A rack is a non-empty set X endowed with an map $\triangleright : X \times X \rightarrow X$ satisfying

- (a) $x \triangleright _$ is a bijection for any $x \in X$,
- (b) $x \triangleright (y \triangleright z) = (x \triangleright y) \triangleright (x \triangleright z)$ for all $x, y, z \in X$.

The archetypical example of a rack is a conjugacy class in a group.

We say that a rack is *simple* if $|X| > 1$ and any rack epimorphism $X \twoheadrightarrow Y$ is bijective or $|Y| = 1$.

Definition

A rack X is of type D if

Definition

A rack X is of type D if it contains a decomposable subrack $Y = R \amalg S$ and elements $r \in R$, $s \in S$ such that $r \triangleright (s \triangleright (r \triangleright s)) \neq s$.

Definition

A rack X is of type D if it contains a decomposable subrack $Y = R \amalg S$ and elements $r \in R$, $s \in S$ such that $r \triangleright (s \triangleright (r \triangleright s)) \neq s$.

If \mathcal{O} is a conjugacy class in a finite group G ,

Definition

A rack X is of type D if it contains a decomposable subrack $Y = R \amalg S$ and elements $r \in R$, $s \in S$ such that $r \triangleright (s \triangleright (r \triangleright s)) \neq s$.

If \mathcal{O} is a conjugacy class in a finite group G , then the rack \mathcal{O} is of type D if and only if the answer to our first question is positive.

Definition

A rack X is of type D if it contains a decomposable subrack $Y = R \amalg S$ and elements $r \in R$, $s \in S$ such that $r \triangleright (s \triangleright (r \triangleright s)) \neq s$.

If \mathcal{O} is a conjugacy class in a finite group G , then the rack \mathcal{O} is of type D if and only if the answer to our first question is positive.

Facts:

Definition

A rack X is of type D if it contains a decomposable subrack $Y = R \amalg S$ and elements $r \in R$, $s \in S$ such that $r \triangleright (s \triangleright (r \triangleright s)) \neq s$.

If \mathcal{O} is a conjugacy class in a finite group G , then the rack \mathcal{O} is of type D if and only if the answer to our first question is positive.

Facts: ► If Z is a finite rack that admits a rack epimorphism $Z \twoheadrightarrow X$, where X is of type D, then Z is of type D.

Definition

A rack X is of type D if it contains a decomposable subrack $Y = R \amalg S$ and elements $r \in R, s \in S$ such that $r \triangleright (s \triangleright (r \triangleright s)) \neq s$.

If \mathcal{O} is a conjugacy class in a finite group G , then the rack \mathcal{O} is of type D if and only if the answer to our first question is positive.

Facts: ► If Z is a finite rack that admits a rack epimorphism $Z \twoheadrightarrow X$, where X is of type D, then Z is of type D.

► If Z is indecomposable, then it admits a rack epimorphism $Z \twoheadrightarrow X$ with X simple.

Definition

A rack X is of type D if it contains a decomposable subrack $Y = R \amalg S$ and elements $r \in R, s \in S$ such that $r \triangleright (s \triangleright (r \triangleright s)) \neq s$.

If \mathcal{O} is a conjugacy class in a finite group G , then the rack \mathcal{O} is of type D if and only if the answer to our first question is positive.

Facts: ► If Z is a finite rack that admits a rack epimorphism $Z \twoheadrightarrow X$, where X is of type D, then Z is of type D.

► If Z is indecomposable, then it admits a rack epimorphism $Z \twoheadrightarrow X$ with X simple.

Hence

Problem

Definition

A rack X is of type D if it contains a decomposable subrack $Y = R \amalg S$ and elements $r \in R, s \in S$ such that $r \triangleright (s \triangleright (r \triangleright s)) \neq s$.

If \mathcal{O} is a conjugacy class in a finite group G , then the rack \mathcal{O} is of type D if and only if the answer to our first question is positive.

Facts: ► If Z is a finite rack that admits a rack epimorphism $Z \twoheadrightarrow X$, where X is of type D, then Z is of type D.

► If Z is indecomposable, then it admits a rack epimorphism $Z \twoheadrightarrow X$ with X simple.

Hence

Problem

Determine all simple racks of type D

Finite simple racks where classified in [AG]¹ and [J]².

¹N. Andruskiewitsch and M. Graña, From racks to pointed Hopf algebras, *Adv. Math.* **178** (2003), 177–243.

²D. Joyce, Simple quandles, *J. Algebra* **79** 2 (1982), 307–318. 

Finite simple racks were classified in [AG]¹ and [J]².

We are interested in the (twisted) conjugacy classes of type (L^t, u) :

¹N. Andruskiewitsch and M. Graña, From racks to pointed Hopf algebras, *Adv. Math.* **178** (2003), 177–243.

²D. Joyce, Simple quandles, *J. Algebra* **79** 2 (1982), 307–318. 

Finite simple racks were classified in [AG]¹ and [J]².

We are interested in the (twisted) conjugacy classes of type (L^t, u) :

► (L, t, θ) a triple where L is a simple non-abelian finite group,

¹N. Andruskiewitsch and M. Graña, From racks to pointed Hopf algebras, *Adv. Math.* **178** (2003), 177–243.

²D. Joyce, Simple quandles, *J. Algebra* **79** 2 (1982), 307–318.

Finite simple racks were classified in [AG]¹ and [J]².

We are interested in the (twisted) conjugacy classes of type (L^t, u) :

► (L, t, θ) a triple where L is a simple non-abelian finite group,
 $t \in \mathbb{N}$

¹N. Andruskiewitsch and M. Graña, From racks to pointed Hopf algebras, *Adv. Math.* **178** (2003), 177–243.

²D. Joyce, Simple quandles, *J. Algebra* **79** 2 (1982), 307–318.

Finite simple racks were classified in [AG]¹ and [J]².

We are interested in the (twisted) conjugacy classes of type (L^t, u) :

► (L, t, θ) a triple where L is a simple non-abelian finite group, $t \in \mathbb{N}$ and $\theta \in \text{Aut}(L)$.

¹N. Andruskiewitsch and M. Graña, From racks to pointed Hopf algebras, *Adv. Math.* **178** (2003), 177–243.

²D. Joyce, Simple quandles, *J. Algebra* **79** 2 (1982), 307–318.

Finite simple racks were classified in [AG]¹ and [J]².

We are interested in the (twisted) conjugacy classes of type (L^t, u) :

- ▶ (L, t, θ) a triple where L is a simple non-abelian finite group, $t \in \mathbb{N}$ and $\theta \in \text{Aut}(L)$.
- ▶ u an automorphism of L^t defined by

¹N. Andruskiewitsch and M. Graña, From racks to pointed Hopf algebras, *Adv. Math.* **178** (2003), 177–243.

²D. Joyce, Simple quandles, *J. Algebra* **79** 2 (1982), 307–318. 

Finite simple racks were classified in [AG]¹ and [J]².

We are interested in the (twisted) conjugacy classes of type (L^t, u) :

► (L, t, θ) a triple where L is a simple non-abelian finite group, $t \in \mathbb{N}$ and $\theta \in \text{Aut}(L)$.

► u an automorphism of L^t defined by

$u(\ell_1, \dots, \ell_t) = (\theta(\ell_t), \ell_1, \dots, \ell_{t-1})$ with $\ell_1, \dots, \ell_t \in L$.

¹N. Andruskiewitsch and M. Graña, From racks to pointed Hopf algebras, *Adv. Math.* **178** (2003), 177–243.

²D. Joyce, Simple quandles, *J. Algebra* **79** 2 (1982), 307–318.

Finite simple racks were classified in [AG]¹ and [J]².

We are interested in the (twisted) conjugacy classes of type (L^t, u) :

► (L, t, θ) a triple where L is a simple non-abelian finite group, $t \in \mathbb{N}$ and $\theta \in \text{Aut}(L)$.

► u an automorphism of L^t defined by $u(\ell_1, \dots, \ell_t) = (\theta(\ell_t), \ell_1, \dots, \ell_{t-1})$ with $\ell_1, \dots, \ell_t \in L$.

Definition

¹N. Andruskiewitsch and M. Graña, From racks to pointed Hopf algebras, *Adv. Math.* **178** (2003), 177–243.

²D. Joyce, Simple quandles, *J. Algebra* **79** 2 (1982), 307–318.

Finite simple racks were classified in [AG]¹ and [J]².

We are interested in the (twisted) conjugacy classes of type (L^t, u) :

► (L, t, θ) a triple where L is a simple non-abelian finite group, $t \in \mathbb{N}$ and $\theta \in \text{Aut}(L)$.

► u an automorphism of L^t defined by $u(\ell_1, \dots, \ell_t) = (\theta(\ell_t), \ell_1, \dots, \ell_{t-1})$ with $\ell_1, \dots, \ell_t \in L$.

Definition

If L is a finite group of Lie type, we call a (twisted) conjugacy class of type (L^t, u) a **Lie rack**.

¹N. Andruskiewitsch and M. Graña, From racks to pointed Hopf algebras, *Adv. Math.* **178** (2003), 177–243.

²D. Joyce, Simple quandles, *J. Algebra* **79** 2 (1982), 307–318.

Finite simple racks were classified in [AG]¹ and [J]².

We are interested in the (twisted) conjugacy classes of type (L^t, u) :

► (L, t, θ) a triple where L is a simple non-abelian finite group, $t \in \mathbb{N}$ and $\theta \in \text{Aut}(L)$.

► u an automorphism of L^t defined by $u(\ell_1, \dots, \ell_t) = (\theta(\ell_t), \ell_1, \dots, \ell_{t-1})$ with $\ell_1, \dots, \ell_t \in L$.

Definition

If L is a finite group of Lie type, we call a (twisted) conjugacy class of type (L^t, u) a **Lie rack**.

Thus, non-trivial (twisted) conjugacy classes in simple groups of Lie type are Lie racks.

¹N. Andruskiewitsch and M. Graña, From racks to pointed Hopf algebras, *Adv. Math.* **178** (2003), 177–243.

²D. Joyce, Simple quandles, *J. Algebra* **79** 2 (1982), 307–318.

The rack formulation is more effective for applications to the classification of finite dimensional pointed Hopf algebras over non-abelian groups.

³N. Andruskiewitsch, F. Fantino, M. Graña and L. Vendramin, Finite-dimensional pointed Hopf algebras with alternating groups are trivial. *Ann. Mat. Pura Appl.* (4), to appear.

The rack formulation is more effective for applications to the classification of finite dimensional pointed Hopf algebras over non-abelian groups.

Question

³N. Andruskiewitsch, F. Fantino, M. Graña and L. Vendramin, Finite-dimensional pointed Hopf algebras with alternating groups are trivial. *Ann. Mat. Pura Appl.* (4), to appear.

The rack formulation is more effective for applications to the classification of finite dimensional pointed Hopf algebras over non-abelian groups.

Question

Let X be a finite rack and \mathbf{q} a 2-cocycle with values in $GL(n, \mathbb{C})$.

³N. Andruskiewitsch, F. Fantino, M. Graña and L. Vendramin, Finite-dimensional pointed Hopf algebras with alternating groups are trivial. *Ann. Mat. Pura Appl.* (4), to appear.

The rack formulation is more effective for applications to the classification of finite dimensional pointed Hopf algebras over non-abelian groups.

Question

Let X be a finite rack and \mathbf{q} a 2-cocycle with values in $GL(n, \mathbb{C})$. When is the Nichols algebra $\mathfrak{B}(X, \mathbf{q})$ finite dimensional?

³N. Andruskiewitsch, F. Fantino, M. Graña and L. Vendramin, Finite-dimensional pointed Hopf algebras with alternating groups are trivial. *Ann. Mat. Pura Appl.* (4), to appear.

The rack formulation is more effective for applications to the classification of finite dimensional pointed Hopf algebras over non-abelian groups.

Question

Let X be a finite rack and \mathbf{q} a 2-cocycle with values in $GL(n, \mathbb{C})$. When is the Nichols algebra $\mathfrak{B}(X, \mathbf{q})$ finite dimensional?

An indecomposable finite rack X collapses if for any cocycle \mathbf{q} , $\dim \mathfrak{B}(X, \mathbf{q}) = \infty$.

³N. Andruskiewitsch, F. Fantino, M. Graña and L. Vendramin, Finite-dimensional pointed Hopf algebras with alternating groups are trivial. *Ann. Mat. Pura Appl.* (4), to appear.

The rack formulation is more effective for applications to the classification of finite dimensional pointed Hopf algebras over non-abelian groups.

Question

Let X be a finite rack and \mathbf{q} a 2-cocycle with values in $GL(n, \mathbb{C})$. When is the Nichols algebra $\mathfrak{B}(X, \mathbf{q})$ finite dimensional?

An indecomposable finite rack X collapses if for any cocycle \mathbf{q} , $\dim \mathfrak{B}(X, \mathbf{q}) = \infty$.

Moreover, in [AFGV]³ it was proved

³N. Andruskiewitsch, F. Fantino, M. Graña and L. Vendramin, Finite-dimensional pointed Hopf algebras with alternating groups are trivial. *Ann. Mat. Pura Appl.* (4), to appear.

The rack formulation is more effective for applications to the classification of finite dimensional pointed Hopf algebras over non-abelian groups.

Question

Let X be a finite rack and \mathbf{q} a 2-cocycle with values in $GL(n, \mathbb{C})$. When is the Nichols algebra $\mathfrak{B}(X, \mathbf{q})$ finite dimensional?

An indecomposable finite rack X collapses if for any cocycle \mathbf{q} , $\dim \mathfrak{B}(X, \mathbf{q}) = \infty$.

Moreover, in [AFGV]³ it was proved

Theorem

If X is a finite rack of type D, then X collapses.

³N. Andruskiewitsch, F. Fantino, M. Graña and L. Vendramin, Finite-dimensional pointed Hopf algebras with alternating groups are trivial. *Ann. Mat. Pura Appl.* (4), to appear.

Let p be a prime number, $m \in \mathbb{N}$ and $q = p^m$.

Let p be a prime number, $m \in \mathbb{N}$ and $q = p^m$.

► \mathbb{G} algebraic reductive group defined over the algebraic closure of \mathbb{F}_q

Let p be a prime number, $m \in \mathbb{N}$ and $q = p^m$.

► \mathbb{G} algebraic reductive group defined over the algebraic closure of \mathbb{F}_q and $G = \mathbb{G}(\mathbb{F}_q)$ finite group of \mathbb{F}_q -points.

Let p be a prime number, $m \in \mathbb{N}$ and $q = p^m$.

▶ \mathbb{G} algebraic reductive group defined over the algebraic closure of \mathbb{F}_q and $G = \mathbb{G}(\mathbb{F}_q)$ finite group of \mathbb{F}_q -points.

▶ Take $x \in G$;

Let p be a prime number, $m \in \mathbb{N}$ and $q = p^m$.

- ▶ \mathbb{G} algebraic reductive group defined over the algebraic closure of \mathbb{F}_q and $G = \mathbb{G}(\mathbb{F}_q)$ finite group of \mathbb{F}_q -points.
- ▶ Take $x \in G$; we want to investigate the orbit \mathcal{O}_x^G .

Let p be a prime number, $m \in \mathbb{N}$ and $q = p^m$.

- ▶ \mathbb{G} algebraic reductive group defined over the algebraic closure of \mathbb{F}_q and $G = \mathbb{G}(\mathbb{F}_q)$ finite group of \mathbb{F}_q -points.
- ▶ Take $x \in G$; we want to investigate the orbit \mathcal{O}_x^G .
- ▶ If $x = x_s x_u$ is the Chevalley-Jordan decomposition,

Let p be a prime number, $m \in \mathbb{N}$ and $q = p^m$.

- ▶ \mathbb{G} algebraic reductive group defined over the algebraic closure of \mathbb{F}_q and $G = \mathbb{G}(\mathbb{F}_q)$ finite group of \mathbb{F}_q -points.
- ▶ Take $x \in G$; we want to investigate the orbit \mathcal{O}_x^G .
- ▶ If $x = x_s x_u$ is the Chevalley-Jordan decomposition, then $x_s, x_u \in G$.

Let p be a prime number, $m \in \mathbb{N}$ and $q = p^m$.

- ▶ \mathbb{G} algebraic reductive group defined over the algebraic closure of \mathbb{F}_q and $G = \mathbb{G}(\mathbb{F}_q)$ finite group of \mathbb{F}_q -points.
- ▶ Take $x \in G$; we want to investigate the orbit \mathcal{O}_x^G .
- ▶ If $x = x_s x_u$ is the Chevalley-Jordan decomposition, then $x_s, x_u \in G$.
- ▶ Let $\mathbb{K} = C_{\mathbb{G}}(x_s)$, a reductive subgroup of G .

Let p be a prime number, $m \in \mathbb{N}$ and $q = p^m$.

- ▶ \mathbb{G} algebraic reductive group defined over the algebraic closure of \mathbb{F}_q and $G = \mathbb{G}(\mathbb{F}_q)$ finite group of \mathbb{F}_q -points.
- ▶ Take $x \in G$; we want to investigate the orbit \mathcal{O}_x^G .
- ▶ If $x = x_s x_u$ is the Chevalley-Jordan decomposition, then $x_s, x_u \in G$.
- ▶ Let $\mathbb{K} = C_{\mathbb{G}}(x_s)$, a reductive subgroup of G . Then $K = \mathbb{K} \cap G = C_G(x_s)$.

Let p be a prime number, $m \in \mathbb{N}$ and $q = p^m$.

► \mathbb{G} algebraic reductive group defined over the algebraic closure of \mathbb{F}_q and $G = \mathbb{G}(\mathbb{F}_q)$ finite group of \mathbb{F}_q -points.

► Take $x \in G$; we want to investigate the orbit \mathcal{O}_x^G .

► If $x = x_s x_u$ is the Chevalley-Jordan decomposition, then $x_s, x_u \in G$.

► Let $\mathbb{K} = C_{\mathbb{G}}(x_s)$, a reductive subgroup of G . Then $K = \mathbb{K} \cap G = C_G(x_s)$.

Since $x_u \in K$, $\mathcal{O}_{x_u}^K$ is a subrack of \mathcal{O}_x^G and we can reduce our study to the case when x is either unipotent or semisimple.

Theorem

Assume q is odd and let $u \in SL(n, q)$ be a nontrivial unipotent element. Then \mathcal{O}_u is of type D if and only if $n \neq 2$ or $n = 2$ and q is a square different from 9.

⁴S. Freyre, M. Graña and L. Vendramin, On Nichols algebras over $PGL(2, q)$ and $PSL(2, q)$, *J. Algebra Appl.*, Vol. 9, No. 2 (2010), 195–208.

Theorem

Assume q is odd and let $u \in SL(n, q)$ be a nontrivial unipotent element. Then \mathcal{O}_u is of type D if and only if $n \neq 2$ or $n = 2$ and q is a square different from 9.

Note:

⁴S. Freyre, M. Graña and L. Vendramin, On Nichols algebras over $PGL(2, q)$ and $PSL(2, q)$, *J. Algebra Appl.*, Vol. 9, No. 2 (2010), 195–208.

Theorem

Assume q is odd and let $u \in SL(n, q)$ be a nontrivial unipotent element. Then \mathcal{O}_u is of type D if and only if $n \neq 2$ or $n = 2$ and q is a square different from 9.

Note: Even if the conjugacy classes $\mathcal{O}_u^{SL(2,q)}$ are not of type D for q not a square, by [FGV]⁴, $\mathcal{O}_u^{SL(2,q)}$ collapses for q odd.

⁴S. Freyre, M. Graña and L. Vendramin, On Nichols algebras over $PGL(2, q)$ and $PSL(2, q)$, *J. Algebra Appl.*, Vol. 9, No. 2 (2010), 195–208.

For $\mathbf{a} \in (\mathbb{F}_q^*)^{n-1}$, define

$$r_{\mathbf{a}} = \begin{pmatrix} 1 & a_1 & 0 & \dots & 0 \\ 0 & 1 & a_2 & \dots & 0 \\ \vdots & & \ddots & \ddots & 0 \\ 0 & \dots & \dots & 1 & a_{n-1} \\ 0 & \dots & \dots & 0 & 1 \end{pmatrix}.$$

For $\mathbf{a} \in (\mathbb{F}_q^*)^{n-1}$, define

$$r_{\mathbf{a}} = \begin{pmatrix} 1 & a_1 & 0 & \dots & 0 \\ 0 & 1 & a_2 & \dots & 0 \\ \vdots & & \ddots & \ddots & 0 \\ 0 & \dots & \dots & 1 & a_{n-1} \\ 0 & \dots & \dots & 0 & 1 \end{pmatrix}.$$

A unipotent element $u \in GL(n, q)$ is of type $\lambda = (\lambda_1, \dots, \lambda_k)$ if it is conjugate to the element

$$u = \begin{pmatrix} u_1 & 0 & \dots & 0 \\ 0 & u_2 & \dots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & u_k \end{pmatrix} \quad \text{where } u_i = r_{\mathbf{1}} \in \mathbb{F}_q^{\lambda_i \times \lambda_i}.$$

For $\mathbf{a} \in (\mathbb{F}_q^*)^{n-1}$, define

$$r_{\mathbf{a}} = \begin{pmatrix} 1 & a_1 & 0 & \dots & 0 \\ 0 & 1 & a_2 & \dots & 0 \\ \vdots & & \ddots & \ddots & 0 \\ 0 & \dots & \dots & 1 & a_{n-1} \\ 0 & \dots & \dots & 0 & 1 \end{pmatrix}.$$

A unipotent element $u \in GL(n, q)$ is of type $\lambda = (\lambda_1, \dots, \lambda_k)$ if it is conjugate to the element

$$u = \begin{pmatrix} u_1 & 0 & \dots & 0 \\ 0 & u_2 & \dots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & u_k \end{pmatrix} \quad \text{where } u_i = r_{\mathbf{1}} \in \mathbb{F}_q^{\lambda_i \times \lambda_i}.$$

If $x \in GL(n, q)$ is of type (n) , we call it *regular*.

Lemma 1

Let $u \in SL(n, q)$ be a unipotent element of type $(\lambda_1, \dots, \lambda_k)$ with $u_i \in \mathbb{F}_q^{\lambda_i \times \lambda_i}$.

Lemma 1

Let $u \in SL(n, q)$ be a unipotent element of type $(\lambda_1, \dots, \lambda_k)$ with $u_i \in \mathbb{F}_q^{\lambda_i \times \lambda_i}$. If $\mathcal{O}_{u_i}^{SL(\lambda_i, q)}$ is of type D , then \mathcal{O}_u is of type D .

Lemma 1

Let $u \in SL(n, q)$ be a unipotent element of type $(\lambda_1, \dots, \lambda_k)$ with $u_i \in \mathbb{F}_q^{\lambda_i \times \lambda_i}$. If $\mathcal{O}_{u_i}^{SL(\lambda_i, q)}$ is of type D , then \mathcal{O}_u is of type D .

Sketch of the proof:

Lemma 1

Let $u \in SL(n, q)$ be a unipotent element of type $(\lambda_1, \dots, \lambda_k)$ with $u_i \in \mathbb{F}_q^{\lambda_i \times \lambda_i}$. If $\mathcal{O}_{u_i}^{SL(\lambda_i, q)}$ is of type D , then \mathcal{O}_u is of type D .

Sketch of the proof: Assume $\mathcal{O}_{u_1}^{SL(\lambda_1, q)}$ is of type D .

Lemma 1

Let $u \in SL(n, q)$ be a unipotent element of type $(\lambda_1, \dots, \lambda_k)$ with $u_i \in \mathbb{F}_q^{\lambda_i \times \lambda_i}$. If $\mathcal{O}_{u_i}^{SL(\lambda_i, q)}$ is of type D , then \mathcal{O}_u is of type D .

Sketch of the proof: Assume $\mathcal{O}_{u_1}^{SL(\lambda_1, q)}$ is of type D .

Let H be the image of the natural embedding of $SL(\lambda_1, q) \times \dots \times SL(\lambda_k, q)$ into $SL(n, q)$ containing u .

Lemma 1

Let $u \in SL(n, q)$ be a unipotent element of type $(\lambda_1, \dots, \lambda_k)$ with $u_i \in \mathbb{F}_q^{\lambda_i \times \lambda_i}$. If $\mathcal{O}_{u_i}^{SL(\lambda_i, q)}$ is of type D , then \mathcal{O}_u is of type D .

Sketch of the proof: Assume $\mathcal{O}_{u_1}^{SL(\lambda_1, q)}$ is of type D .

Let H be the image of the natural embedding of $SL(\lambda_1, q) \times \dots \times SL(\lambda_k, q)$ into $SL(n, q)$ containing u .

Then \mathcal{O}_u^H is a subrack of $\mathcal{O}_u^{SL(n, q)}$ and the projection of H onto its first factor induces a rack epimorphism of \mathcal{O}_u^H onto $\mathcal{O}_{u_1}^{SL(\lambda_1, q)}$.

Lemma 2

Assume that $n > 2$ and q is odd.

Lemma 2

Assume that $n > 2$ and q is odd. Then the conjugacy class of a unipotent regular element in $SL(n, q)$ is of type D .

Lemma 2

Assume that $n > 2$ and q is odd. Then the conjugacy class of a unipotent regular element in $SL(n, q)$ is of type D .

Corollary

Lemma 2

Assume that $n > 2$ and q is odd. Then the conjugacy class of a unipotent regular element in $SL(n, q)$ is of type D .

Corollary

Assume q is odd and let $u \in SL(n, q)$ be a unipotent element of type $(\lambda_1, \dots, \lambda_k)$.

Lemma 2

Assume that $n > 2$ and q is odd. Then the conjugacy class of a unipotent regular element in $SL(n, q)$ is of type D .

Corollary

Assume q is odd and let $u \in SL(n, q)$ be a unipotent element of type $(\lambda_1, \dots, \lambda_k)$. If $\lambda_1 > 2$, then the conjugacy class \mathcal{O}_u is of type D .

Lemma 3

Assume q is odd. Let $u \in SL(n, q)$ be a unipotent element of type $(2, 2)$ or $(2, 1)$. Then the conjugacy class \mathcal{O}_u is of type D.

Lemma 3

Assume q is odd. Let $u \in SL(n, q)$ be a unipotent element of type $(2, 2)$ or $(2, 1)$. Then the conjugacy class \mathcal{O}_u is of type D.

Case $(2, 2)$ in $SL(4, 3)$ done with GAP.

Lemma 3

Assume q is odd. Let $u \in SL(n, q)$ be a unipotent element of type $(2, 2)$ or $(2, 1)$. Then the conjugacy class \mathcal{O}_u is of type D.

Case $(2, 2)$ in $SL(4, 3)$ done with GAP.

Lemma 4

Assume $q = p^m$ is odd. Then any non-trivial unipotent conjugacy class in $SL(2, q)$ is of type D if and only if q is a square different from 9.

All results can be used to prove the main theorem for $PSL(n, q)$ and $PGL(n, q)$ since the rack structures are not deformed by taking the quotients.

All results can be used to prove the main theorem for $PSL(n, q)$ and $PGL(n, q)$ since the rack structures are not deformed by taking the quotients.

Assume \mathbb{G} is a simple algebraic group and $F : \mathbb{G} \rightarrow \mathbb{G}$ is a Steinberg automorphism which is \mathbb{F}_q -split.

All results can be used to prove the main theorem for $PSL(n, q)$ and $PGL(n, q)$ since the rack structures are not deformed by taking the quotients.

Assume \mathbb{G} is a simple algebraic group and $F : \mathbb{G} \rightarrow \mathbb{G}$ is a Steinberg automorphism which is \mathbb{F}_q -split.

Translating the techniques used for $SL(n, q)$ using Lie theory one can extend the result to other types, for example

All results can be used to prove the main theorem for $PSL(n, q)$ and $PGL(n, q)$ since the rack structures are not deformed by taking the quotients.

Assume \mathbb{G} is a simple algebraic group and $F : \mathbb{G} \rightarrow \mathbb{G}$ is a Steinberg automorphism which is \mathbb{F}_q -split.

Translating the techniques used for $SL(n, q)$ using Lie theory one can extend the result to other types, for example

If the Dynkin diagram of \mathbb{G} has at least a component of rank greater than 1,

All results can be used to prove the main theorem for $PSL(n, q)$ and $PGL(n, q)$ since the rack structures are not deformed by taking the quotients.

Assume \mathbb{G} is a simple algebraic group and $F : \mathbb{G} \rightarrow \mathbb{G}$ is a Steinberg automorphism which is \mathbb{F}_q -split.

Translating the techniques used for $SL(n, q)$ using Lie theory one can extend the result to other types, for example

If the Dynkin diagram of \mathbb{G} has at least a component of rank greater than 1, then every regular unipotent class of \mathbb{G}^F is of type D .

THANK YOU

MERCI

GRACIAS

GRAZZIE

OBRIGADO

DANKE