

# Knörr lattices for elementary abelian $p$ -groups

Michael Geline  
Northern Illinois University  
with Geoffrey Robinson

Regina, SK  
June 2012

# Blocks, Characters

Consider:

$(K/\mathbb{Q}_p, R, k) = p$ -modular system

# Blocks, Characters

Consider:

$(K/\mathbb{Q}_p, R, k) = p$ -modular system

$$RG \rightarrow KG$$

$$Z(RG) \rightarrow Z(KG)$$

# Blocks, Characters

Consider:

$(K/\mathbb{Q}_p, R, k) = p$ -modular system

$$RG \rightarrow KG$$

$$Z(RG) \rightarrow Z(KG)$$

A primitive idempotent in  $Z(RG)$  becomes a sum of primitive idempotents in  $Z(KG)$ .

Obtain: Blocks of irreducible characters of  $G$ .

# Blocks, Characters

Consider:

$(K/\mathbb{Q}_p, R, k) = p$ -modular system

$$RG \rightarrow KG$$

$$Z(RG) \rightarrow Z(KG)$$

A primitive idempotent in  $Z(RG)$  becomes a sum of primitive idempotents in  $Z(KG)$ .

Obtain: Blocks of irreducible characters of  $G$ .

- ▶ What do irreducibles in same block have in common?
- ▶ How many irreducibles appear in a given block?
- ▶ In what terms can we expect answers to questions like this?

# Defect Groups

If  $H \subseteq G$ , there is natural map

$$\mathrm{Tr}_H^G : (RG)^H \rightarrow (RG)^G = Z(RG)$$

$$\alpha \mapsto \sum_{g \in G/H} g^{-1} \alpha g$$

# Defect Groups

If  $H \subseteq G$ , there is natural map

$$\mathrm{Tr}_H^G : (RG)^H \rightarrow (RG)^G = Z(RG)$$

$$\alpha \mapsto \sum_{g \in G/H} g^{-1} \alpha g$$

If  $e$  is a block idempotent,  $D$  is called a *defect group* of  $e$  if  $D$  is minimal such that

$$e = \mathrm{Tr}_D^G(\beta) \text{ for some } \beta \in (RG)^D.$$

# Defect Groups

If  $H \subseteq G$ , there is natural map

$$\mathrm{Tr}_H^G : (RG)^H \rightarrow (RG)^G = Z(RG)$$

$$\alpha \mapsto \sum_{g \in G/H} g^{-1} \alpha g$$

If  $e$  is a block idempotent,  $D$  is called a *defect group* of  $e$  if  $D$  is minimal such that

$$e = \mathrm{Tr}_D^G(\beta) \text{ for some } \beta \in (RG)^D.$$

Fact:  $D$  is determined up to conjugacy by  $e$ .

# Heights

Let  $\chi$  be an irreducible character associated to  $e$ :

# Heights

Let  $\chi$  be an irreducible character associated to  $e$ :

$$\chi(1) = \chi(e) = \chi(\mathrm{Tr}_D^G(\beta)) = |G : D|\chi(\beta).$$

# Heights

Let  $\chi$  be an irreducible character associated to  $e$ :

$$\chi(1) = \chi(e) = \chi(\mathrm{Tr}_D^G(\beta)) = |G : D|\chi(\beta).$$

Let  $|G|_p = p^a$ , and  $|D| = p^d$ .

# Heights

Let  $\chi$  be an irreducible character associated to  $e$ :

$$\chi(1) = \chi(e) = \chi(\mathrm{Tr}_D^G(\beta)) = |G : D|\chi(\beta).$$

Let  $|G|_p = p^a$ , and  $|D| = p^d$ . Then

$$\chi(1)_p = p^{a-d+h}$$

and  $h$  is called the *height* of  $\chi$ .

# Heights

Let  $\chi$  be an irreducible character associated to  $e$ :

$$\chi(1) = \chi(e) = \chi(\mathrm{Tr}_D^G(\beta)) = |G : D|\chi(\beta).$$

Let  $|G|_p = p^a$ , and  $|D| = p^d$ . Then

$$\chi(1)_p = p^{a-d+h}$$

and  $h$  is called the *height* of  $\chi$ .

Brauer's Height Zero Conjecture:

$h = 0$  for all  $\chi$  in  $e$  if and only if  $D$  is abelian.

# Lattices, Vertices, Sources

An *RG-lattice* is an *RG*-module that is free of finite rank over  $R$ .

# Lattices, Vertices, Sources

An *RG-lattice* is an *RG*-module that is free of finite rank over  $R$ .

- ▶ Let  $\hat{M}$  be a *KG*-module affording  $\chi$ .

# Lattices, Vertices, Sources

An *RG-lattice* is an *RG*-module that is free of finite rank over  $R$ .

- ▶ Let  $\hat{M}$  be a *KG*-module affording  $\chi$ .
- ▶ Use a  $K$ -basis of  $\hat{M}$  to generate an *RG*-lattice,  $M$ .

# Lattices, Vertices, Sources

An *RG-lattice* is an *RG*-module that is free of finite rank over *R*.

- ▶ Let  $\hat{M}$  be a *KG*-module affording  $\chi$ .
- ▶ Use a *K*-basis of  $\hat{M}$  to generate an *RG*-lattice,  $M$ .
- ▶ Let  $V$  be subgroup minimal such that

$$\mathrm{Ind}_V^G(S) \cong M \oplus \dots \text{ for some } RV\text{-lattice } S$$

# Lattices, Vertices, Sources

An *RG-lattice* is an *RG*-module that is free of finite rank over *R*.

- ▶ Let  $\hat{M}$  be a *KG*-module affording  $\chi$ .
- ▶ Use a *K*-basis of  $\hat{M}$  to generate an *RG*-lattice,  $M$ .
- ▶ Let  $V$  be subgroup minimal such that

$$\mathrm{Ind}_V^G(S) \cong M \oplus \dots \text{ for some } RV\text{-lattice } S$$

Green:

- ▶  $V$  is determined up to conjugacy by (isomorphism class of)  $M$ .

# Lattices, Vertices, Sources

An *RG-lattice* is an *RG*-module that is free of finite rank over *R*.

- ▶ Let  $\hat{M}$  be a *KG*-module affording  $\chi$ .
- ▶ Use a *K*-basis of  $\hat{M}$  to generate an *RG*-lattice,  $M$ .
- ▶ Let  $V$  be subgroup minimal such that

$$\mathrm{Ind}_V^G(S) \cong M \oplus \dots \text{ for some } RV\text{-lattice } S$$

Green:

- ▶  $V$  is determined up to conjugacy by (isomorphism class of)  $M$ .
- ▶ The isomorphism class of  $S$  is determined up to conjugacy by  $N_G(V)$ .

# Lattices, Vertices, Sources

An *RG-lattice* is an *RG*-module that is free of finite rank over *R*.

- ▶ Let  $\hat{M}$  be a *KG*-module affording  $\chi$ .
- ▶ Use a *K*-basis of  $\hat{M}$  to generate an *RG*-lattice,  $M$ .
- ▶ Let  $V$  be subgroup minimal such that

$$\mathrm{Ind}_V^G(S) \cong M \oplus \dots \text{ for some } RV\text{-lattice } S$$

Green:

- ▶  $V$  is determined up to conjugacy by (isomorphism class of)  $M$ .
- ▶ The isomorphism class of  $S$  is determined up to conjugacy by  $N_G(V)$ .
- ▶ Can choose  $V \subseteq D$ .

## Knörr 1978

Again:  $V$  is minimal subgroup such that

$$\mathrm{Ind}_V^G(S) \cong M \oplus \dots \text{ for some } RV\text{-lattice } S.$$

# Knörr 1978

Again:  $V$  is minimal subgroup such that

$$\mathrm{Ind}_V^G(S) \cong M \oplus \dots \text{ for some } RV\text{-lattice } S.$$

## Theorem

*Let  $\chi$  be irreducible, and  $M$ ,  $V$ ,  $D$ , and  $S$  as above. Then*

$$h = \text{height of } \chi = \nu|D : V| + \nu(\mathrm{rank}(S)).$$

## Knörr 1978

Again:  $V$  is minimal subgroup such that

$$\mathrm{Ind}_V^G(S) \cong M \oplus \dots \text{ for some } RV\text{-lattice } S.$$

### Theorem

*Let  $\chi$  be irreducible, and  $M$ ,  $V$ ,  $D$ , and  $S$  as above. Then*

$$h = \text{height of } \chi = \nu|D : V| + \nu(\mathrm{rank}(S)).$$

### Theorem

*With the same assumptions*

$$C_D(V) \subseteq V.$$

*So if  $D$  is abelian, then  $V = D$ .*

# Knörr 1988

## Definition

Let  $L$  be an indecomposable  $RH$ -lattice. Let  $A = \text{End}_{RH}(L)$ .

# Knörr 1988

## Definition

Let  $L$  be an indecomposable  $RH$ -lattice. Let  $A = \text{End}_{RH}(L)$ .  
Assume that  $A = R \cdot \text{Id}_L + J(A)$ .

# Knörr 1988

## Definition

Let  $L$  be an indecomposable  $RH$ -lattice. Let  $A = \text{End}_{RH}(L)$ .

Assume that  $A = R \cdot \text{Id}_L + J(A)$ .

Then  $L$  is a *Knörr lattice* if

# Knörr 1988

## Definition

Let  $L$  be an indecomposable  $RH$ -lattice. Let  $A = \text{End}_{RH}(L)$ .

Assume that  $A = R \cdot \text{Id}_L + J(A)$ .

Then  $L$  is a *Knörr lattice* if

$$\text{trace}(J(A)) \neq \text{trace}(A) \subseteq R$$

# Knörr 1988

## Definition

Let  $L$  be an indecomposable  $RH$ -lattice. Let  $A = \text{End}_{RH}(L)$ .

Assume that  $A = R \cdot \text{Id}_L + J(A)$ .

Then  $L$  is a *Knörr lattice* if

$$\text{trace}(J(A)) \neq \text{trace}(A) \subseteq R$$

Examples:

- ▶  $L$  affords an absolutely irreducible character.
- ▶  $\text{rank}(L)$  is not divisible by  $p$ .

# Knörr 1988

## Definition

Let  $L$  be an indecomposable  $RH$ -lattice. Let  $A = \text{End}_{RH}(L)$ .

Assume that  $A = R \cdot \text{Id}_L + J(A)$ .

Then  $L$  is a *Knörr lattice* if

$$\text{trace}(J(A)) \neq \text{trace}(A) \subseteq R$$

Examples:

- ▶  $L$  affords an absolutely irreducible character.
- ▶  $\text{rank}(L)$  is not divisible by  $p$ .

## Theorem

*The source of a Knörr lattice is itself a Knörr lattice.*

# Knörr 1988

## Definition

Let  $L$  be an indecomposable  $RH$ -lattice. Let  $A = \text{End}_{RH}(L)$ .

Assume that  $A = R \cdot \text{Id}_L + J(A)$ .

Then  $L$  is a *Knörr lattice* if

$$\text{trace}(J(A)) \neq \text{trace}(A) \subseteq R$$

Examples:

- ▶  $L$  affords an absolutely irreducible character.
- ▶  $\text{rank}(L)$  is not divisible by  $p$ .

## Theorem

*The source of a Knörr lattice is itself a Knörr lattice.*

Question:

- ▶ Can an abelian  $p$ -group have a Knörr lattice of rank divisible by  $p$ ?

## Knörr 1988, example

Let  $R = \mathbb{Z}_2[2^{1/3}]$  and  $D = C_2 \times C_2 \times C_2 \times C_2$ .

THEN: there is a rank 6 Knörr  $RD$ -lattice (gives four explicit  $6 \times 6$  matrices).

But...

The group  $\text{Gal}(\mathbb{Q}_2(2^{1/3}, \zeta_3)/\mathbb{Q}_2)$  is not abelian, yet  $\chi$  can be afforded by a  $KG$  module in which  $K/\mathbb{Q}_2$  is cyclotomic!

But...

The group  $\text{Gal}(\mathbb{Q}_2(2^{1/3}, \zeta_3)/\mathbb{Q}_2)$  is not abelian, yet  $\chi$  can be afforded by a  $KG$  module in which  $K/\mathbb{Q}_2$  is cyclotomic!

Conclude: Knörr's example does not count!

Conclude: Knörr's strategy is still alive!

But...

The group  $\text{Gal}(\mathbb{Q}_2(2^{1/3}, \zeta_3)/\mathbb{Q}_2)$  is not abelian, yet  $\chi$  can be afforded by a  $KG$  module in which  $K/\mathbb{Q}_2$  is cyclotomic!

Conclude: Knörr's example does not count!

Conclude: Knörr's strategy is still alive!

Specifically, if a positive height irreducible character of some finite group in a 2-block with elementary abelian 16 defect group exists, THEN a Knörr lattice of even rank *defined over a cyclotomic extension* of  $\mathbb{Z}_2$  exists. And Knörr does not display such a lattice.

# A better splitting field

Facts:

- ▶ Associated to  $\chi$  is a division algebra component  $\Delta$  of the group algebra  $\mathbb{Q}_p G$ .

# A better splitting field

Facts:

- ▶ Associated to  $\chi$  is a division algebra component  $\Delta$  of the group algebra  $\mathbb{Q}_p G$ .
- ▶  $\mathbb{Q}_p \subseteq Z(\Delta) \cong \mathbb{Q}_p(\chi) \subseteq K' \subseteq \Delta$ , where  $K'$  is a maximal subfield.

# A better splitting field

Facts:

- ▶ Associated to  $\chi$  is a division algebra component  $\Delta$  of the group algebra  $\mathbb{Q}_p G$ .
- ▶  $\mathbb{Q}_p \subseteq Z(\Delta) \cong \mathbb{Q}_p(\chi) \subseteq K' \subseteq \Delta$ , where  $K'$  is a maximal subfield.
- ▶ There is a  $K'G$ -module affording  $\chi$  for each choice of  $K'$ .

# A better splitting field

Facts:

- ▶ Associated to  $\chi$  is a division algebra component  $\Delta$  of the group algebra  $\mathbb{Q}_p G$ .
- ▶  $\mathbb{Q}_p \subseteq Z(\Delta) \cong \mathbb{Q}_p(\chi) \subseteq K' \subseteq \Delta$ , where  $K'$  is a maximal subfield.
- ▶ There is a  $K'G$ -module affording  $\chi$  for each choice of  $K'$ .
- ▶ There is a  $K'$  which is unramified over  $Z(\Delta) \cong \mathbb{Q}_p(\chi)$ .

# A better splitting field

Facts:

- ▶ Associated to  $\chi$  is a division algebra component  $\Delta$  of the group algebra  $\mathbb{Q}_p G$ .
- ▶  $\mathbb{Q}_p \subseteq Z(\Delta) \cong \mathbb{Q}_p(\chi) \subseteq K' \subseteq \Delta$ , where  $K'$  is a maximal subfield.
- ▶ There is a  $K'G$ -module affording  $\chi$  for each choice of  $K'$ .
- ▶ There is a  $K'$  which is unramified over  $Z(\Delta) \cong \mathbb{Q}_p(\chi)$ .

Also (Green/Brauer)

- ▶ If  $g \in G$  and  $g_p$  is not conjugate to an element of  $D$ , then  $\chi(g) = 0$ .

# A better splitting field

## Lemma

Let

$$K = \mathbb{Q}_p(\text{all } p' \text{ roots of unity, } \zeta_{p^m})$$

where  $p^m$  is the exponent of  $D$ . Let  $R$  be the valuation ring of  $K$ .  
Then  $\chi$  is afforded by an RG-lattice.

# A better splitting field

## Lemma

Let

$$K = \mathbb{Q}_p(\text{all } p' \text{ roots of unity}, \zeta_{p^m})$$

where  $p^m$  is the exponent of  $D$ . Let  $R$  be the valuation ring of  $K$ . Then  $\chi$  is afforded by an  $RG$ -lattice.

The point:

- ▶ A uniformizer for above  $R$  is  $1 - \zeta_{p^m}$ , and this *shrinks* the category of  $RD$ -lattices.

# A better splitting field

## Lemma

Let

$$K = \mathbb{Q}_p(\text{all } p' \text{ roots of unity}, \zeta_{p^m})$$

where  $p^m$  is the exponent of  $D$ . Let  $R$  be the valuation ring of  $K$ . Then  $\chi$  is afforded by an  $RG$ -lattice.

The point:

- ▶ A uniformizer for above  $R$  is  $1 - \zeta_{p^m}$ , and this *shrinks* the category of  $RD$ -lattices.

The hope:

- ▶ Knörr lattices (with rank divisible by  $p$ ) for abelian  $p$ -groups with above uniformizer might be understandable.

# First results

## Theorem

*Let  $D$  be elementary abelian of order  $2^d$  with  $d \geq 2$ . Let  $M$  be a Knörr RD-lattice. Then*

$$\nu(\text{rank}(M)) \leq d - 2.$$

# First results

## Theorem

*Let  $D$  be elementary abelian of order  $2^d$  with  $d \geq 2$ . Let  $M$  be a Knörr  $RD$ -lattice. Then*

$$\nu(\text{rank}(M)) \leq d - 2.$$

Idea: If  $g \in D$ , then  $\chi_M(g) \neq 0$ . This forces  $M_{\langle g \rangle}$  to have a rank 1 summand. Transfer the projection to an  $RD$ -endomorphism of  $M$ .

# First results

## Theorem

*Let  $D$  be elementary abelian of order  $2^d$  with  $d \geq 2$ . Let  $M$  be a Knörr  $RD$ -lattice. Then*

$$\nu(\text{rank}(M)) \leq d - 2.$$

Idea: If  $g \in D$ , then  $\chi_M(g) \neq 0$ . This forces  $M_{\langle g \rangle}$  to have a rank 1 summand. Transfer the projection to an  $RD$ -endomorphism of  $M$ .

## Theorem

*Let  $D$  be elementary abelian of order 8, and assume  $R/\mathbb{Z}_2$  is unramified. Then an even rank Knörr  $RD$ -lattice must have rank at least 14.*

## Idea for a rank 6 lattice with $D \cong C_2 \times C_2 \times C_2$

- ▶ Let  $H$  be a maximal subgroup of  $D$ . If  $M_H$  has an odd-rank summand, consider the transfer of the projection to an  $RD$ -endomorphism of  $M$ . We conclude that  $M_H$  has no odd-rank summands.

## Idea for a rank 6 lattice with $D \cong C_2 \times C_2 \times C_2$

- ▶ Let  $H$  be a maximal subgroup of  $D$ . If  $M_H$  has an odd-rank summand, consider the transfer of the projection to an  $RD$ -endomorphism of  $M$ . We conclude that  $M_H$  has no odd-rank summands.
- ▶ If  $M_H$  has a projective summand, then  $M_H \cong P \oplus 2$ . This forces  $\chi_M(g) = 0$  for some  $g \in H$ .

## Idea for a rank 6 lattice with $D \cong C_2 \times C_2 \times C_2$

- ▶ Let  $H$  be a maximal subgroup of  $D$ . If  $M_H$  has an odd-rank summand, consider the transfer of the projection to an  $RD$ -endomorphism of  $M$ . We conclude that  $M_H$  has no odd-rank summands.
- ▶ If  $M_H$  has a projective summand, then  $M_H \cong P \oplus 2$ . This forces  $\chi_M(g) = 0$  for some  $g \in H$ .
- ▶ If  $M$  has no projective summands, then a result of MCR Butler shows that (using fact that  $R/(2) = k$ )

$$2M \subseteq (M_H)_{ss} \subseteq M.$$

Now  $(M_H)_{ss}$  is an  $RD$ -lattice with a rank-1 summand. Consider multiplication by 2 followed by projection.

## Cyclic examples

A *cyclic* lattice is a quotient of the regular lattice  $M \cong RD/I$ , for some pure ideal  $I \subseteq RD$ .

## Cyclic examples

A *cyclic* lattice is a quotient of the regular lattice  $M \cong RD/I$ , for some pure ideal  $I \subseteq RD$ .

Assume now that  $D$  is an abelian  $p$ -group.

# Cyclic examples

A *cyclic* lattice is a quotient of the regular lattice  $M \cong RD/I$ , for some pure ideal  $I \subseteq RD$ .

Assume now that  $D$  is an abelian  $p$ -group.

- ▶ Cyclic  $RD$ -lattices are determined by the characters they afford.

# Cyclic examples

A *cyclic* lattice is a quotient of the regular lattice  $M \cong RD/I$ , for some pure ideal  $I \subseteq RD$ .

Assume now that  $D$  is an abelian  $p$ -group.

- ▶ Cyclic  $RD$ -lattices are determined by the characters they afford.
- ▶ If  $R$  contains  $\zeta_{p^m}$ , and  $\chi$  is a multiplicity-free character of  $D$ , there will be a unique cyclic  $RD$ -lattice affording  $\chi$ .

# Cyclic examples

A *cyclic* lattice is a quotient of the regular lattice  $M \cong RD/I$ , for some pure ideal  $I \subseteq RD$ .

Assume now that  $D$  is an abelian  $p$ -group.

- ▶ Cyclic  $RD$ -lattices are determined by the characters they afford.
- ▶ If  $R$  contains  $\zeta_{p^m}$ , and  $\chi$  is a multiplicity-free character of  $D$ , there will be a unique cyclic  $RD$ -lattice affording  $\chi$ .

## Lemma

Let  $M$  be a cyclic  $RD$ -lattice with  $\text{rank}(M)_p = p^h$ . Then  $M$  is a Knörr lattice if and only if

$$\chi_M(g) \equiv \chi_M(1) \pmod{\pi p^h}$$

for all  $g \in D$ .

# Cyclic examples

## Lemma

*Let  $g$  have order  $p$ , and let  $\chi$  be a character of  $D$ . Then*

$$\chi(g) \equiv \chi(1) \pmod{\pi p^h}$$

*if and only if  $[\chi_{\langle g \rangle}, \lambda] \equiv 0 \pmod{p^h}$  for all  $\lambda \in \langle \hat{g} \rangle$ .*

# Cyclic examples

## Lemma

*Let  $g$  have order  $p$ , and let  $\chi$  be a character of  $D$ . Then*

$$\chi(g) \equiv \chi(1) \pmod{\pi p^h}$$

*if and only if  $[\chi_{\langle g \rangle}, \lambda] \equiv 0 \pmod{p^h}$  for all  $\lambda \in \langle \hat{g} \rangle$ .*

If  $D$  is an elementary abelian 2-group, and  $h = 1$ , this is equivalent to saying that

$$\det(\chi)(g) = 1.$$

# Cyclic examples

## Lemma

Let  $g$  have order  $p$ , and let  $\chi$  be a character of  $D$ . Then

$$\chi(g) \equiv \chi(1) \pmod{\pi p^h}$$

if and only if  $[\chi_{\langle g \rangle}, \lambda] \equiv 0 \pmod{p^h}$  for all  $\lambda \in \langle \hat{g} \rangle$ .

If  $D$  is an elementary abelian 2-group, and  $h = 1$ , this is equivalent to saying that

$$\det(\chi)(g) = 1.$$

For example

$$\chi = 1_D + \lambda_1 + \lambda_2 + \lambda_3 + \lambda_4 + \lambda_1\lambda_2\lambda_3\lambda_4$$

is the character afforded by a cyclic rank-6 lattice for the e.a. group of order 16. Here  $\hat{D} = \langle \lambda_i \mid 1 \leq i \leq 4 \rangle$ .

## Subsets of $\mathbb{F}_p^d$

Assume now that  $D$  is elementary abelian. If  $\chi$  is multiplicity-free, then  $\chi$  can be thought of as a subset of the  $\mathbb{F}_p$  vector space  $\hat{D}$ .

$$\chi \subseteq \hat{D}$$

## Subsets of $\mathbb{F}_p^d$

Assume now that  $D$  is elementary abelian. If  $\chi$  is multiplicity-free, then  $\chi$  can be thought of as a subset of the  $\mathbb{F}_p$  vector space  $\hat{D}$ .

$$\chi \subseteq \hat{D}$$

### Lemma

*Let  $\chi$  be multiplicity-free character of the e.a.  $p$ -group  $D$ . Say  $\chi(1)_p = p^h$ . Then  $\chi$  is the character of a cyclic Knörr RD-lattice if and only if*

$$|\chi \cap W| \equiv 0 \pmod{p^h}$$

*for every codimension 1 hyperplane  $W$  in  $\hat{D}$ .*

## Subsets of $\mathbb{F}_p^d$

Assume now that  $D$  is elementary abelian. If  $\chi$  is multiplicity-free, then  $\chi$  can be thought of as a subset of the  $\mathbb{F}_p$  vector space  $\hat{D}$ .

$$\chi \subseteq \hat{D}$$

### Lemma

*Let  $\chi$  be multiplicity-free character of the e.a.  $p$ -group  $D$ . Say  $\chi(1)_p = p^h$ . Then  $\chi$  is the character of a cyclic Knörr RD-lattice if and only if*

$$|\chi \cap W| \equiv 0 \pmod{p^h}$$

*for every codimension 1 hyperplane  $W$  in  $\hat{D}$ .*

### Theorem

*If  $|D| = p^d$ , then  $h \leq \frac{d}{2}$ . This bound is tight.*

# Open Problems

- ▶ Does the e.a. group of order  $p^3$  have a height 1 cyclic Knörr lattice for all  $p \geq 5$ ?

# Open Problems

- ▶ Does the e.a. group of order  $p^3$  have a height 1cyclic Knörr lattice for all  $p \geq 5$ ?
- ▶ Count the height 1cyclic Knörr lattices for e.a.  $p^3$  as a function of  $p$ .

# Open Problems

- ▶ Does the e.a. group of order  $p^3$  have a height 1cyclic Knörr lattice for all  $p \geq 5$ ?
- ▶ Count the height 1cyclic Knörr lattices for e.a.  $p^3$  as a function of  $p$ .
- ▶ If  $R$  has uniformizer as above, does there exist a positive height Knörr lattice which is not a syzygy of a cyclic lattice?

# Open Problems

- ▶ Does the e.a. group of order  $p^3$  have a height 1cyclic Knörr lattice for all  $p \geq 5$ ?
- ▶ Count the height 1cyclic Knörr lattices for e.a.  $p^3$  as a function of  $p$ .
- ▶ If  $R$  has uniformizer as above, does there exist a positive height Knörr lattice which is not a syzygy of a cyclic lattice?
- ▶ If  $R/\mathbb{Z}_2$  is unramified, does the e.a. group of order 8 have a height 1 Knörr lattice at all?

# Open Problems

- ▶ Does the e.a. group of order  $p^3$  have a height 1cyclic Knörr lattice for all  $p \geq 5$ ?
- ▶ Count the height 1cyclic Knörr lattices for e.a.  $p^3$  as a function of  $p$ .
- ▶ If  $R$  has uniformizer as above, does there exist a positive height Knörr lattice which is not a syzygy of a cyclic lattice?
- ▶ If  $R/\mathbb{Z}_2$  is unramified, does the e.a. group of order 8 have a height 1 Knörr lattice at all?
- ▶ If  $\chi$  is an irreducible character of  $G$ , is  $\chi$  necessarily afforded by a lattice with a cyclic source?

# Open Problems

- ▶ Does the e.a. group of order  $p^3$  have a height 1 cyclic Knörr lattice for all  $p \geq 5$ ?
- ▶ Count the height 1 cyclic Knörr lattices for e.a.  $p^3$  as a function of  $p$ .
- ▶ If  $R$  has uniformizer as above, does there exist a positive height Knörr lattice which is not a syzygy of a cyclic lattice?
- ▶ If  $R/\mathbb{Z}_2$  is unramified, does the e.a. group of order 8 have a height 1 Knörr lattice at all?
- ▶ If  $\chi$  is an irreducible character of  $G$ , is  $\chi$  necessarily afforded by a lattice with a cyclic source?
- ▶ Examples:  $G = S_n$  etc.

# Open Problems

- ▶ Does the e.a. group of order  $p^3$  have a height 1 cyclic Knörr lattice for all  $p \geq 5$ ?
- ▶ Count the height 1 cyclic Knörr lattices for e.a.  $p^3$  as a function of  $p$ .
- ▶ If  $R$  has uniformizer as above, does there exist a positive height Knörr lattice which is not a syzygy of a cyclic lattice?
- ▶ If  $R/\mathbb{Z}_2$  is unramified, does the e.a. group of order 8 have a height 1 Knörr lattice at all?
- ▶ If  $\chi$  is an irreducible character of  $G$ , is  $\chi$  necessarily afforded by a lattice with a cyclic source?
- ▶ Examples:  $G = S_n$  etc.
- ▶ Powers of radical.

# Open Problems

- ▶ Does the e.a. group of order  $p^3$  have a height 1 cyclic Knörr lattice for all  $p \geq 5$ ?
- ▶ Count the height 1 cyclic Knörr lattices for e.a.  $p^3$  as a function of  $p$ .
- ▶ If  $R$  has uniformizer as above, does there exist a positive height Knörr lattice which is not a syzygy of a cyclic lattice?
- ▶ If  $R/\mathbb{Z}_2$  is unramified, does the e.a. group of order 8 have a height 1 Knörr lattice at all?
- ▶ If  $\chi$  is an irreducible character of  $G$ , is  $\chi$  necessarily afforded by a lattice with a cyclic source?
- ▶ Examples:  $G = S_n$  etc.
- ▶ Powers of radical.
- ▶ Stabilizer of source in  $N_G(D)$ ?