

Weil Representation of the groups $GL_{*}^{\varepsilon}(2, A)$ and $SL_{*}^{\varepsilon}(2, A)$

José Pantoja

Pontificia Universidad Católica de Valparaíso
Valparaíso
Chile

CMS 2012
Regina, Canada
June 2012

partially supported by Fondecyt Grant 1120578

The groups $GL_{*}^{\varepsilon}(2, A)$ and $SL_{*}^{\varepsilon}(2, A)$

Basic Idea

To look at higher rank groups as non commutative analogues of related lower rank groups

- ▶ Example: To look at the symplectic similitude group $GSp(2n, F)$ in $2n$ variables over a field F as a group $GL(2)$ with coefficients in the matrix ring over F , satisfying suitable commutation relations which involve the transpose map*
- ▶ To extend to higher rank groups, successful methods used for lower rank groups. In particular, having presentations for higher rank classical groups as non commutative versions of known presentations for the lower rank case. These presentations can be used in constructing linear representations for higher rank groups (generalized Weil representations, for example)

The groups $GL_{*}^{\varepsilon}(2, A)$ and $SL_{*}^{\varepsilon}(2, A)$

Basic Idea

To look at higher rank groups as non commutative analogues of related lower rank groups

- ▶ Example: To look at the symplectic similitude group $GSp(2n, F)$ in $2n$ variables over a field F as a group $GL(2)$ with coefficients in the matrix ring over F , satisfying suitable commutation relations which involve the transpose map*
- ▶ To extend to higher rank groups, successful methods used for lower rank groups. In particular, having presentations for higher rank classical groups as non commutative versions of known presentations for the lower rank case. These presentations can be used in constructing linear representations for higher rank groups (generalized Weil representations, for example)

The groups $GL_{*}^{\varepsilon}(2, A)$ and $SL_{*}^{\varepsilon}(2, A)$

Basic Idea

To look at higher rank groups as non commutative analogues of related lower rank groups

- ▶ Example: To look at the symplectic similitude group $GSp(2n, F)$ in $2n$ variables over a field F as a group $GL(2)$ with coefficients in the matrix ring over F , satisfying suitable commutation relations which involve the transpose map*
- ▶ To extend to higher rank groups, successful methods used for lower rank groups. In particular, having presentations for higher rank classical groups as non commutative versions of known presentations for the lower rank case. These presentations can be used in constructing linear representations for higher rank groups (generalized Weil representations, for example)

Non commutative generalizations of classical groups: $S^{\varepsilon}_*(2, A)$

Let $(A, *)$ be a unitary ring with involution. Let $\varepsilon = \pm 1$

A left ε - Hermitian form H on a left A -module M is a function which is biadditive, left linear in the first variable and such that $H(y, x) = \varepsilon H(x, y)^*$, for all $x, y \in M$.

Let us consider the special case where $M = A^2$

We have a matrix description of left ε - Hermitian forms.

Let us put $T^* = (t_{ji}^*)_{1 \leq i, j \leq 2}$ for any

$T = (t_{ij})_{1 \leq i, j \leq m} \in M(2, A)$. Now, when a basis $B = \{e_1, e_2\}$ of the free A - module M has been chosen, we define the matrix $[H]$ of H with respect to B by $[H] = (H(e_i, e_j))_{1 \leq i, j \leq 2}$.

Then $H(u, v) = u[H]v^*$ ($u, v \in M$) and conversely, given a matrix T such that $T^* = \varepsilon T$ we recover an ε - Hermitian form H_T by $H_T(u, v) = uTv^*$ ($u, v \in M$).

Non commutative generalizations of classical groups: $S^{\varepsilon}_*(2, A)$

Let $(A, *)$ be a unitary ring with involution. Let $\varepsilon = \pm 1$

A left ε - Hermitian form H on a left A -module M is a function which is biadditive, left linear in the first variable and such that $H(y, x) = \varepsilon H(x, y)^*$, for all $x, y \in M$.

Let us consider the special case where $M = A^2$

We have a matrix description of left ε - Hermitian forms.

Let us put $T^* = (t_{ji}^*)_{1 \leq i, j \leq 2}$ for any

$T = (t_{ij})_{1 \leq i, j \leq m} \in M(2, A)$. Now, when a basis $B = \{e_1, e_2\}$ of the free A - module M has been chosen, we define the matrix $[H]$ of H with respect to B by $[H] = (H(e_i, e_j))_{1 \leq i, j \leq 2}$.

Then $H(u, v) = u[H]v^*$ ($u, v \in M$) and conversely, given a matrix T such that $T^* = \varepsilon T$ we recover an ε - Hermitian form H_T by $H_T(u, v) = uTv^*$ ($u, v \in M$).

Non commutative generalizations of classical groups: $S^{\varepsilon}_*(2, A)$

Let $(A, *)$ be a unitary ring with involution. Let $\varepsilon = \pm 1$

A left ε - Hermitian form H on a left A -module M is a function which is biadditive, left linear in the first variable and such that $H(y, x) = \varepsilon H(x, y)^*$, for all $x, y \in M$.

Let us consider the special case where $M = A^2$

We have a matrix description of left ε - Hermitian forms.

Let us put $T^* = (t_{ji}^*)_{1 \leq i, j \leq 2}$ for any

$T = (t_{ij})_{1 \leq i, j \leq m} \in M(2, A)$. Now, when a basis $B = \{e_1, e_2\}$ of the free A - module M has been chosen, we define the matrix $[H]$ of H with respect to B by $[H] = (H(e_i, e_j))_{1 \leq i, j \leq 2}$.

Then $H(u, v) = u[H]v^*$ ($u, v \in M$) and conversely, given a matrix T such that $T^* = \varepsilon T$ we recover an ε - Hermitian form H_T by $H_T(u, v) = uTv^*$ ($u, v \in M$).

Non commutative generalizations of classical groups: $SI_*^\varepsilon(2, A)$

Let $(A, *)$ be a unitary ring with involution. Let $\varepsilon = \pm 1$

A left ε - Hermitian form H on a left A -module M is a function which is biadditive, left linear in the first variable and such that $H(y, x) = \varepsilon H(x, y)^*$, for all $x, y \in M$.

Let us consider the special case where $M = A^2$

We have a matrix description of left ε - Hermitian forms.

Let us put $T^* = (t_{ji}^*)_{1 \leq i, j \leq 2}$ for any

$T = (t_{ij})_{1 \leq i, j \leq m} \in M(2, A)$. Now, when a basis $B = \{e_1, e_2\}$ of the free A - module M has been chosen, we define the matrix $[H]$ of H with respect to B by $[H] = (H(e_i, e_j))_{1 \leq i, j \leq 2}$.

Then $H(u, v) = u[H]v^*$ ($u, v \in M$) and conversely, given a matrix T such that $T^* = \varepsilon T$ we recover an ε - Hermitian form H_T by $H_T(u, v) = uTv^*$ ($u, v \in M$).

The groups $GL_*^\varepsilon(2, A)$ and $SL_*^\varepsilon(2, A)$ can be described as follow:

The group $GL_*^\varepsilon(2, A)$ is the set of all automorphisms g of the A -module $M = A \times A$ such that $H_\varepsilon \circ (g \times g) = \mu_g H_\varepsilon$ ($\mu_g \in A$ central and symmetric with respect to $*$) or in matrix form as

$$GL_*^\varepsilon(2, A) = \{M \in M(2, A) : MJ_\varepsilon M^* = \mu_g J_\varepsilon\}$$

where $J_\varepsilon = \begin{pmatrix} 0 & 1 \\ \varepsilon & 0 \end{pmatrix}$ and $\varepsilon = \pm 1$

while $SL_*^\varepsilon(2, A)$ is the subgroup of $GL_*^\varepsilon(2, A)$ of those g such that $\mu_g = 1$

The groups $GL_*^\varepsilon(2, A)$ and $SL_*^\varepsilon(2, A)$ can be described as follow:

The group $GL_*^\varepsilon(2, A)$ is the set of all automorphisms g of the A - module $M = A \times A$ such that $H_\varepsilon \circ (g \times g) = \mu_g H_\varepsilon$ ($\mu_g \in A$ central and symmetric with respect to $*$) or in matrix form as

$$GL_*^\varepsilon(2, A) = \{M \in M(2, A) : MJ_\varepsilon M^* = \mu_g J_\varepsilon\}$$

where $J_\varepsilon = \begin{pmatrix} 0 & 1 \\ \varepsilon & 0 \end{pmatrix}$ and $\varepsilon = \pm 1$

while $SL_*^\varepsilon(2, A)$ is the subgroup of $GL_*^\varepsilon(2, A)$ of those g such that $\mu_g = 1$

In other words

$Sl_*^\varepsilon(2, A)$ is the group of $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ such that

$$ab^* = -\varepsilon ba^*$$

$$cd^* = -\varepsilon dc^*,$$

$$a^*c = -\varepsilon c^*a,$$

$$b^*d = -\varepsilon d^*b,$$

$$ad^* + \varepsilon bc^* = a^*d + \varepsilon c^*b = 1$$

Weil representations for $G = S^{\varepsilon}_*(2, A)$, G with a Bruhat presentation

We assume that the group $G = S^{\varepsilon}_*(2, A)$ has a Bruhat presentation, i.e.,

if $A_s = A^{sym}$ (ε -symmetric elements) is the set of a such that $a^* + \varepsilon a = 0$, G is generated by the following matrices satisfying the minimal relations

$$h_t = \begin{pmatrix} t & 0 \\ 0 & t^{*-1} \end{pmatrix} (t \in A^\times), w = w_\varepsilon = \begin{pmatrix} 0 & 1 \\ \varepsilon & 0 \end{pmatrix} \text{ and}$$

$$u_s = \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix} (s \in A^{sym})$$

$$h_t h_{t'} = h_{tt'}, u_b u_{b'} = u_{b+b'};$$

$$w^2 = h_\varepsilon;$$

$$h_t u_b = u_{tb} h_t;$$

$$w h_t = h_{t^{*-1}} w;$$

$$w u_{t-1} w u_{-\varepsilon t} w u_{t-1} = h_{-\varepsilon t}, \text{ with } t \text{ an invertible } \varepsilon\text{-symmetric element in } A.$$

Weil representations for $G = S/_{*}^{\varepsilon}(2, A)$, G with a Bruhat presentation

We assume that the group $G = S/_{*}^{\varepsilon}(2, A)$ has a Bruhat presentation, i.e.,

if $A_s = A^{sym}$ (ε -symmetric elements) is the set of a such that $a^* + \varepsilon a = 0$, G is generated by the following matrices satisfying the minimal relations

$$h_t = \begin{pmatrix} t & 0 \\ 0 & t^{*-1} \end{pmatrix} (t \in A^{\times}), w = w_{\varepsilon} = \begin{pmatrix} 0 & 1 \\ \varepsilon & 0 \end{pmatrix} \text{ and}$$

$$u_s = \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix} (s \in A^{sym})$$

$$h_t h_{t'} = h_{tt'}, u_b u_{b'} = u_{b+b'};$$

$$w^2 = h_{\varepsilon};$$

$$h_t u_b = u_{tb} h_t;$$

$$w h_t = h_{t^{*-1}} w;$$

$$w u_{t^{-1}} w u_{-\varepsilon t} w u_{t^{-1}} = h_{-\varepsilon t}, \text{ with } t \text{ an invertible } \varepsilon\text{-symmetric element in } A.$$

Weakly euclidean rigs

- ▶ $(A, *)$ is a weakly euclidean ring if given $a, b \in A$ such that $a^*b = -\varepsilon b^*a$, $Aa + Ab = A$, then there are finite sequences of elements $s_0, s_1, \dots, s_{n-1} \in A_s$ and $r_1, r_2, \dots, r_n \in A$ with $r_n \in A^\times$ such that

$$a = s_0 b + r_1$$

$$b = s_1 r_1 + r_2$$

...

$$r_{n-2} = s_{n-1} r_{n-1} + r_n$$

- ▶ We observe that if $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_*^\varepsilon(2, A)$, then a and b (or c) satisfy above
- ▶ and we have:

If $(A, *)$ is a weakly euclidean ring, then the elements u_s, h_t and w , generate the group $SL_*^\varepsilon(2, A)$.

Weakly euclidean rigs

- ▶ $(A, *)$ is a weakly euclidean ring if given $a, b \in A$ such that $a^*b = -\varepsilon b^*a$, $Aa + Ab = A$, then there are finite sequences of elements $s_0, s_1, \dots, s_{n-1} \in A_s$ and $r_1, r_2, \dots, r_n \in A$ with $r_n \in A^\times$ such that

$$a = s_0 b + r_1$$

$$b = s_1 r_1 + r_2$$

...

$$r_{n-2} = s_{n-1} r_{n-1} + r_n$$

- ▶ We observe that if $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_*^\varepsilon(2, A)$, then a and b (or c) satisfy above
- ▶ and we have:

If $(A, *)$ is a weakly euclidean ring, then the elements u_s, h_t and w , generate the group $SL_*^\varepsilon(2, A)$.

Weakly euclidean rigs

- ▶ $(A, *)$ is a weakly euclidean ring if given $a, b \in A$ such that $a^*b = -\varepsilon b^*a$, $Aa + Ab = A$, then there are finite sequences of elements $s_0, s_1, \dots, s_{n-1} \in A_s$ and $r_1, r_2, \dots, r_n \in A$ with $r_n \in A^\times$ such that

$$a = s_0 b + r_1$$

$$b = s_1 r_1 + r_2$$

...

$$r_{n-2} = s_{n-1} r_{n-1} + r_n$$

- ▶ We observe that if $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_*^\varepsilon(2, A)$, then a and b (or c) satisfy above
- ▶ and we have:

If $(A, *)$ is a weakly euclidean ring, then the elements u_s, h_t and w , generate the group $SL_*^\varepsilon(2, A)$.

Weakly euclidean rigs

- ▶ $(A, *)$ is a weakly euclidean ring if given $a, b \in A$ such that $a^*b = -\varepsilon b^*a$, $Aa + Ab = A$, then there are finite sequences of elements $s_0, s_1, \dots, s_{n-1} \in A_s$ and $r_1, r_2, \dots, r_n \in A$ with $r_n \in A^\times$ such that

$$a = s_0 b + r_1$$

$$b = s_1 r_1 + r_2$$

...

$$r_{n-2} = s_{n-1} r_{n-1} + r_n$$

- ▶ We observe that if $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_*^\varepsilon(2, A)$, then a and b (or c) satisfy above
- ▶ and we have:

If $(A, *)$ is a weakly euclidean ring, then the elements u_s, h_t and w , generate the group $SL_*^\varepsilon(2, A)$.

In what follows we suppose that:

A is a finite involutive (unitary) ring

We fix the following functions with the below properties, which will be useful in the construction of the Weil representation of G . Let M be a finite (right) A -module

In what follows we suppose that:

A is a finite involutive (unitary) ring

We fix the following functions with the below properties, which will be useful in the construction of the Weil representation of G . Let M be a finite (right) A -module

In what follows we suppose that:

A is a finite involutive (unitary) ring

We fix the following functions with the below properties, which will be useful in the construction of the Weil representation of G . Let M be a finite (right) A —module

1. A bi-additive function $\chi : M \times M \rightarrow \mathbb{C}^\times$ and a character $\alpha \in \widehat{A}^\times$ such that:

1.1 $\chi(xt, y) = \alpha(tt^*)\chi(x, yt^*)$ for $x, y \in M$ and $t \in A^\times$ (χ is α -balanced).

1.2 $\chi(y, x) = [\chi(x, y)]^{-\varepsilon}$. We observe that $[\chi(x, y)]^{-\varepsilon} = \chi(-\varepsilon x, y)$ (χ is ε -symmetric).

1.3 $\chi(x, y) = 1$ for any $x \in M$, implies $y = 0$ (χ is non-degenerate).

2. A function $\gamma : A^{\text{sym}} \times M \rightarrow \mathbb{C}^\times$ such that:

2.1 $\gamma(b + b', x) = \gamma(b, x)\gamma(b', x)$, for all $b, b' \in A^{\text{sym}}$ and $x \in M$.

2.2 $\gamma(b, xt) = \gamma(tbt^*, x)$ or equivalently $\gamma(b, x) = \gamma(tbt^*, xt^{-1})$, for all $b \in A^{\text{sym}}$, $t \in A^\times$ and $x \in M$.

2.3 $\gamma(t, x + z) = \gamma(t, x)\gamma(t, z)\chi(x, zt)$ for all $x, z \in M, t \in A^{\text{sym}}$, where t is ε -symmetric invertible in A and $c \in \mathbb{C}^\times$ satisfies $c^2 |M| = \alpha(\varepsilon)$.

3. We assume moreover that these data are related by the equation:

$$c\gamma(-\varepsilon t, x) \sum_{y \in M} \chi(-\varepsilon x, y)\gamma(t^{-1}, y) = \alpha(-t).$$

1. A bi-additive function $\chi : M \times M \rightarrow \mathbb{C}^\times$ and a character $\alpha \in \widehat{A}^\times$ such that:

1.1 $\chi(xt, y) = \alpha(tt^*)\chi(x, yt^*)$ for $x, y \in M$ and $t \in A^\times$ (χ is α -balanced).

1.2 $\chi(y, x) = [\chi(x, y)]^{-\varepsilon}$. We observe that $[\chi(x, y)]^{-\varepsilon} = \chi(-\varepsilon x, y)$ (χ is ε -symmetric).

1.3 $\chi(x, y) = 1$ for any $x \in M$, implies $y = 0$ (χ is non-degenerate).

2. A function $\gamma : A^{\text{sym}} \times M \rightarrow \mathbb{C}^\times$ such that:

2.1 $\gamma(b + b', x) = \gamma(b, x)\gamma(b', x)$, for all $b, b' \in A^{\text{sym}}$ and $x \in M$.

2.2 $\gamma(b, xt) = \gamma(tbt^*, x)$ or equivalently $\gamma(b, x) = \gamma(tbt^*, xt^{-1})$, for all $b \in A^{\text{sym}}$, $t \in A^\times$ and $x \in M$.

2.3 $\gamma(t, x + z) = \gamma(t, x)\gamma(t, z)\chi(x, zt)$ for all $x, z \in M, t \in A^{\text{sym}}$, where t is ε -symmetric invertible in A and $c \in \mathbb{C}^\times$ satisfies $c^2 |M| = \alpha(\varepsilon)$.

3. We assume moreover that these data are related by the equation:

$$c\gamma(-\varepsilon t, x) \sum_{y \in M} \chi(-\varepsilon x, y)\gamma(t^{-1}, y) = \alpha(-t).$$

1. A bi-additive function $\chi : M \times M \rightarrow \mathbb{C}^\times$ and a character $\alpha \in \widehat{A^\times}$ such that:

1.1 $\chi(xt, y) = \alpha(tt^*)\chi(x, yt^*)$ for $x, y \in M$ and $t \in A^\times$ (χ is α -balanced).

1.2 $\chi(y, x) = [\chi(x, y)]^{-\varepsilon}$. We observe that $[\chi(x, y)]^{-\varepsilon} = \chi(-\varepsilon x, y)$ (χ is ε -symmetric).

1.3 $\chi(x, y) = 1$ for any $x \in M$, implies $y = 0$ (χ is non-degenerate).

2. A function $\gamma : A^{\text{sym}} \times M \rightarrow \mathbb{C}^\times$ such that:

2.1 $\gamma(b + b', x) = \gamma(b, x)\gamma(b', x)$, for all $b, b' \in A^{\text{sym}}$ and $x \in M$.

2.2 $\gamma(b, xt) = \gamma(tbt^*, x)$ or equivalently $\gamma(b, x) = \gamma(tbt^*, xt^{-1})$, for all $b \in A^{\text{sym}}$, $t \in A^\times$ and $x \in M$.

2.3 $\gamma(t, x + z) = \gamma(t, x)\gamma(t, z)\chi(x, zt)$ for all $x, z \in M, t \in A^{\text{sym}}$, where t is ε -symmetric invertible in A and $c \in \mathbb{C}^\times$ satisfies $c^2 |M| = \alpha(\varepsilon)$.

3. We assume moreover that these data are related by the equation:

$$c\gamma(-\varepsilon t, x) \sum_{y \in M} \chi(-\varepsilon x, y)\gamma(t^{-1}, y) = \alpha(-t).$$

1. A bi-additive function $\chi : M \times M \rightarrow \mathbb{C}^\times$ and a character $\alpha \in \widehat{A^\times}$ such that:

1.1 $\chi(xt, y) = \alpha(tt^*)\chi(x, yt^*)$ for $x, y \in M$ and $t \in A^\times$ (χ is α -balanced).

1.2 $\chi(y, x) = [\chi(x, y)]^{-\varepsilon}$. We observe that $[\chi(x, y)]^{-\varepsilon} = \chi(-\varepsilon x, y)$ (χ is ε -symmetric).

1.3 $\chi(x, y) = 1$ for any $x \in M$, implies $y = 0$ (χ is non-degenerate).

2. A function $\gamma : A^{\text{sym}} \times M \rightarrow \mathbb{C}^\times$ such that:

2.1 $\gamma(b + b', x) = \gamma(b, x)\gamma(b', x)$, for all $b, b' \in A^{\text{sym}}$ and $x \in M$.

2.2 $\gamma(b, xt) = \gamma(tbt^*, x)$ or equivalently $\gamma(b, x) = \gamma(tbt^*, xt^{-1})$, for all $b \in A^{\text{sym}}$, $t \in A^\times$ and $x \in M$.

2.3 $\gamma(t, x + z) = \gamma(t, x)\gamma(t, z)\chi(x, zt)$ for all $x, z \in M, t \in A^{\text{sym}}$, where t is ε -symmetric invertible in A and $c \in \mathbb{C}^\times$ satisfies $c^2 |M| = \alpha(\varepsilon)$.

3. We assume moreover that these data are related by the equation:

$$c\gamma(-\varepsilon t, x) \sum_{y \in M} \chi(-\varepsilon x, y)\gamma(t^{-1}, y) = \alpha(-t).$$

1. A bi-additive function $\chi : M \times M \rightarrow \mathbb{C}^\times$ and a character $\alpha \in \widehat{A}^\times$ such that:

1.1 $\chi(xt, y) = \alpha(tt^*)\chi(x, yt^*)$ for $x, y \in M$ and $t \in A^\times$ (χ is α -balanced).

1.2 $\chi(y, x) = [\chi(x, y)]^{-\varepsilon}$. We observe that $[\chi(x, y)]^{-\varepsilon} = \chi(-\varepsilon x, y)$ (χ is ε -symmetric).

1.3 $\chi(x, y) = 1$ for any $x \in M$, implies $y = 0$ (χ is non-degenerate).

2. A function $\gamma : A^{\text{sym}} \times M \rightarrow \mathbb{C}^\times$ such that:

2.1 $\gamma(b + b', x) = \gamma(b, x)\gamma(b', x)$, for all $b, b' \in A^{\text{sym}}$ and $x \in M$.

2.2 $\gamma(b, xt) = \gamma(tbt^*, x)$ or equivalently $\gamma(b, x) = \gamma(tbt^*, xt^{-1})$, for all $b \in A^{\text{sym}}$, $t \in A^\times$ and $x \in M$.

2.3 $\gamma(t, x + z) = \gamma(t, x)\gamma(t, z)\chi(x, zt)$ for all $x, z \in M$, $t \in A^{\text{sym}}$, where t is ε -symmetric invertible in A and $c \in \mathbb{C}^\times$ satisfies $c^2 |M| = \alpha(\varepsilon)$.

3. We assume moreover that these data are related by the equation:

$$c\gamma(-\varepsilon t, x) \sum_{y \in M} \chi(-\varepsilon x, y)\gamma(t^{-1}, y) = \alpha(-t).$$

1. A bi-additive function $\chi : M \times M \rightarrow \mathbb{C}^\times$ and a character $\alpha \in \widehat{A}^\times$ such that:

1.1 $\chi(xt, y) = \alpha(tt^*)\chi(x, yt^*)$ for $x, y \in M$ and $t \in A^\times$ (χ is α -balanced).

1.2 $\chi(y, x) = [\chi(x, y)]^{-\varepsilon}$. We observe that $[\chi(x, y)]^{-\varepsilon} = \chi(-\varepsilon x, y)$ (χ is ε -symmetric).

1.3 $\chi(x, y) = 1$ for any $x \in M$, implies $y = 0$ (χ is non-degenerate).

2. A function $\gamma : A^{\text{sym}} \times M \rightarrow \mathbb{C}^\times$ such that:

2.1 $\gamma(b + b', x) = \gamma(b, x)\gamma(b', x)$, for all b, b'^{sym} and $x \in M$.

2.2 $\gamma(b, xt) = \gamma(tbt^*, x)$ or equivalently $\gamma(b, x) = \gamma(tbt^*, xt^{-1})$, for all $b \in A^{\text{sym}}$, $t \in A^\times$ and $x \in M$.

2.3 $\gamma(t, x + z) = \gamma(t, x)\gamma(t, z)\chi(x, zt)$ for all $x, z \in M, t \in A^{\text{sym}}$. where t is ε -symmetric invertible in A and $c \in \mathbb{C}^\times$ satisfies $c^2 |M| = \alpha(\varepsilon)$.

3. We assume moreover that these data are related by the equation:

$$c\gamma(-\varepsilon t, x) \sum_{y \in M} \chi(-\varepsilon x, y)\gamma(t^{-1}, y) = \alpha(-t).$$

1. A bi-additive function $\chi : M \times M \rightarrow \mathbb{C}^\times$ and a character $\alpha \in \widehat{A^\times}$ such that:

1.1 $\chi(xt, y) = \alpha(tt^*)\chi(x, yt^*)$ for $x, y \in M$ and $t \in A^\times$ (χ is α -balanced).

1.2 $\chi(y, x) = [\chi(x, y)]^{-\varepsilon}$. We observe that $[\chi(x, y)]^{-\varepsilon} = \chi(-\varepsilon x, y)$ (χ is ε -symmetric).

1.3 $\chi(x, y) = 1$ for any $x \in M$, implies $y = 0$ (χ is non-degenerate).

2. A function $\gamma : A^{\text{sym}} \times M \rightarrow \mathbb{C}^\times$ such that:

2.1 $\gamma(b + b', x) = \gamma(b, x)\gamma(b', x)$, for all b, b'^{sym} and $x \in M$.

2.2 $\gamma(b, xt) = \gamma(tbt^*, x)$ or equivalently $\gamma(b, x) = \gamma(tbt^*, xt^{-1})$, for all $b \in A^{\text{sym}}$, $t \in A^\times$ and $x \in M$.

2.3 $\gamma(t, x + z) = \gamma(t, x)\gamma(t, z)\chi(x, zt)$ for all $x, z \in M, t \in A^{\text{sym}}$. where t is ε -symmetric invertible in A and $c \in \mathbb{C}^\times$ satisfies $c^2 |M| = \alpha(\varepsilon)$.

3. We assume moreover that these data are related by the equation:

$$c\gamma(-\varepsilon t, x) \sum_{y \in M} \chi(-\varepsilon x, y)\gamma(t^{-1}, y) = \alpha(-t).$$

1. A bi-additive function $\chi : M \times M \rightarrow \mathbb{C}^\times$ and a character $\alpha \in \widehat{A}^\times$ such that:

1.1 $\chi(xt, y) = \alpha(tt^*)\chi(x, yt^*)$ for $x, y \in M$ and $t \in A^\times$ (χ is α -balanced).

1.2 $\chi(y, x) = [\chi(x, y)]^{-\varepsilon}$. We observe that $[\chi(x, y)]^{-\varepsilon} = \chi(-\varepsilon x, y)$ (χ is ε -symmetric).

1.3 $\chi(x, y) = 1$ for any $x \in M$, implies $y = 0$ (χ is non-degenerate).

2. A function $\gamma : A^{\text{sym}} \times M \rightarrow \mathbb{C}^\times$ such that:

2.1 $\gamma(b + b', x) = \gamma(b, x)\gamma(b', x)$, for all b, b'^{sym} and $x \in M$.

2.2 $\gamma(b, xt) = \gamma(tbt^*, x)$ or equivalently $\gamma(b, x) = \gamma(tbt^*, xt^{-1})$, for all $b \in A^{\text{sym}}$, $t \in A^\times$ and $x \in M$.

2.3 $\gamma(t, x + z) = \gamma(t, x)\gamma(t, z)\chi(x, zt)$ for all $x, z \in M, t \in A^{\text{sym}}$. where t is ε -symmetric invertible in A and $c \in \mathbb{C}^\times$ satisfies $c^2 |M| = \alpha(\varepsilon)$.

3. We assume moreover that these data are related by the equation:

$$c\gamma(-\varepsilon t, x) \sum_{y \in M} \chi(-\varepsilon x, y)\gamma(t^{-1}, y) = \alpha(-t).$$

1. A bi-additive function $\chi : M \times M \rightarrow \mathbb{C}^\times$ and a character $\alpha \in \widehat{A^\times}$ such that:

1.1 $\chi(xt, y) = \alpha(tt^*)\chi(x, yt^*)$ for $x, y \in M$ and $t \in A^\times$ (χ is α -balanced).

1.2 $\chi(y, x) = [\chi(x, y)]^{-\varepsilon}$. We observe that $[\chi(x, y)]^{-\varepsilon} = \chi(-\varepsilon x, y)$ (χ is ε -symmetric).

1.3 $\chi(x, y) = 1$ for any $x \in M$, implies $y = 0$ (χ is non-degenerate).

2. A function $\gamma : A^{\text{sym}} \times M \rightarrow \mathbb{C}^\times$ such that:

2.1 $\gamma(b + b', x) = \gamma(b, x)\gamma(b', x)$, for all b, b'^{sym} and $x \in M$.

2.2 $\gamma(b, xt) = \gamma(tbt^*, x)$ or equivalently $\gamma(b, x) = \gamma(tbt^*, xt^{-1})$, for all $b \in A^{\text{sym}}$, $t \in A^\times$ and $x \in M$.

2.3 $\gamma(t, x + z) = \gamma(t, x)\gamma(t, z)\chi(x, zt)$ for all $x, z \in M, t \in A^{\text{sym}}$. where t is ε -symmetric invertible in A and $c \in \mathbb{C}^\times$ satisfies $c^2 |M| = \alpha(\varepsilon)$.

3. We assume moreover that these data are related by the equation:

$$c\gamma(-\varepsilon t, x) \sum_{y \in M} \chi(-\varepsilon x, y)\gamma(t^{-1}, y) = \alpha(-t).$$

As a consequence we have

The following two relations are equivalent:

1. $c\gamma(-\varepsilon t, x) \sum_{y \in M} \chi(-\varepsilon x, y) \gamma(t^{-1}, y) = \alpha(-t).$
2. $\sum_{y \in M} \gamma(t, y) = \frac{\alpha(\varepsilon t)}{c}.$

As a consequence we have

The following two relations are equivalent:

1. $c\gamma(-\varepsilon t, x) \sum_{y \in M} \chi(-\varepsilon x, y) \gamma(t^{-1}, y) = \alpha(-t).$
2. $\sum_{y \in M} \gamma(t, y) = \frac{\alpha(\varepsilon t)}{c}.$

As a consequence we have

The following two relations are equivalent:

1. $c\gamma(-\varepsilon t, x) \sum_{y \in M} \chi(-\varepsilon x, y) \gamma(t^{-1}, y) = \alpha(-t).$
2. $\sum_{y \in M} \gamma(t, y) = \frac{\alpha(\varepsilon t)}{c}.$

As a consequence we have

The following two relations are equivalent:

1. $c\gamma(-\varepsilon t, x) \sum_{y \in M} \chi(-\varepsilon x, y) \gamma(t^{-1}, y) = \alpha(-t).$
2. $\sum_{y \in M} \gamma(t, y) = \frac{\alpha(\varepsilon t)}{c}.$

Weil Representation defined via a presentation

We have the following result:

► Theorem

There is a representation (\mathbb{C}^M, ρ) of G defined as follows on the basis of Dirac delta functions $\{e_x\}_{x \in M}$ (given by $e_x(y) = 1$ if $y = x$; $e_x(y) = 0$ otherwise):

$$\rho_{u_b}(e_x) = \gamma(b, x)e_x$$

$$\rho_{h_t}(e_x) = \alpha(t)e_{xt^{-1}}$$

$$\rho_w(e_x) = c \sum_{y \in M} \chi(-\varepsilon x, y)e_y$$

for $x \in M, b \in A^\times \cap A^\times, t \in A^\times$.

This representation is called the generalized Weil representation of $SL_^\varepsilon(2, A)$ associated to the data $(M, \alpha, \gamma, \chi)$.*

Weil Representation defined via a presentation

We have the following result:

► Theorem

There is a representation (\mathbb{C}^M, ρ) of G defined as follows on the basis of Dirac delta functions $\{e_x\}_{x \in M}$ (given by $e_x(y) = 1$ if $y = x$; $e_x(y) = 0$ otherwise):

$$\rho_{U_b}(e_x) = \gamma(b, x)e_x$$

$$\rho_{h_t}(e_x) = \alpha(t)e_{xt^{-1}}$$

$$\rho_w(e_x) = c \sum_{y \in M} \chi(-\varepsilon x, y)e_y$$

for $x \in M, b \in A^\times \cap A^\times, t \in A^\times$.

This representation is called the generalized Weil representation of $SL_^\varepsilon(2, A)$ associated to the data $(M, \alpha, \gamma, \chi)$.*

Weil Representation defined via a presentation

We have the following result:

► Theorem

There is a representation (\mathbb{C}^M, ρ) of G defined as follows on the basis of Dirac delta functions $\{e_x\}_{x \in M}$ (given by $e_x(y) = 1$ if $y = x$; $e_x(y) = 0$ otherwise):

$$\rho_{u_b}(e_x) = \gamma(b, x)e_x$$

$$\rho_{h_t}(e_x) = \alpha(t)e_{xt^{-1}}$$

$$\rho_w(e_x) = c \sum_{y \in M} \chi(-\varepsilon x, y)e_y$$

for $x \in M, b \in A^\times \cap A^\times, t \in A^\times$.

This representation is called the generalized Weil representation of $SL_^\varepsilon(2, A)$ associated to the data $(M, \alpha, \gamma, \chi)$.*

Proof: It is enough to verify that $\rho_{u_b}, \rho_{h_t}, \rho_w$ defined as above, satisfy the relations corresponding to the universal relations in the Bruhat presentation of G .

To this end, we observe that

- ▶ a) $(\rho_{h_t} \circ \rho_{h_{t'}})(e_x) = \rho_{h_t}(\alpha(t')e_{xt'^{-1}}) = \alpha(t')\alpha(t)e_{xt'^{-1}t^{-1}} = \alpha(tt')\alpha(t)e_{x(tt')^{-1}} = \rho_{h_{tt'}}(e_x)$
- ▶ b) $(\rho_{u_b} \circ \rho_{u_{b'}})(e_x) = \gamma(b', x)\gamma(b, x)e_x = \gamma(b + b', x)e_x = (\rho_{u_{b+b'}})(e_x)$
- ▶ c) $(\rho_w \circ \rho_w)(e_x) = c^2 \sum_{z \in M} \sum_{y \in M} \chi(-\varepsilon x, y)\chi(-\varepsilon y, z)e_z = \alpha(\varepsilon)e_{x\varepsilon} = \rho_{h_\varepsilon}e_x$
- ▶ d) $(\rho_{h_t} \circ \rho_{u_b})(e_x) = \gamma(b, x)\alpha(t)e_{xt^{-1}} = \alpha(t)\gamma(tbt^*, xt^{-1})e_{xt^{-1}} = (\rho_{tbt^*} \circ \rho_{h_t})(e_x)$

Proof: It is enough to verify that $\rho_{u_b}, \rho_{h_t}, \rho_w$ defined as above, satisfy the relations corresponding to the universal relations in the Bruhat presentation of G .

To this end, we observe that

- ▶ a) $(\rho_{h_t} \circ \rho_{h_{t'}})(e_x) = \rho_{h_t}(\alpha(t')e_{xt'-1}) = \alpha(t')\alpha(t)e_{xt'-1t-1} = \alpha(t't)\alpha(t)e_{x(tt')^{-1}} = \alpha(tt')e_{x(tt')^{-1}} = \rho_{h_{tt'}}(e_x)$
- ▶ b) $(\rho_{u_b} \circ \rho_{u_{b'}})(e_x) = \gamma(b', x)\gamma(b, x)e_x = \gamma(b + b', x)e_x = (\rho_{u_{b+b'}})(e_x)$
- ▶ c) $(\rho_w \circ \rho_w)(e_x) = c^2 \sum_{z \in M} \sum_{y \in M} \chi(-\varepsilon x, y)\chi(-\varepsilon y, z)e_z = \alpha(\varepsilon)e_{x\varepsilon} = \rho_{h_\varepsilon}e_x$
- ▶ d) $(\rho_{h_t} \circ \rho_{u_b})(e_x) = \gamma(b, x)\alpha(t)e_{xt-1} = \alpha(t)\gamma(tbt^*, xt^{-1})e_{xt-1} = (\rho_{tbt^*} \circ \rho_{h_t})(e_x)$

Proof: It is enough to verify that $\rho_{u_b}, \rho_{h_t}, \rho_w$ defined as above, satisfy the relations corresponding to the universal relations in the Bruhat presentation of G .

To this end, we observe that

- ▶ a) $(\rho_{h_t} \circ \rho_{h_{t'}})(e_x) = \rho_{h_t}(\alpha(t')e_{xt'-1}) = \alpha(t')\alpha(t)e_{xt'-1t-1} = \alpha(tt')\alpha(t)e_{x(tt')^{-1}} = \rho_{h_{tt'}}(e_x)$
- ▶ b) $(\rho_{u_b} \circ \rho_{u_{b'}})(e_x) = \gamma(b', x)\gamma(b, x)e_x = \gamma(b + b', x)e_x = (\rho_{u_{b+b'}})(e_x)$
- ▶ c) $(\rho_w \circ \rho_w)(e_x) = c^2 \sum_{z \in M} \sum_{y \in M} \chi(-\varepsilon x, y)\chi(-\varepsilon y, z)e_z = \alpha(\varepsilon)e_{x\varepsilon} = \rho_{h_\varepsilon}e_x$
- ▶ d) $(\rho_{h_t} \circ \rho_{u_b})(e_x) = \gamma(b, x)\alpha(t)e_{xt-1} = \alpha(t)\gamma(tbt^*, xt^{-1})e_{xt-1} = (\rho_{tbt^*} \circ \rho_{h_t})(e_x)$

Proof: It is enough to verify that $\rho_{u_b}, \rho_{h_t}, \rho_w$ defined as above, satisfy the relations corresponding to the universal relations in the Bruhat presentation of G .

To this end, we observe that

- ▶ a) $(\rho_{h_t} \circ \rho_{h_{t'}})(e_x) = \rho_{h_t}(\alpha(t')e_{xt'-1}) = \alpha(t')\alpha(t)e_{xt'-1t-1} = \alpha(tt')\alpha(t)e_{x(tt')^{-1}} = \rho_{h_{tt'}}(e_x)$
- ▶ b) $(\rho_{u_b} \circ \rho_{u_{b'}})(e_x) = \gamma(b', x)\gamma(b, x)e_x = \gamma(b + b', x)e_x = (\rho_{u_{b+b'}})(e_x)$
- ▶ c) $(\rho_w \circ \rho_w)(e_x) = c^2 \sum_{z \in M} \sum_{y \in M} \chi(-\varepsilon x, y)\chi(-\varepsilon y, z)e_z = \alpha(\varepsilon)e_{x\varepsilon} = \rho_{h_\varepsilon}e_x$
- ▶ d) $(\rho_{h_t} \circ \rho_{u_b})(e_x) = \gamma(b, x)\alpha(t)e_{xt-1} = \alpha(t)\gamma(tbt^*, xt^{-1})e_{xt-1} = (\rho_{tbt^*} \circ \rho_{h_t})(e_x)$

Proof: It is enough to verify that $\rho_{u_b}, \rho_{h_t}, \rho_w$ defined as above, satisfy the relations corresponding to the universal relations in the Bruhat presentation of G .

To this end, we observe that

- ▶ a) $(\rho_{h_t} \circ \rho_{h_{t'}})(e_x) = \rho_{h_t}(\alpha(t')e_{xt'-1}) = \alpha(t')\alpha(t)e_{xt'-1t-1} = \alpha(tt')\alpha(t)e_{x(tt')^{-1}} = \rho_{h_{tt'}}(e_x)$
- ▶ b) $(\rho_{u_b} \circ \rho_{u_{b'}})(e_x) = \gamma(b', x)\gamma(b, x)e_x = \gamma(b + b', x)e_x = (\rho_{u_{b+b'}})(e_x)$
- ▶ c) $(\rho_w \circ \rho_w)(e_x) = c^2 \sum_{z \in M} \sum_{y \in M} \chi(-\varepsilon x, y)\chi(-\varepsilon y, z)e_z = \alpha(\varepsilon)e_{x\varepsilon} = \rho_{h_\varepsilon}e_x$
- ▶ d) $(\rho_{h_t} \circ \rho_{u_b})(e_x) = \gamma(b, x)\alpha(t)e_{xt-1} = \alpha(t)\gamma(tbt^*, xt^{-1})e_{xt-1} = (\rho_{tbt^*} \circ \rho_{h_t})(e_x)$

$$\begin{aligned}
 \blacktriangleright \text{ e) } \rho_w \circ \rho_{h_t})(e_x) &= c\alpha(t) \sum_y \chi(-\varepsilon x t^{-1}, y) e_y = \\
 &= c \sum_y \chi(-\varepsilon x, y t^{*-1}) \alpha(t^{*-1}) e_y = \\
 &= \rho_{h_{t^{*-1}}}(c \sum_y \chi(-\varepsilon x, y) e_y) = (\rho_{h_{t^{*-1}}} \circ \rho_w)(e_x)
 \end{aligned}$$

\blacktriangleright f) Finally, a computation shows that

$$\begin{aligned}
 &(\rho_w \rho_{u_{t^{-1}}} \rho_w \rho_{u_{-\varepsilon t}})(e_x) = \\
 &\sum_z c^2 \gamma(-\varepsilon t, x) \left(\sum_y \chi(-\varepsilon x, y) \gamma(t^{-1}, y) \chi(-\varepsilon y, z) \right) e_z \\
 &= (\rho_{h_{-\varepsilon t}} \rho_{u_{-t^{-1}}} \rho_w \rho_{h_\varepsilon})(e_x) = \\
 &= c\alpha(-t) \sum_z \chi(-\varepsilon x, -\varepsilon z t) \gamma(-t^{-1}, -\varepsilon z t) e_z.
 \end{aligned}$$

So we want to prove that

$$\begin{aligned}
 &c\gamma(-\varepsilon t, x) \sum_y \chi(-\varepsilon x, y) \gamma(t^{-1}, y) \chi(-\varepsilon y, z) = \\
 &\alpha(-t) \chi(-\varepsilon x, -\varepsilon z t) \gamma(-t^{-1}, -\varepsilon z t)
 \end{aligned}$$

But by hypothesis

$$c\gamma(-\varepsilon t, x - \varepsilon z) \sum_{y \in M} \chi(-\varepsilon(x - \varepsilon z), y) \gamma(-\varepsilon t, y t^{-1}) = \alpha(-t),$$

$$\begin{aligned}
 \blacktriangleright \text{ e) } \rho_w \circ \rho_{h_t})(e_x) &= c\alpha(t) \sum_y \chi(-\varepsilon x t^{-1}, y) e_y = \\
 &= c \sum_y \chi(-\varepsilon x, y t^{*-1}) \alpha(t^{*-1}) e_y = \\
 &= \rho_{h_{t^{*-1}}}(c \sum_y \chi(-\varepsilon x, y) e_y) = (\rho_{h_{t^{*-1}}} \circ \rho_w)(e_x)
 \end{aligned}$$

\blacktriangleright f) Finally, a computation shows that

$$\begin{aligned}
 &(\rho_w \rho_{u_{t-1}} \rho_w \rho_{u_{-\varepsilon t}})(e_x) = \\
 &\sum_z c^2 \gamma(-\varepsilon t, x) \left(\sum_y \chi(-\varepsilon x, y) \gamma(t^{-1}, y) \chi(-\varepsilon y, z) \right) e_z \\
 &= (\rho_{h_{-\varepsilon t}} \rho_{u_{t-1}} \rho_w \rho_{h_\varepsilon})(e_x) = \\
 &c\alpha(-t) \sum_z \chi(-\varepsilon x, -\varepsilon z t) \gamma(-t^{-1}, -\varepsilon z t) e_z.
 \end{aligned}$$

So we want to prove that

$$\begin{aligned}
 &c\gamma(-\varepsilon t, x) \sum_y \chi(-\varepsilon x, y) \gamma(t^{-1}, y) \chi(-\varepsilon y, z) = \\
 &\alpha(-t) \chi(-\varepsilon x, -\varepsilon z t) \gamma(-t^{-1}, -\varepsilon z t)
 \end{aligned}$$

But by hypothesis

$$c\gamma(-\varepsilon t, x - \varepsilon z) \sum_{y \in M} \chi(-\varepsilon(x - \varepsilon z), y) \gamma(-\varepsilon t, y t^{-1}) = \alpha(-t),$$

$$\begin{aligned}
 \blacktriangleright \text{ e) } \rho_w \circ \rho_{h_t})(e_x) &= c\alpha(t) \sum_y \chi(-\varepsilon x t^{-1}, y) e_y = \\
 &= c \sum_y \chi(-\varepsilon x, y t^{*-1}) \alpha(t^{*-1}) e_y = \\
 &= \rho_{h_{t^{*-1}}}(c \sum_y \chi(-\varepsilon x, y) e_y) = (\rho_{h_{t^{*-1}}} \circ \rho_w)(e_x)
 \end{aligned}$$

\blacktriangleright f) Finally, a computation shows that

$$\begin{aligned}
 &(\rho_w \rho_{u_{t-1}} \rho_w \rho_{u_{-\varepsilon t}})(e_x) = \\
 &\sum_z c^2 \gamma(-\varepsilon t, x) \left(\sum_y \chi(-\varepsilon x, y) \gamma(t^{-1}, y) \chi(-\varepsilon y, z) \right) e_z \\
 &= (\rho_{h_{-\varepsilon t}} \rho_{u_{t-1}} \rho_w \rho_{h_\varepsilon})(e_x) = \\
 &= c\alpha(-t) \sum_z \chi(-\varepsilon x, -\varepsilon z t) \gamma(-t^{-1}, -\varepsilon z t) e_z.
 \end{aligned}$$

So we want to prove that

$$\begin{aligned}
 &c\gamma(-\varepsilon t, x) \sum_y \chi(-\varepsilon x, y) \gamma(t^{-1}, y) \chi(-\varepsilon y, z) = \\
 &\alpha(-t) \chi(-\varepsilon x, -\varepsilon z t) \gamma(-t^{-1}, -\varepsilon z t)
 \end{aligned}$$

But by hypothesis

$$c\gamma(-\varepsilon t, x - \varepsilon z) \sum_{y \in M} \chi(-\varepsilon(x - \varepsilon z), y) \gamma(-\varepsilon t, y t^{-1}) = \alpha(-t),$$

$$\begin{aligned}
 \blacktriangleright \text{ e) } \rho_w \circ \rho_{h_t})(e_x) &= c\alpha(t) \sum_y \chi(-\varepsilon x t^{-1}, y) e_y = \\
 &= c \sum_y \chi(-\varepsilon x, y t^{*-1}) \alpha(t^{*-1}) e_y = \\
 &= \rho_{h_{t^{*-1}}} (c \sum_y \chi(-\varepsilon x, y) e_y) = (\rho_{h_{t^{*-1}}} \circ \rho_w)(e_x)
 \end{aligned}$$

\blacktriangleright f) Finally, a computation shows that

$$\begin{aligned}
 &(\rho_w \rho_{u_{t-1}} \rho_w \rho_{u_{-\varepsilon t}})(e_x) = \\
 &\sum_z c^2 \gamma(-\varepsilon t, x) \left(\sum_y \chi(-\varepsilon x, y) \gamma(t^{-1}, y) \chi(-\varepsilon y, z) \right) e_z \\
 &= (\rho_{h_{-\varepsilon t}} \rho_{u_{-t-1}} \rho_w \rho_{h_\varepsilon})(e_x) = \\
 &c\alpha(-t) \sum_z \chi(-\varepsilon x, -\varepsilon z t) \gamma(-t^{-1}, -\varepsilon z t) e_z.
 \end{aligned}$$

So we want to prove that

$$\begin{aligned}
 &c\gamma(-\varepsilon t, x) \sum_y \chi(-\varepsilon x, y) \gamma(t^{-1}, y) \chi(-\varepsilon y, z) = \\
 &\alpha(-t) \chi(-\varepsilon x, -\varepsilon z t) \gamma(-t^{-1}, -\varepsilon z t)
 \end{aligned}$$

But by hypothesis

$$c\gamma(-\varepsilon t, x - \varepsilon z) \sum_{y \in M} \chi(-\varepsilon(x - \varepsilon z), y) \gamma(-\varepsilon t, y t^{-1}) = \alpha(-t),$$

so

$$c\gamma(-\varepsilon t, x) \sum_y \chi(-\varepsilon x, y) \gamma(t^{-1}, y) \chi(-\varepsilon y, z) =$$

$$c\gamma(-\varepsilon t, x) \sum_y \chi(-\varepsilon x + z, y) \gamma(-\varepsilon t, yt^{-1}) =$$

$$c\gamma(-\varepsilon t, x) \frac{\alpha(-t)}{c\gamma(-\varepsilon t, x - \varepsilon z)} = \frac{\gamma(-\varepsilon t, x) \alpha(-t)}{\gamma(-\varepsilon t, x) \gamma(-\varepsilon t, -\varepsilon z) \chi(x, zt)} =$$

$$\alpha(-t) \chi(-\varepsilon x, -\varepsilon zt) \gamma(-t^{-1}, -\varepsilon zt),$$

from which our theorem follows.

so

$$c\gamma(-\varepsilon t, x) \sum_y \chi(-\varepsilon x, y) \gamma(t^{-1}, y) \chi(-\varepsilon y, z) =$$

$$c\gamma(-\varepsilon t, x) \sum_y \chi(-\varepsilon x + z, y) \gamma(-\varepsilon t, yt^{-1}) =$$

$$c\gamma(-\varepsilon t, x) \frac{\alpha(-t)}{c\gamma(-\varepsilon t, x - \varepsilon z)} = \frac{\gamma(-\varepsilon t, x) \alpha(-t)}{\gamma(-\varepsilon t, x) \gamma(-\varepsilon t, -\varepsilon z) \chi(x, zt)} =$$

$$\alpha(-t) \chi(-\varepsilon x, -\varepsilon zt) \gamma(-t^{-1}, -\varepsilon zt),$$

from which our theorem follows.

so

$$c\gamma(-\varepsilon t, x) \sum_y \chi(-\varepsilon x, y) \gamma(t^{-1}, y) \chi(-\varepsilon y, z) =$$

$$c\gamma(-\varepsilon t, x) \sum_y \chi(-\varepsilon x + z, y) \gamma(-\varepsilon t, yt^{-1}) =$$

$$c\gamma(-\varepsilon t, x) \frac{\alpha(-t)}{c\gamma(-\varepsilon t, x - \varepsilon z)} = \frac{\gamma(-\varepsilon t, x) \alpha(-t)}{\gamma(-\varepsilon t, x) \gamma(-\varepsilon t, -\varepsilon z) \chi(x, zt)} =$$

$$\alpha(-t) \chi(-\varepsilon x, -\varepsilon zt) \gamma(-t^{-1}, -\varepsilon zt),$$

from which our theorem follows.

We mention an example:

We take A to be the full matrix ring $M_n(k)$, where k denotes the finite field \mathbb{F}_q with q elements, endowed with the transpose involution $*$.

It has been proved that the group $SL_*^{-1}(2, A)$ has a Bruhat presentation

Let M be the k -vector space $\bigoplus_{1 \leq i \leq m} E_i$, where $E_i = k^n$, endowed with a non-degenerate k -quadratic form Q_0 with associated k -bilinear form B_0 , defined by

$$B_0(u, v) = Q_0(u + v) - Q_0(u) - Q_0(v)$$

$$u, v \in k^n$$

This induces canonically a non degenerate A -valued quadratic form \mathbb{Q} on M

given by

1. $Q(x)_{ii} = Q_0(x_i)$
 2. $Q(x)_{ij} = B_0(x_i, x_j)$
 3. $Q(x)_{ji} = 0$
- for all $x = (x_1, \dots, x_m) \in M, 1 \leq i, j \leq n$.

and a k bilinear form \mathbb{B} on M with values in A given by $\mathbb{B}(x, y)_{ij} = B_0(x_i, y_j)$

If we pass now to the quotient modulo “anti-traces”, i.e. if we define $\overline{Q} = pr \circ Q : M \rightarrow \overline{A}$, and $\overline{\mathbb{B}} = pr \circ \mathbb{B} : M \times M \rightarrow \overline{A}$ where pr denotes the canonical projection of A onto A/A^0 , with $A^0 = \{a - a^* | a \in A\}$, We fix moreover a non trivial character ψ of k^+ and we denote by tr the usual matrix trace from A onto k .

given by

1. $Q(x)_{ii} = Q_0(x_i)$
 2. $Q(x)_{ij} = B_0(x_i, x_j)$
 3. $Q(x)_{ji} = 0$
- for all $x = (x_1, \dots, x_m) \in M, 1 \leq i, j \leq n$.

and a k bilinear form \mathbb{B} on M with values in A given by $\mathbb{B}(x, y)_{ij} = B_0(x_i, y_j)$

If we pass now to the quotient modulo “anti-traces”, i.e. if we define $\overline{Q} = pr \circ Q : M \rightarrow \overline{A}$, and $\overline{\mathbb{B}} = pr \circ \mathbb{B} : M \times M \rightarrow \overline{A}$ where pr denotes the canonical projection of A onto A/A^0 , with $A^0 = \{a - a^* | a \in A\}$, We fix moreover a non trivial character ψ of k^+ and we denote by tr the usual matrix trace from A onto k .

given by

1. $Q(x)_{ii} = Q_0(x_i)$
2. $Q(x)_{ij} = B_0(x_i, x_j)$
3. $Q(x)_{ji} = 0$
for all $x = (x_1, \dots, x_m) \in M, 1 \leq i, j \leq n$.

and a k bilinear form \mathbb{B} on M with values in A given by $\mathbb{B}(x, y)_{ij} = B_0(x_i, y_j)$

If we pass now to the quotient modulo “anti-traces”, i.e. if we define $\overline{Q} = pr \circ Q : M \rightarrow \overline{A}$, and $\overline{\mathbb{B}} = pr \circ \mathbb{B} : M \times M \rightarrow \overline{A}$ where pr denotes the canonical projection of A onto A/A^0 , with $A^0 = \{a - a^* | a \in A\}$, We fix moreover a non trivial character ψ of k^+ and we denote by tr the usual matrix trace from A onto k .

given by

1. $Q(x)_{ii} = Q_0(x_i)$
 2. $Q(x)_{ij} = B_0(x_i, x_j)$
 3. $Q(x)_{ji} = 0$
- for all $x = (x_1, \dots, x_m) \in M, 1 \leq i, j \leq n$.

and a k bilinear form \mathbb{B} on M with values in A given by $\mathbb{B}(x, y)_{ij} = B_0(x_i, y_j)$

If we pass now to the quotient modulo “anti-traces”, i.e. if we define $\overline{Q} = pr \circ Q : M \rightarrow \overline{A}$, and $\overline{\mathbb{B}} = pr \circ \mathbb{B} : M \times M \rightarrow \overline{A}$ where pr denotes the canonical projection of A onto A/A^0 , with $A^0 = \{a - a^* | a \in A\}$, We fix moreover a non trivial character ψ of k^+ and we denote by tr the usual matrix trace from A onto k .

given by

1. $Q(x)_{ii} = Q_0(x_i)$
 2. $Q(x)_{ij} = B_0(x_i, x_j)$
 3. $Q(x)_{ji} = 0$
- for all $x = (x_1, \dots, x_m) \in M, 1 \leq i, j \leq n$.

and a k bilinear form \mathbb{B} on M with values in A given by $\mathbb{B}(x, y)_{ij} = B_0(x_i, y_j)$

If we pass now to the quotient modulo “anti-traces”, i.e. if we define $\overline{Q} = pr \circ Q : M \rightarrow \overline{A}$, and $\overline{\mathbb{B}} = pr \circ \mathbb{B} : M \times M \rightarrow \overline{A}$ where pr denotes the canonical projection of A onto A/A^0 , with $A^0 = \{a - a^* | a \in A\}$, We fix moreover a non trivial character ψ of k^+ and we denote by tr the usual matrix trace from A onto k .

given by

1. $Q(x)_{ii} = Q_0(x_i)$
 2. $Q(x)_{ij} = B_0(x_i, x_j)$
 3. $Q(x)_{ji} = 0$
- for all $x = (x_1, \dots, x_m) \in M, 1 \leq i, j \leq n$.

and a k bilinear form \mathbb{B} on M with values in A given by $\mathbb{B}(x, y)_{ij} = B_0(x_i, y_j)$

If we pass now to the quotient modulo “anti-traces”, i.e. if we define $\overline{Q} = pr \circ Q : M \rightarrow \overline{A}$, and $\overline{\mathbb{B}} = pr \circ \mathbb{B} : M \times M \rightarrow \overline{A}$ where pr denotes the canonical projection of A onto A/A^0 , with $A^0 = \{a - a^* | a \in A\}$, We fix moreover a non trivial character ψ of k^+ and we denote by tr the usual matrix trace from A onto k .

Then $\underline{\psi} = \psi \circ \text{tr}$ is a non trivial character of A^+ such that $\underline{\psi}(ab) = \underline{\psi}(ba)$ and $\underline{\psi}(a^*) = \underline{\psi}(a)$ for all $a, b \in A$. On the other hand, we have

$$\underline{\psi} \circ \mathbb{Q} = \psi \circ Q$$

where Q denotes the k -valued non-degenerate quadratic form $\text{tr} \circ \mathbb{Q}$ over k , whose associated k -bilinear form will be denote by B .

We put here

1. $\gamma(s, x) = \underline{\psi}(s\overline{Q}(x))$
2. $\chi(x, y) = \underline{\psi}(s\overline{B}(x, y))$.
3. $\alpha = 1$ (when m is even)

To get a Weil representation

We put here

1. $\gamma(\mathbf{s}, x) = \underline{\psi}(\mathbf{s}\overline{\mathbf{Q}}(x))$
2. $\chi(x, y) = \underline{\psi}(\mathbf{s}\overline{\mathbf{B}}(x, y)).$
3. $\alpha = 1$ (when m is even)

To get a Weil representation

We put here

1. $\gamma(\mathbf{s}, x) = \underline{\psi}(\mathbf{s}\overline{\mathbf{Q}}(x))$
2. $\chi(x, y) = \underline{\psi}(\mathbf{s}\overline{\mathbf{B}}(x, y)).$
3. $\alpha = 1$ (*when m is even*)

To get a Weil representation

We put here

1. $\gamma(\mathbf{s}, x) = \underline{\psi}(\mathbf{s}\overline{\mathbf{Q}}(x))$
2. $\chi(x, y) = \underline{\psi}(\mathbf{s}\overline{\mathbf{B}}(x, y)).$
3. $\alpha = 1$ (*when m is even*)

To get a Weil representation

We put here

1. $\gamma(\mathbf{s}, x) = \underline{\psi}(\mathbf{s}\overline{\mathbf{Q}}(x))$
2. $\chi(x, y) = \underline{\psi}(\mathbf{s}\overline{\mathbf{B}}(x, y)).$
3. $\alpha = 1$ (*when m is even*)

To get a Weil representation

Weil representations for $G = SI_*^\varepsilon(2, A)$, G via vector bundles

- ▶ A G -Hilbert vector bundle is a tuple (E, p, B, τ) where E is the total space, B is the base, $p : E \rightarrow B$ is the projection, and $\tau = (\tau, \tau^B)$ is the action of G on E and B , respectively, such that $p \circ \tau_g^E = \tau_g^B \circ p$ for all $g \in G$. The fiber $p^{-1}(b)$ ($b \in B$) is denoted by E_b , and is a (finite dimensional) Hilbert space with inner product \langle, \rangle .
- ▶ A G -equivariant connection on G is a family of Hilbert space isomorphisms $\mathcal{H} = (E, p, B, \tau)$
 $\Gamma = \{\gamma_{b',b} \mid \gamma_{b',b} : E_b \rightarrow E_{b'}\}_{b,b' \in B}$ such that
 - $\langle \gamma_{b',b}(f), \gamma_{b',b}(h) \rangle = \langle f, h \rangle \quad (f, h \in E_b)$
 - $\langle \gamma_{b',b}(f), h \rangle = \langle f, \gamma_{b,b'}(h) \rangle \quad (f \in E_b, h \in E_{b'})$
 - $\gamma_{b,b'} \circ \gamma_{b',b} = \gamma_{b,b} = id_{E_b} \quad (b, b' \in B)$
 - $\gamma_{b',b'} \circ \gamma_{b',b} = \mu_\Gamma(b'', b', b) \gamma_{b',b} \quad (b, b', b'' \in B)$
for a suitable mapping $\mu_\Gamma : B \times B \times B \rightarrow \mathbb{C}^\times$, called the multiplier of Γ .
 - $\tau_g \circ \gamma_{b',b} = \gamma_{\tau_g(b'), \tau_g(b)} \circ \tau_g \quad (b, b' \in B, g \in G).$

Weil representations for $G = SI_*^\varepsilon(2, A)$, G via vector bundles

- ▶ A G -Hilbert vector bundle is a tuple (E, p, B, τ) where E is the total space, B is the base, $p : E \rightarrow B$ is the projection, and $\tau = (\tau, \tau^B)$ is the action of G on E and B , respectively, such that $p \circ \tau_g^E = \tau_g^B \circ p$ for all $g \in G$. The fiber $p^{-1}(b)$ ($b \in B$) is denoted by E_b , and is a (finite dimensional) Hilbert space with inner product \langle, \rangle .
- ▶ A G -equivariant connection on G is a family of Hilbert space isomorphisms $\mathcal{H} = (E, p, B, \tau)$
 $\Gamma = \{\gamma_{b',b} \mid \gamma_{b',b} : E_b \rightarrow E_{b'}\}_{b,b' \in B}$ such that
 - $\langle \gamma_{b',b}(f), \gamma_{b',b}(h) \rangle = \langle f, h \rangle \quad (f, h \in E_b)$
 - $\langle \gamma_{b',b}(f), h \rangle = \langle f, \gamma_{b,b'}(h) \rangle \quad (f \in E_b, h \in E_{b'})$
 - $\gamma_{b,b'} \circ \gamma_{b',b} = \gamma_{b,b} = id_{E_b} \quad (b, b' \in B)$
 - $\gamma_{b'',b'} \circ \gamma_{b',b} = \mu_\Gamma(b'', b', b) \gamma_{b'',b} \quad (b, b', b'' \in B)$
for a suitable mapping $\mu_\Gamma : B \times B \times B \rightarrow \mathbb{C}^\times$, called the multiplier of Γ .
- ▶ $\tau_g \circ \gamma_{b',b} = \gamma_{\tau_g(b'), \tau_g(b)} \circ \tau_g \quad (b, b' \in B, g \in G).$

Weil representations for $G = SL_*^\varepsilon(2, A)$, G via vector bundles

- ▶ A G -Hilbert vector bundle is a tuple (E, p, B, τ) where E is the total space, B is the base, $p : E \rightarrow B$ is the projection, and $\tau = (\tau, \tau^B)$ is the action of G on E and B , respectively, such that $p \circ \tau_g^E = \tau_g^B \circ p$ for all $g \in G$. The fiber $p^{-1}(b)$ ($b \in B$) is denoted by E_b , and is a (finite dimensional) Hilbert space with inner product \langle, \rangle .
- ▶ A G -equivariant connection on G is a family of Hilbert space isomorphisms $\mathcal{H} = (E, p, B, \tau)$
 $\Gamma = \{\gamma_{b',b} \mid \gamma_{b',b} : E_b \rightarrow E_{b'}\}_{b,b' \in B}$ such that
 - $\langle \gamma_{b',b}(f), \gamma_{b',b}(h) \rangle = \langle f, h \rangle \quad (f, h \in E_b)$
 - $\langle \gamma_{b',b}(f), h \rangle = \langle f, \gamma_{b,b'}(h) \rangle \quad (f \in E_b, h \in E_{b'})$
 - $\gamma_{b,b'} \circ \gamma_{b',b} = \gamma_{b,b} = id_{E_b} \quad (b, b' \in B)$
 - $\gamma_{b'',b'} \circ \gamma_{b',b} = \mu_\Gamma(b'', b', b) \gamma_{b'',b} \quad (b, b', b'' \in B)$
for a suitable mapping $\mu_\Gamma : B \times B \times B \rightarrow \mathbb{C}^\times$, called the multiplier of Γ .
- ▶ $\tau_g \circ \gamma_{b',b} = \gamma_{\tau_g(b'), \tau_g(b)} \circ \tau_g \quad (b, b' \in B, g \in G).$

Weil representations for $G = SI_*^\varepsilon(2, A)$, G via vector bundles

- ▶ A G -Hilbert vector bundle is a tuple (E, p, B, τ) where E is the total space, B is the base, $p : E \rightarrow B$ is the projection, and $\tau = (\tau, \tau^B)$ is the action of G on E and B , respectively, such that $p \circ \tau_g^E = \tau_g^B \circ p$ for all $g \in G$. The fiber $p^{-1}(b)$ ($b \in B$) is denoted by E_b , and is a (finite dimensional) Hilbert space with inner product \langle, \rangle .
- ▶ A G -equivariant connection on G is a family of Hilbert space isomorphisms $\mathcal{H} = (E, p, B, \tau)$
 $\Gamma = \{ \gamma_{b',b} \mid \gamma_{b',b} : E_b \rightarrow E_{b'} \}_{b,b' \in B}$ such that
 - $\langle \gamma_{b',b}(f), \gamma_{b',b}(h) \rangle = \langle f, h \rangle \quad (f, h \in E_b)$
 - $\langle \gamma_{b',b}(f), h \rangle = \langle f, \gamma_{b,b'}(h) \rangle \quad (f \in E_b, h \in E_{b'})$
 - $\gamma_{b,b'} \circ \gamma_{b',b} = \gamma_{b,b} = id_{E_b} \quad (b, b' \in B)$
 - $\gamma_{b'',b'} \circ \gamma_{b',b} = \mu_\Gamma(b'', b', b) \gamma_{b'',b} \quad (b, b', b'' \in B)$
for a suitable mapping $\mu_\Gamma : B \times B \times B \rightarrow \mathbb{C}^\times$, called the multiplier of Γ .
 - $\tau_g \circ \gamma_{b',b} = \gamma_{\tau_g(b'), \tau_g(b)} \circ \tau_g \quad (b, b' \in B, g \in G).$

Weil representations for $G = SL_*^\varepsilon(2, A)$, G via vector bundles

- ▶ A G -Hilbert vector bundle is a tuple (E, p, B, τ) where E is the total space, B is the base, $p : E \rightarrow B$ is the projection, and $\tau = (\tau, \tau^B)$ is the action of G on E and B , respectively, such that $p \circ \tau_g^E = \tau_g^B \circ p$ for all $g \in G$. The fiber $p^{-1}(b)$ ($b \in B$) is denoted by E_b , and is a (finite dimensional) Hilbert space with inner product \langle, \rangle .
- ▶ A G -equivariant connection on G is a family of Hilbert space isomorphisms $\mathcal{H} = (E, p, B, \tau)$
 $\Gamma = \{\gamma_{b',b} \mid \gamma_{b',b} : E_b \rightarrow E_{b'}\}_{b,b' \in B}$ such that
 - $\langle \gamma_{b',b}(f), \gamma_{b',b}(h) \rangle = \langle f, h \rangle \quad (f, h \in E_b)$
 - $\langle \gamma_{b',b}(f), h \rangle = \langle f, \gamma_{b,b'}(h) \rangle \quad (f \in E_b, h \in E_{b'})$
 - $\gamma_{b,b'} \circ \gamma_{b',b} = \gamma_{b,b} = id_{E_b} \quad (b, b' \in B)$
 - $\gamma_{b'',b'} \circ \gamma_{b',b} = \mu_\Gamma(b'', b', b) \gamma_{b'',b} \quad (b, b', b'' \in B)$
for a suitable mapping $\mu_\Gamma : B \times B \times B \rightarrow \mathbb{C}^\times$, called the multiplier of Γ .
 - $\tau_g \circ \gamma_{b',b} = \gamma_{\tau_g(b'), \tau_g(b)} \circ \tau_g \quad (b, b' \in B, g \in G).$

Weil representations for $G = SI_*^\varepsilon(2, A)$, G via vector bundles

- ▶ A G -Hilbert vector bundle is a tuple (E, p, B, τ) where E is the total space, B is the base, $p : E \rightarrow B$ is the projection, and $\tau = (\tau, \tau^B)$ is the action of G on E and B , respectively, such that $p \circ \tau_g^E = \tau_g^B \circ p$ for all $g \in G$. The fiber $p^{-1}(b)$ ($b \in B$) is denoted by E_b , and is a (finite dimensional) Hilbert space with inner product \langle, \rangle .
- ▶ A G -equivariant connection on G is a family of Hilbert space isomorphisms $\mathcal{H} = (E, p, B, \tau)$
 $\Gamma = \{\gamma_{b',b} \mid \gamma_{b',b} : E_b \rightarrow E_{b'}\}_{b,b' \in B}$ such that
 - $\langle \gamma_{b',b}(f), \gamma_{b',b}(h) \rangle = \langle f, h \rangle \quad (f, h \in E_b)$
 - $\langle \gamma_{b',b}(f), h \rangle = \langle f, \gamma_{b,b'}(h) \rangle \quad (f \in E_b, h \in E_{b'})$
 - $\gamma_{b,b'} \circ \gamma_{b',b} = \gamma_{b,b} = id_{E_b} \quad (b, b' \in B)$
 - $\gamma_{b'',b'} \circ \gamma_{b',b} = \mu_\Gamma(b'', b', b) \gamma_{b'',b} \quad (b, b', b'' \in B)$for a suitable mapping $\mu_\Gamma : B \times B \times B \rightarrow \mathbb{C}^\times$, called the multiplier of Γ .

v. $\tau_g \circ \gamma_{b',b} = \gamma_{\tau_g(b'), \tau_g(b)} \circ \tau_g \quad (b, b' \in B, g \in G).$

Weil representations for $G = SL_*^\varepsilon(2, A)$, G via vector bundles

- ▶ A G -Hilbert vector bundle is a tuple (E, p, B, τ) where E is the total space, B is the base, $p : E \rightarrow B$ is the projection, and $\tau = (\tau, \tau^B)$ is the action of G on E and B , respectively, such that $p \circ \tau_g^E = \tau_g^B \circ p$ for all $g \in G$. The fiber $p^{-1}(b)$ ($b \in B$) is denoted by E_b , and is a (finite dimensional) Hilbert space with inner product \langle, \rangle .
- ▶ A G -equivariant connection on G is a family of Hilbert space isomorphisms $\mathcal{H} = (E, p, B, \tau)$
 $\Gamma = \{ \gamma_{b',b} \mid \gamma_{b',b} : E_b \rightarrow E_{b'} \}_{b,b' \in B}$ such that
 - $\langle \gamma_{b',b}(f), \gamma_{b',b}(h) \rangle = \langle f, h \rangle \quad (f, h \in E_b)$
 - $\langle \gamma_{b',b}(f), h \rangle = \langle f, \gamma_{b,b'}(h) \rangle \quad (f \in E_b, h \in E_{b'})$
 - $\gamma_{b,b'} \circ \gamma_{b',b} = \gamma_{b,b} = id_{E_b} \quad (b, b' \in B)$
 - $\gamma_{b'',b'} \circ \gamma_{b',b} = \mu_\Gamma(b'', b', b) \gamma_{b'',b} \quad (b, b', b'' \in B)$
for a suitable mapping $\mu_\Gamma : B \times B \times B \rightarrow \mathbb{C}^\times$, called the multiplier of Γ .
- ▶ $\tau_g \circ \gamma_{b',b} = \gamma_{\tau_g(b'), \tau_g(b)} \circ \tau_g \quad (b, b' \in B, g \in G).$

Representation of G by contraction of the fiber bundle over b

- ▶ The projective unitary representation (V_b, ρ^b) of G is defined as follows:

- $V_b = E_b$ as a Hilbert space,
- $\rho_g^b(v) = \gamma_{b, \tau_g^B(b)} \tau_g^E(v)$ for all $g \in G, v \in V_b$

whose cocycle c is given by

$c(g, h) = \mu_\Gamma(b, g.b, gh.b)$ for all $g, h \in G$. If Γ is flat, i.e., if the function μ_Γ is 1, then ρ^b is a true representation of G .

- ▶ We briefly present now a construction of $G = SI_{*}^{-}(2, A)$ using Lagrangian fiber bundles:

Representation of G by contraction of the fiber bundle over b

- ▶ The projective unitary representation (V_b, ρ^b) of G is defined as follows:
 - i. $V_b = E_b$ as a Hilbert space,
 - ii. $\rho_g^b(v) = \gamma_{b, \tau_g^B(b)} \tau_g^E(v)$ for all $g \in G, v \in V_b$

whose cocycle c is given by

$c(g, h) = \mu_\Gamma(b, g.b, gh.b)$ for all $g, h \in G$. If Γ is flat, i.e., if the function μ_Γ is 1, then ρ^b is a true representation of G .

- ▶ We briefly present now a construction of $G = SI_{*}^{-}(2, A)$ using Lagrangian fiber bundles:

Representation of G by contraction of the fiber bundle over b

- ▶ The projective unitary representation (V_b, ρ^b) of G is defined as follows:
 - i. $V_b = E_b$ as a Hilbert space,
 - ii. $\rho_g^b(v) = \gamma_{b, \tau_g^B(b)} \tau_g^E(v)$ for all $g \in G, v \in V_b$

whose cocycle c is given by

$c(g, h) = \mu_\Gamma(b, g.b, gh.b)$ for all $g, h \in G$. If Γ is flat, i.e., if the function μ_Γ is 1, then ρ^b is a true representation of G .

- ▶ We briefly present now a construction of $G = SI_{*}^{-}(2, A)$ using Lagrangian fiber bundles:

Representation of G by contraction of the fiber bundle over b

- ▶ The projective unitary representation (V_b, ρ^b) of G is defined as follows:
 - i. $V_b = E_b$ as a Hilbert space,
 - ii. $\rho_g^b(v) = \gamma_{b, \tau_g^B(b)} \tau_g^E(v)$ for all $g \in G, v \in V_b$

whose cocycle c is given by

$c(g, h) = \mu_\Gamma(b, g.b, gh.b)$ for all $g, h \in G$. If Γ is flat, i.e., if the function μ_Γ is 1, then ρ^b is a true representation of G .

- ▶ We briefly present now a construction of $G = SI_{*}^{-}(2, A)$ using Lagrangian fiber bundles:

Representation of G by contraction of the fiber bundle over b

- ▶ The projective unitary representation (V_b, ρ^b) of G is defined as follows:
 - i. $V_b = E_b$ as a Hilbert space,
 - ii. $\rho_g^b(v) = \gamma_{b, \tau_g^B(b)} \tau_g^E(v)$ for all $g \in G, v \in V_b$

whose cocycle c is given by

$c(g, h) = \mu_\Gamma(b, g.b, gh.b)$ for all $g, h \in G$. If Γ is flat, i.e., if the function μ_Γ is 1, then ρ^b is a true representation of G .

- ▶ We briefly present now a construction of $G = SL_*^-(2, A)$ using Lagrangian fiber bundles:

Weil representation of G by contraction of Lagrangian fiber bundles

- ▶ We assume that A is a finite k -algebra with an involution that fixes the finite base field.
- ▶ Let S be a left A -module (which is right A -module via $s.a = a^*.s$ for $a \in A, s \in S$), finite dimensional as k -vector space
- ▶ Let $\eta : S \times S \rightarrow k$. be a non-degenerate, k -bilinear, symmetric A -balanced pairing (i.e. $\eta(s.a, t) = \eta(s, a.t)$ for $a \in A, s, t \in S$)
- ▶ We set $W = S \oplus S$ and we define a symplectic form B on W by

$$B((s, t), (s', t')) = \eta(s, t') - \eta(t, s')$$

for all $(s, t), (s', t') \in W$. We fix a non trivial character ψ of the additive group k^+ of k and we put $\chi = \psi \circ B$.

Weil representation of G by contraction of Lagrangian fiber bundles

- ▶ We assume that A is a finite k -algebra with an involution that fixes the finite base field.
- ▶ Let S be a left A -module (which is right A -module via $s.a = a^*.s$ for $a \in A, s \in S$), finite dimensional as k -vector space
- ▶ Let $\eta : S \times S \rightarrow k$. be a non-degenerate, k -bilinear, symmetric A -balanced pairing (i.e. $\eta(s.a, t) = \eta(s, a.t)$ for $a \in A, s, t \in S$)
- ▶ We set $W = S \oplus S$ and we define a symplectic form B on W by

$$B((s, t), (s', t')) = \eta(s, t') - \eta(t, s')$$

for all $(s, t), (s', t') \in W$. We fix a non trivial character ψ of the additive group k^+ of k and we put $\chi = \psi \circ B$.

Weil representation of G by contraction of Lagrangian fiber bundles

- ▶ We assume that A is a finite k -algebra with an involution that fixes the finite base field.
- ▶ Let S be a left A -module (which is right A -module via $s.a = a^*.s$ for $a \in A, s \in S$), finite dimensional as k -vector space
- ▶ Let $\eta : S \times S \rightarrow k$ be a non-degenerate, k -bilinear, symmetric A -balanced pairing (i.e. $\eta(s.a, t) = \eta(s, a.t)$ for $a \in A, s, t \in S$)
- ▶ We set $W = S \oplus S$ and we define a symplectic form B on W by

$$B((s, t), (s', t')) = \eta(s, t') - \eta(t, s')$$

for all $(s, t), (s', t') \in W$. We fix a non trivial character ψ of the additive group k^+ of k and we put $\chi = \psi \circ B$.

Weil representation of G by contraction of Lagrangian fiber bundles

- ▶ We assume that A is a finite k -algebra with an involution that fixes the finite base field.
- ▶ Let S be a left A -module (which is right A -module via $s.a = a^*.s$ for $a \in A, s \in S$), finite dimensional as k -vector space
- ▶ Let $\eta : S \times S \rightarrow k$ be a non-degenerate, k -bilinear, symmetric A -balanced pairing (i.e. $\eta(s.a, t) = \eta(s, a.t)$ for $a \in A, s, t \in S$)
- ▶ We set $W = S \oplus S$ and we define a symplectic form B on W by

$$B((s, t), (s', t')) = \eta(s, t') - \eta(t, s')$$

for all $(s, t), (s', t') \in W$. We fix a non trivial character ψ of the additive group k^+ of k and we put $\chi = \psi \circ B$.

- ▶ A Lagrangian L in W is a A -submodule which is maximal totally isotropic for B , i.e., $L = L^\perp$
- ▶ The group G acts on W by matrix multiplication, and we have $B(gw, gw') = B(w, w')$
- ▶ Then the Lagrangian bundle of G associated to S is given by

- i B is the set of all Lagrangians of $W = S \oplus S$
- ii E is the disjoint union of the spaces

$$E_L = \{f : W \rightarrow \mathbb{C} \mid f(w + \zeta) = \chi(w, \zeta)f(w); \quad w \in W, \zeta \in L\},$$

for $L \in B$, each endowed with the inner product given by

$$\langle f, h \rangle = \sum_{w \in W} f(w) \overline{h(w)}$$

- ▶ A Lagrangian L in W is a A -submodule which is maximal totally isotropic for B , i.e., $L = L^\perp$
- ▶ The group G acts on W by matrix multiplication, and we have $B(gw, gw') = B(w, w')$
- ▶ Then the Lagrangian bundle of G associated to S is given by

- i B is the set of all Lagrangians of $W = S \oplus S$
- ii E is the disjoint union of the spaces

$$E_L = \{f : W \rightarrow \mathbb{C} \mid f(w + \zeta) = \chi(w, \zeta)f(w); \quad w \in W, \zeta \in L\},$$

for $L \in B$, each endowed with the inner product given by

$$\langle f, h \rangle = \sum_{w \in W} f(w) \overline{h(w)}$$

- ▶ A Lagrangian L in W is a A -submodule which is maximal totally isotropic for B , i.e., $L = L^\perp$
- ▶ The group G acts on W by matrix multiplication, and we have $B(gw, gw') = B(w, w')$
- ▶ Then the Lagrangian bundle of G associated to S is given by

- i \mathcal{B} is the set of all Lagrangians of $W = S \oplus S$
- ii E is the disjoint union of the spaces

$$E_L = \{f : W \rightarrow \mathbb{C} \mid f(w + \zeta) = \chi(w, \zeta)f(w); \quad w \in W, \zeta \in L\},$$

for $L \in \mathcal{B}$, each endowed with the inner product given by

$$\langle f, h \rangle = \sum_{w \in W} f(w) \overline{h(w)}$$

- ▶ A Lagrangian L in W is a A -submodule which is maximal totally isotropic for B , i.e., $L = L^\perp$
- ▶ The group G acts on W by matrix multiplication, and we have $B(gw, gw') = B(w, w')$
- ▶ Then the Lagrangian bundle of G associated to S is given by
 - \mathcal{B} is the set of all Lagrangians of $W = S \oplus S$
 - E is the disjoint union of the spaces

$$E_L = \{f : W \rightarrow \mathbb{C} \mid f(w + \zeta) = \chi(w, \zeta)f(w); \quad w \in W, \zeta \in L\},$$

for $L \in \mathcal{B}$, each endowed with the inner product given by

$$\langle f, h \rangle = \sum_{w \in W} f(w) \overline{h(w)}$$

- ▶ A Lagrangian L in W is a A -submodule which is maximal totally isotropic for B , i.e., $L = L^\perp$
- ▶ The group G acts on W by matrix multiplication, and we have $B(gw, gw') = B(w, w')$
- ▶ Then the Lagrangian bundle of G associated to S is given by
 - i \mathcal{B} is the set of all Lagrangians of $W = S \oplus S$
 - ii E is the disjoint union of the spaces

$$E_L = \{f : W \rightarrow \mathbb{C} \mid f(w + \zeta) = \chi(w, \zeta)f(w); \ w \in W, \ \zeta \in L\},$$

for $L \in \mathcal{B}$, each endowed with the inner product given by

$$\langle f, h \rangle = \sum_{w \in W} f(w) \overline{h(w)}$$

- iii $p : E \rightarrow \mathcal{B}$ is given by $p(f) = L$ if $f \in E_L$
- iv τ denotes the action of G in E and \mathcal{B} given by

$$(\tau_g(f))(w) = f(g^{-1}w), \tau_g(L) = g(L),$$

for $g \in G, f \in E, w \in W, L \in \mathcal{B}$.

iii $p : E \rightarrow \mathcal{B}$ is given by $p(f) = L$ if $f \in E_L$

iv τ denotes the action of G in E and \mathcal{B} given by

$$(\tau_g(f))(w) = f(g^{-1}w), \tau_g(L) = g(L),$$

for $g \in G, f \in E, w \in W, L \in \mathcal{B}$.

- iii $p : E \rightarrow \mathcal{B}$ is given by $p(f) = L$ if $f \in E_L$
- iv τ denotes the action of G in E and \mathcal{B} given by

$$(\tau_g(f))(w) = f(g^{-1}w), \tau_g(L) = g(L),$$

for $g \in G, f \in E, w \in W, L \in \mathcal{B}$.

G -equivariant connections

Assume that $2 \in A$ is invertible. Then the family

$$\Gamma = \{\gamma_{L',L} \mid \gamma_{L',L} : E_L \rightarrow E_{L'}\}_{L,L' \in \mathcal{B}}$$

of linear isomorphisms

$$\gamma_{L',L}(f)(w) = \frac{1}{\sqrt{|L||L \cap L'|}} \sum_{\zeta' \in L'} \overline{\chi(w, \zeta')} f(w + \zeta'), \quad (f \in E_L, w \in W)$$

is a G -equivariant connection with multiplier

$$\mu_\Gamma(L'', L', L) = \sqrt{\frac{|L'' \cap L'|}{|L \cap L''||L' \cap L||L|}} S_W(L; L', L'') \quad (L, L', L'' \in \mathcal{B}),$$

where the geometric Gauss sum $S_W(L; L', L'')$ is given by

$$S_W(L; L', L'') = \sum_{\zeta \in L \cap (L' + L'')} \chi(\zeta')$$

where $\zeta \in L \cap (L' + L'')$ is written as $\zeta' + \zeta''$ with $\zeta' \in L', \zeta'' \in L''$.

References

-  L. Gutiérrez, *A generalized Weil Representation for $SL_*(2, A_m)$, where $A_m = F_q / \langle x^m \rangle$* , J. of Algebra, 322, (2009) 42-53.
-  L. Gutiérrez, J. Pantoja, J. Soto-Andrade, *On generalized Weil Representations over Involutive Rings*, Contemporary Math., 544, (2011) 109-121.
-  L. Gutiérrez, J. Pantoja, J. Soto-Andrade, *Geometric Weil Representations for star-analogues of $SL(2, K)$* , Contemporary Math., 537, (2011) 211-226.
-  J. Pantoja, *A presentation of the group $SL_*(2, A)$, A a simple artinian ring with involution*, Manuscripta Math., 121, (2006), 97-104
-  J. Pantoja, J. Soto-Andrade, *A Bruhat decomposition of the group $SL_*(2, A)$* , J. of Algebra, 262 (2003), 401-412

References

-  L. Gutiérrez, *A generalized Weil Representation for $SL_*(2, A_m)$, where $A_m = F_q / \langle x^m \rangle$* , J. of Algebra, 322, (2009) 42-53.
-  L. Gutiérrez, J. Pantoja, J. Soto-Andrade, *On generalized Weil Representations over Involutive Rings*, Contemporary Math., 544, (2011) 109-121.
-  L. Gutiérrez, J. Pantoja, J. Soto-Andrade, *Geometric Weil Representations for star-analogues of $SL(2, K)$* , Contemporary Math., 537, (2011) 211-226.
-  J. Pantoja, *A presentation of the group $SL_*(2, A)$, A a simple artinian ring with involution*, Manuscripta Math., 121, (2006), 97-104
-  J. Pantoja, J. Soto-Andrade, *A Bruhat decomposition of the group $SL_*(2, A)$* , J. of Algebra, 262 (2003), 401-412

References

-  L. Gutiérrez, *A generalized Weil Representation for $SL_*(2, A_m)$, where $A_m = F_q / \langle x^m \rangle$* , J. of Algebra, 322, (2009) 42-53.
-  L. Gutiérrez, J. Pantoja, J. Soto-Andrade, *On generalized Weil Representations over Involutive Rings*, Contemporary Math., 544, (2011) 109-121.
-  L. Gutiérrez, J. Pantoja, J. Soto-Andrade, *Geometric Weil Representations for star-analogues of $SL(2, K)$* , Contemporary Math., 537, (2011) 211-226.
-  J. Pantoja, *A presentation of the group $SL_*(2, A)$, A a simple artinian ring with involution*, Manuscripta Math., 121, (2006), 97-104
-  J. Pantoja, J. Soto-Andrade, *A Bruhat decomposition of the group $SL_*(2, A)$* , J. of Algebra, 262 (2003), 401-412

References

-  L. Gutiérrez, *A generalized Weil Representation for $SL_*(2, A_m)$, where $A_m = F_q / \langle x^m \rangle$* , J. of Algebra, 322, (2009) 42-53.
-  L. Gutiérrez, J. Pantoja, J. Soto-Andrade, *On generalized Weil Representations over Involutive Rings*, Contemporary Math., 544, (2011) 109-121.
-  L. Gutiérrez, J. Pantoja, J. Soto-Andrade, *Geometric Weil Representations for star-analogues of $SL(2, K)$* , Contemporary Math., 537, (2011) 211-226.
-  J. Pantoja, *A presentation of the group $SL_*(2, A)$, A a simple artinian ring with involution*, Manuscripta Math., 121, (2006), 97-104
-  J. Pantoja, J. Soto-Andrade, *A Bruhat decomposition of the group $SL_*(2, A)$* , J. of Algebra, 262 (2003), 401-412

References

-  L. Gutiérrez, *A generalized Weil Representation for $SL_*(2, A_m)$, where $A_m = F_q / \langle x^m \rangle$* , J. of Algebra, 322, (2009) 42-53.
-  L. Gutiérrez, J. Pantoja, J. Soto-Andrade, *On generalized Weil Representations over Involutive Rings*, Contemporary Math., 544, (2011) 109-121.
-  L. Gutiérrez, J. Pantoja, J. Soto-Andrade, *Geometric Weil Representations for star-analogues of $SL(2, K)$* , Contemporary Math., 537, (2011) 211-226.
-  J. Pantoja, *A presentation of the group $SL_*(2, A)$, A a simple artinian ring with involution*, Manuscripta Math., 121, (2006), 97-104
-  J. Pantoja, J. Soto-Andrade, *A Bruhat decomposition of the group $SL_*(2, A)$* , J. of Algebra, 262 (2003), 401-412