

Extensions of tensor categories

Sonia Natale¹ Alain Bruguières²

¹Universidad Nacional de Córdoba. CIEM-CONICET.

²Université Montpellier II.

Representation Theory of Groups, Lie Algebras, and Hopf
Algebras

CMS Summer Meeting - Regina 2012

Outline

Exact sequences of tensor categories

Definitions

Examples

Exact sequences and Hopf monads

Classification of exact sequences

Applications

Tensor categories

Let k be a field.

Tensor categories

Let k be a field.

A *tensor category over k* is a k -linear abelian rigid monoidal category \mathcal{C} where:

- ▶ Hom spaces are finite dimensional.
- ▶ Objects have finite length.
- ▶ The monoidal product $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ is k -bilinear.
- ▶ The unit object 1 is scalar.

Tensor categories

Let k be a field.

A *tensor category over k* is a k -linear abelian rigid monoidal category \mathcal{C} where:

- ▶ Hom spaces are finite dimensional.
- ▶ Objects have finite length.
- ▶ The monoidal product $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ is k -bilinear.
- ▶ The unit object $\mathbb{1}$ is scalar.

If \mathcal{C} is a tensor category, \otimes is bi-exact and $\mathbb{1}$ is simple.

Tensor categories

Let k be a field.

A *tensor category over k* is a k -linear abelian rigid monoidal category \mathcal{C} where:

- ▶ Hom spaces are finite dimensional.
- ▶ Objects have finite length.
- ▶ The monoidal product $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ is k -bilinear.
- ▶ The unit object $\mathbb{1}$ is scalar.

If \mathcal{C} is a tensor category, \otimes is bi-exact and $\mathbb{1}$ is simple.

A tensor category \mathcal{C} is *finite* if it is finite as a k -linear abelian category.

Tensor categories

Let k be a field.

A *tensor category over k* is a k -linear abelian rigid monoidal category \mathcal{C} where:

- ▶ Hom spaces are finite dimensional.
- ▶ Objects have finite length.
- ▶ The monoidal product $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ is k -bilinear.
- ▶ The unit object $\mathbb{1}$ is scalar.

If \mathcal{C} is a tensor category, \otimes is bi-exact and $\mathbb{1}$ is simple.

A tensor category \mathcal{C} is *finite* if it is finite as a k -linear abelian category.

A *tensor functor* $F : \mathcal{C} \rightarrow \mathcal{D}$ is a k -linear exact strong monoidal functor.

Tensor categories

Let k be a field.

A *tensor category over k* is a k -linear abelian rigid monoidal category \mathcal{C} where:

- ▶ Hom spaces are finite dimensional.
- ▶ Objects have finite length.
- ▶ The monoidal product $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ is k -bilinear.
- ▶ The unit object $\mathbb{1}$ is scalar.

If \mathcal{C} is a tensor category, \otimes is bi-exact and $\mathbb{1}$ is simple.

A tensor category \mathcal{C} is *finite* if it is finite as a k -linear abelian category.

A *tensor functor* $F : \mathcal{C} \rightarrow \mathcal{D}$ is a k -linear exact strong monoidal functor.

A tensor functor preserves duals. Moreover it is automatically faithful.

Normal tensor functors

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a tensor functor.

Normal tensor functors

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a tensor functor.

F is *dominant* if any object Y of \mathcal{D} is a subobject of $F(X)$, for some X in \mathcal{C} .

Normal tensor functors

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a tensor functor.

F is *dominant* if any object Y of \mathcal{D} is a subobject of $F(X)$, for some X in \mathcal{C} .

$\mathcal{Ret}_F := F^{-1}(\langle \mathbb{1} \rangle) \subseteq \mathcal{C}$: full tensor subcategory of objects X of \mathcal{C} such that $F(X)$ is a trivial object of \mathcal{D} .

Normal tensor functors

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a tensor functor.

F is *dominant* if any object Y of \mathcal{D} is a subobject of $F(X)$, for some X in \mathcal{C} .

$\mathcal{Ker}_F := F^{-1}(\langle 1 \rangle) \subseteq \mathcal{C}$: full tensor subcategory of objects X of \mathcal{C} such that $F(X)$ is a trivial object of \mathcal{D} .

Definition

A tensor functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is *normal* if for any object X of \mathcal{C} , there exists a subobject $X_0 \subset X$, such that $F(X_0)$ is the largest trivial subobject of $F(X)$.

Normal tensor functors

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a tensor functor.

F is *dominant* if any object Y of \mathcal{D} is a subobject of $F(X)$, for some X in \mathcal{C} .

$\mathcal{Ker}_F := F^{-1}(\langle 1 \rangle) \subseteq \mathcal{C}$: full tensor subcategory of objects X of \mathcal{C} such that $F(X)$ is a trivial object of \mathcal{D} .

Definition

A tensor functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is *normal* if for any object X of \mathcal{C} , there exists a subobject $X_0 \subset X$, such that $F(X_0)$ is the largest trivial subobject of $F(X)$.

Characterizations:

Normal tensor functors

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a tensor functor.

F is *dominant* if any object Y of \mathcal{D} is a subobject of $F(X)$, for some X in \mathcal{C} .

$\mathcal{Ker}_F := F^{-1}(\langle \mathbb{I} \rangle) \subseteq \mathcal{C}$: full tensor subcategory of objects X of \mathcal{C} such that $F(X)$ is a trivial object of \mathcal{D} .

Definition

A tensor functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is *normal* if for any object X of \mathcal{C} , there exists a subobject $X_0 \subset X$, such that $F(X_0)$ is the largest trivial subobject of $F(X)$.

Characterizations:

- Suppose F admits a right (or left) adjoint R . Then:
 F is normal $\Leftrightarrow R(\mathbb{I})$ belongs to \mathcal{Ker}_F .

Normal tensor functors

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a tensor functor.

F is *dominant* if any object Y of \mathcal{D} is a subobject of $F(X)$, for some X in \mathcal{C} .

$\mathfrak{Ker}_F := F^{-1}(\langle \mathbb{1} \rangle) \subseteq \mathcal{C}$: full tensor subcategory of objects X of \mathcal{C} such that $F(X)$ is a trivial object of \mathcal{D} .

Definition

A tensor functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is *normal* if for any object X of \mathcal{C} , there exists a subobject $X_0 \subset X$, such that $F(X_0)$ is the largest trivial subobject of $F(X)$.

Characterizations:

- ▶ Suppose F admits a right (or left) adjoint R . Then:
 F is normal $\Leftrightarrow R(\mathbb{1})$ belongs to \mathfrak{Ker}_F .
- ▶ Suppose \mathcal{C}, \mathcal{D} are fusion categories. F is normal iff
 \forall simple object X , $\text{Hom}(\mathbb{1}, F(X)) \neq 0 \Rightarrow X \in \mathfrak{Ker}_F$.

Exact sequences of tensor categories

Let $\mathcal{C}', \mathcal{C}, \mathcal{C}''$ be tensor categories over k .

Exact sequences of tensor categories

Let $\mathcal{C}', \mathcal{C}, \mathcal{C}''$ be tensor categories over k .

A sequence of tensor functors

$$\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$$

is an *exact sequence of tensor categories* if the following hold:

Exact sequences of tensor categories

Let $\mathcal{C}', \mathcal{C}, \mathcal{C}''$ be tensor categories over k .

A sequence of tensor functors

$$\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$$

is an *exact sequence of tensor categories* if the following hold:

- ▶ The tensor functor F is dominant and normal;

Exact sequences of tensor categories

Let $\mathcal{C}', \mathcal{C}, \mathcal{C}''$ be tensor categories over k .

A sequence of tensor functors

$$\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$$

is an *exact sequence of tensor categories* if the following hold:

- ▶ The tensor functor F is dominant and normal;
- ▶ The tensor functor f is a full embedding;

Exact sequences of tensor categories

Let $\mathcal{C}', \mathcal{C}, \mathcal{C}''$ be tensor categories over k .

A sequence of tensor functors

$$\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$$

is an *exact sequence of tensor categories* if the following hold:

- ▶ The tensor functor F is dominant and normal;
- ▶ The tensor functor f is a full embedding;
- ▶ The essential image of f is $\mathcal{Ker} F$;

Exact sequences of tensor categories

Let $\mathcal{C}', \mathcal{C}, \mathcal{C}''$ be tensor categories over k .

A sequence of tensor functors

$$\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$$

is an *exact sequence of tensor categories* if the following hold:

- ▶ The tensor functor F is dominant and normal;
- ▶ The tensor functor f is a full embedding;
- ▶ The essential image of f is $\mathcal{R}et_F$;

Comments

Exact sequences of tensor categories

Let $\mathcal{C}', \mathcal{C}, \mathcal{C}''$ be tensor categories over k .

A sequence of tensor functors

$$\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$$

is an *exact sequence of tensor categories* if the following hold:

- ▶ The tensor functor F is dominant and normal;
- ▶ The tensor functor f is a full embedding;
- ▶ The essential image of f is \mathcal{Ker}_F ;

Comments

- ▶ $F : \mathcal{C} \rightarrow \mathcal{C}''$ normal dominant \leadsto exact sequence

$$\mathcal{Ker}_F \rightarrow \mathcal{C} \xrightarrow{F} \mathcal{C}''.$$

Exact sequences of tensor categories

Let $\mathcal{C}', \mathcal{C}, \mathcal{C}''$ be tensor categories over k .

A sequence of tensor functors

$$\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$$

is an *exact sequence of tensor categories* if the following hold:

- ▶ The tensor functor F is dominant and normal;
- ▶ The tensor functor f is a full embedding;
- ▶ The essential image of f is \mathcal{Ker}_F ;

Comments

- ▶ $F : \mathcal{C} \rightarrow \mathcal{C}''$ normal dominant \leadsto exact sequence

$$\mathcal{Ker}_F \rightarrow \mathcal{C} \xrightarrow{F} \mathcal{C}''.$$

- ▶ Suppose $\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$ exact $\Rightarrow \exists$ a Hopf algebra H st:

$$\mathcal{C}' \simeq \text{comod-}H.$$

Group extensions

Let

$$1 \rightarrow G'' \xrightarrow{\iota} G \xrightarrow{\pi} G' \rightarrow 1$$

be an exact sequence of finite groups.

Group extensions

Let

$$1 \rightarrow G'' \xrightarrow{\iota} G \xrightarrow{\pi} G' \rightarrow 1$$

be an exact sequence of finite groups.

This gives rise to exact sequences of tensor categories:

Group extensions

Let

$$1 \rightarrow G'' \xrightarrow{\iota} G \xrightarrow{\pi} G' \rightarrow 1$$

be an exact sequence of finite groups.

This gives rise to exact sequences of tensor categories:

$$\mathrm{rep} G' \xrightarrow{\pi^*} \mathrm{rep} G \xrightarrow{\iota^*} \mathrm{rep} G''.$$

Group extensions

Let

$$1 \rightarrow G'' \xrightarrow{\iota} G \xrightarrow{\pi} G' \rightarrow 1$$

be an exact sequence of finite groups.

This gives rise to exact sequences of tensor categories:

$$\mathrm{rep} G' \xrightarrow{\pi^*} \mathrm{rep} G \xrightarrow{\iota^*} \mathrm{rep} G''.$$

and

Group extensions

Let

$$1 \rightarrow G'' \xrightarrow{\iota} G \xrightarrow{\pi} G' \rightarrow 1$$

be an exact sequence of finite groups.

This gives rise to exact sequences of tensor categories:

$$\mathrm{rep} G' \xrightarrow{\pi^*} \mathrm{rep} G \xrightarrow{\iota^*} \mathrm{rep} G''.$$

and

$$\mathcal{C}(G'') \rightarrow \mathcal{C}(G) \rightarrow \mathcal{C}(G'),$$

Group extensions

Let

$$1 \rightarrow G'' \xrightarrow{\iota} G \xrightarrow{\pi} G' \rightarrow 1$$

be an exact sequence of finite groups.

This gives rise to exact sequences of tensor categories:

$$\operatorname{rep} G' \xrightarrow{\pi^*} \operatorname{rep} G \xrightarrow{\iota^*} \operatorname{rep} G''.$$

and

$$\mathcal{C}(G'') \rightarrow \mathcal{C}(G) \rightarrow \mathcal{C}(G'),$$

where $\mathcal{C}(G)$: finite dimensional G -graded vector spaces over k .

Equivariantization

Let \mathcal{C} be a tensor category over k , G a group.

Equivariantization

Let \mathcal{C} be a tensor category over k , G a group.

$\underline{\text{Aut}}_{\otimes} \mathcal{C}$: tensor autoequivalences of \mathcal{C} and monoidal natural transformations.

Equivariantization

Let \mathcal{C} be a tensor category over k , G a group.

$\underline{\text{Aut}}_{\otimes} \mathcal{C}$: tensor autoequivalences of \mathcal{C} and monoidal natural transformations.

\underline{G} : G and identities as only morphisms.

Equivariantization

Let \mathcal{C} be a tensor category over k , G a group.

$\underline{\text{Aut}}_{\otimes} \mathcal{C}$: tensor autoequivalences of \mathcal{C} and monoidal natural transformations.

\underline{G} : G and identities as only morphisms.

An *action* of G on \mathcal{C} is a strong monoidal functor

$$\rho : \underline{G} \rightarrow \underline{\text{Aut}}_{\otimes} \mathcal{C}.$$

Equivariantization

Let \mathcal{C} be a tensor category over k , G a group.

$\underline{\text{Aut}}_{\otimes} \mathcal{C}$: tensor autoequivalences of \mathcal{C} and monoidal natural transformations.

\underline{G} : G and identities as only morphisms.

An *action* of G on \mathcal{C} is a strong monoidal functor

$$\rho : \underline{G} \rightarrow \underline{\text{Aut}}_{\otimes} \mathcal{C}.$$

A G -equivariant object is a pair (X, u) , where:

Equivariantization

Let \mathcal{C} be a tensor category over k , G a group.

$\underline{\text{Aut}}_{\otimes} \mathcal{C}$: tensor autoequivalences of \mathcal{C} and monoidal natural transformations.

\underline{G} : G and identities as only morphisms.

An *action* of G on \mathcal{C} is a strong monoidal functor

$$\rho : \underline{G} \rightarrow \underline{\text{Aut}}_{\otimes} \mathcal{C}.$$

A G -equivariant object is a pair (X, u) , where:

- ▶ X is an object of \mathcal{C} , and
- ▶ $u = (u^g)_{g \in G}$, $u^g : \rho^g X \rightarrow X$ is an isomorphism st:

$$u^g \rho^g(u^h) = u^{gh} \rho_{2_X}^{g,h}, \quad \forall g, h \in G,$$

$$u_1 \rho_0 X = \text{id}_X.$$

Equivariantization

Let \mathcal{C} be a tensor category over k , G a group.

$\underline{\text{Aut}}_{\otimes} \mathcal{C}$: tensor autoequivalences of \mathcal{C} and monoidal natural transformations.

\underline{G} : G and identities as only morphisms.

An *action* of G on \mathcal{C} is a strong monoidal functor

$$\rho : \underline{G} \rightarrow \underline{\text{Aut}}_{\otimes} \mathcal{C}.$$

A *G -equivariant object* is a pair (X, u) , where:

- ▶ X is an object of \mathcal{C} , and
- ▶ $u = (u^g)_{g \in G}$, $u^g : \rho^g X \rightarrow X$ is an isomorphism st:

$$u^g \rho^g(u^h) = u^{gh} \rho_{2_X}^{g,h}, \quad \forall g, h \in G,$$

$$u_1 \rho_0 X = \text{id}_X.$$

A *G -equivariant morphism* $f : (X, u) \rightarrow (Y, v)$ is a morphism $f : X \rightarrow Y$ st: $fu_g = v_g f, \forall g \in G$.

\mathcal{C}^G : the G -equivariantization of \mathcal{C} .

\mathcal{C}^G : the G -equivariantization of \mathcal{C} .

G -equivariant objects and G -equivariant morphisms.

\mathcal{C}^G : the G -equivariantization of \mathcal{C} .

G -equivariant objects and G -equivariant morphisms.

\mathcal{C}^G is a tensor category and the forgetful functor

$$\mathcal{U} : \mathcal{C}^G \rightarrow \mathcal{C}$$

is a normal dominant tensor functor.

\mathcal{C}^G : the G -equivariantization of \mathcal{C} .

G -equivariant objects and G -equivariant morphisms.

\mathcal{C}^G is a tensor category and the forgetful functor

$$\mathcal{U} : \mathcal{C}^G \rightarrow \mathcal{C}$$

is a normal dominant tensor functor.

Example

Let G be a group and let ρ the trivial action of G on vec_k .
Then

\mathcal{C}^G : the G -equivariantization of \mathcal{C} .

G -equivariant objects and G -equivariant morphisms.

\mathcal{C}^G is a tensor category and the forgetful functor

$$\mathcal{U} : \mathcal{C}^G \rightarrow \mathcal{C}$$

is a normal dominant tensor functor.

Example

Let G be a group and let ρ the trivial action of G on vec_k .
Then

$$(\text{vec}_k)^G = \text{rep } G.$$

\mathcal{C}^G : the G -equivariantization of \mathcal{C} .

G -equivariant objects and G -equivariant morphisms.

\mathcal{C}^G is a tensor category and the forgetful functor

$$\mathcal{U} : \mathcal{C}^G \rightarrow \mathcal{C}$$

is a normal dominant tensor functor.

Example

Let G be a group and let ρ the trivial action of G on vec_k .
Then

$$(\text{vec}_k)^G = \text{rep } G.$$

Equivariantization gives rise canonically to an exact sequence

$$\text{rep } G \rightarrow \mathcal{C}^G \rightarrow \mathcal{C}.$$

Some properties of exact sequences

Proposition

Let $\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$ be an exact sequence of tensor categories, with induced Hopf algebra H .

Some properties of exact sequences

Proposition

Let $\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$ be an exact sequence of tensor categories, with induced Hopf algebra H .

The following are equivalent:

Some properties of exact sequences

Proposition

Let $\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$ be an exact sequence of tensor categories, with induced Hopf algebra H .

The following are equivalent:

- ▶ *The functor F has adjoints;*

Some properties of exact sequences

Proposition

Let $\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$ be an exact sequence of tensor categories, with induced Hopf algebra H .

The following are equivalent:

- ▶ *The functor F has adjoints;*
- ▶ *The tensor category \mathcal{C}' is finite;*

Some properties of exact sequences

Proposition

Let $\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$ be an exact sequence of tensor categories, with induced Hopf algebra H .

The following are equivalent:

- ▶ *The functor F has adjoints;*
- ▶ *The tensor category \mathcal{C}' is finite;*
- ▶ *The Hopf algebra H is finite-dimensional.*

Some properties of exact sequences

Proposition

Let $\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$ be an exact sequence of tensor categories, with induced Hopf algebra H .

The following are equivalent:

- ▶ *The functor F has adjoints;*
- ▶ *The tensor category \mathcal{C}' is finite;*
- ▶ *The Hopf algebra H is finite-dimensional.*

In particular, if \mathcal{C}' and \mathcal{C}'' are finite, so is \mathcal{C} .

Some properties of exact sequences

Proposition

Let $\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$ be an exact sequence of tensor categories, with induced Hopf algebra H .

The following are equivalent:

- ▶ *The functor F has adjoints;*
- ▶ *The tensor category \mathcal{C}' is finite;*
- ▶ *The Hopf algebra H is finite-dimensional.*

In particular, if \mathcal{C}' and \mathcal{C}'' are finite, so is \mathcal{C} .

A fusion category \mathcal{C} is integral if $\text{FPdim } X \in \mathbb{Z}, \forall X \in \text{Irr}(\mathcal{C})$.

Some properties of exact sequences

Proposition

Let $\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}'$ be an exact sequence of tensor categories, with induced Hopf algebra H .

The following are equivalent:

- ▶ *The functor F has adjoints;*
- ▶ *The tensor category \mathcal{C}' is finite;*
- ▶ *The Hopf algebra H is finite-dimensional.*

In particular, if \mathcal{C}' and \mathcal{C}'' are finite, so is \mathcal{C} .

A fusion category \mathcal{C} is integral if $\text{FPdim } X \in \mathbb{Z}, \forall X \in \text{Irr}(\mathcal{C})$.

\mathcal{C} is weakly integral if $\text{FPdim } \mathcal{C} = \sum_{X \in \text{Irr}(\mathcal{C})} (\text{FPdim } X)^2 \in \mathbb{Z}$.

Some properties of exact sequences

Proposition

Let $\mathcal{C}' \xrightarrow{f} \mathcal{C} \xrightarrow{F} \mathcal{C}''$ be an exact sequence of tensor categories, with induced Hopf algebra H .

The following are equivalent:

- ▶ The functor F has adjoints;
- ▶ The tensor category \mathcal{C}' is finite;
- ▶ The Hopf algebra H is finite-dimensional.

In particular, if \mathcal{C}' and \mathcal{C}'' are finite, so is \mathcal{C} .

A fusion category \mathcal{C} is *integral* if $\text{FPdim } X \in \mathbb{Z}, \forall X \in \text{Irr}(\mathcal{C})$.

\mathcal{C} is *weakly integral* if $\text{FPdim } \mathcal{C} = \sum_{X \in \text{Irr}(\mathcal{C})} (\text{FPdim } X)^2 \in \mathbb{Z}$.

Proposition

Let $\mathcal{C}' \rightarrow \mathcal{C} \rightarrow \mathcal{C}''$ be an exact sequence of fusion categories. Then \mathcal{C} is (weakly) integral iff \mathcal{C}'' is (weakly) integral.

Proposition

Let $\mathcal{C}, \mathcal{C}', \mathcal{C}''$ be fusion categories, $i : \mathcal{C}' \rightarrow \mathcal{C}$, $F : \mathcal{C} \rightarrow \mathcal{C}''$, tensor functors.

Proposition

Let $\mathcal{C}, \mathcal{C}', \mathcal{C}''$ be fusion categories, $i : \mathcal{C}' \rightarrow \mathcal{C}$, $F : \mathcal{C} \rightarrow \mathcal{C}''$, tensor functors. Assume that F is dominant, i is full and $i(\mathcal{C}) \subset \mathcal{R}\text{ef}_F$.

Proposition

Let $\mathcal{C}, \mathcal{C}', \mathcal{C}''$ be fusion categories, $i : \mathcal{C}' \rightarrow \mathcal{C}$, $F : \mathcal{C} \rightarrow \mathcal{C}''$, tensor functors. Assume that F is dominant, i is full and $i(\mathcal{C}) \subset \mathcal{R}\text{ef}_F$.

Then $\text{FPdim } \mathcal{C} \geq \text{FPdim } \mathcal{C}' \text{ FPdim } \mathcal{C}''$.

Proposition

Let $\mathcal{C}, \mathcal{C}', \mathcal{C}''$ be fusion categories, $i : \mathcal{C}' \rightarrow \mathcal{C}$, $F : \mathcal{C} \rightarrow \mathcal{C}''$, tensor functors. Assume that F is dominant, i is full and $i(\mathcal{C}) \subset \mathcal{R}\text{ef}_F$.

Then $\text{FPdim } \mathcal{C} \geq \text{FPdim } \mathcal{C}' \text{ FPdim } \mathcal{C}''$.

Moreover, the sequence

$$\mathcal{C}' \xrightarrow{i} \mathcal{C} \xrightarrow{F} \mathcal{C}''$$

is exact iff $\text{FPdim } \mathcal{C} = \text{FPdim } \mathcal{C}' \text{ FPdim } \mathcal{C}''$.

Proposition

Let $\mathcal{C}, \mathcal{C}', \mathcal{C}''$ be fusion categories, $i : \mathcal{C}' \rightarrow \mathcal{C}$, $F : \mathcal{C} \rightarrow \mathcal{C}''$, tensor functors. Assume that F is dominant, i is full and $i(\mathcal{C}) \subset \text{Ret}_F$.

Then $\text{FPdim } \mathcal{C} \geq \text{FPdim } \mathcal{C}' \text{ FPdim } \mathcal{C}''$.

Moreover, the sequence

$$\mathcal{C}' \xrightarrow{i} \mathcal{C} \xrightarrow{F} \mathcal{C}''$$

is exact iff $\text{FPdim } \mathcal{C} = \text{FPdim } \mathcal{C}' \text{ FPdim } \mathcal{C}''$.

If so, then \forall simple object $Y \in \mathcal{C}''$,

$$\text{FPdim } Y = \frac{1}{\text{FPdim } \mathcal{C}'} \sum_{X \in \text{Irr}(\mathcal{C})} m_Y(F(X)) \text{FPdim } X,$$

where $m_Y(F(X)) = \dim \text{Hom}_{\mathcal{C}''}(Y, F(X))$.

Hopf monads and their modules

Let \mathcal{C} be a monoidal category. A monad T on \mathcal{C} is an algebra $T \in \text{End}(\mathcal{C})$:

$$\mu : T^2 \rightarrow T, \quad \eta : \text{id}_{\mathcal{C}} \rightarrow T.$$

Hopf monads and their modules

Let \mathcal{C} be a monoidal category. A monad T on \mathcal{C} is an algebra $T \in \text{End}(\mathcal{C})$:

$$\mu : T^2 \rightarrow T, \quad \eta : \text{id}_{\mathcal{C}} \rightarrow T.$$

\mathcal{C}^T : T -modules in \mathcal{C} . Objects: (X, r) , $X \in \mathcal{C}$, $r : T(X) \rightarrow X$, st:

$$rT(r) = r\mu_X, \quad r\eta_X = \text{id}_X.$$

Hopf monads and their modules

Let \mathcal{C} be a monoidal category. A monad T on \mathcal{C} is an algebra $T \in \text{End}(\mathcal{C})$:

$$\mu : T^2 \rightarrow T, \quad \eta : \text{id}_{\mathcal{C}} \rightarrow T.$$

\mathcal{C}^T : T -modules in \mathcal{C} . Objects: (X, r) , $X \in \mathcal{C}$, $r : T(X) \rightarrow X$, st:

$$rT(r) = r\mu_X, \quad r\eta_X = \text{id}_X.$$

Consider the forgetful functor $\mathcal{U} : \mathcal{C}^T \rightarrow \mathcal{C}$.

Hopf monads and their modules

Let \mathcal{C} be a monoidal category. A monad T on \mathcal{C} is an algebra $T \in \text{End}(\mathcal{C})$:

$$\mu : T^2 \rightarrow T, \quad \eta : \text{id}_{\mathcal{C}} \rightarrow T.$$

\mathcal{C}^T : T -modules in \mathcal{C} . Objects: (X, r) , $X \in \mathcal{C}$, $r : T(X) \rightarrow X$, st:

$$rT(r) = r\mu_X, \quad r\eta_X = \text{id}_X.$$

Consider the forgetful functor $\mathcal{U} : \mathcal{C}^T \rightarrow \mathcal{C}$.

A monad T is a *bimonad* iff \mathcal{C}^T is a monoidal category st \mathcal{U} is strict monoidal.

Hopf monads and their modules

Let \mathcal{C} be a monoidal category. A monad T on \mathcal{C} is an algebra $T \in \text{End}(\mathcal{C})$:

$$\mu : T^2 \rightarrow T, \quad \eta : \text{id}_{\mathcal{C}} \rightarrow T.$$

\mathcal{C}^T : T -modules in \mathcal{C} . Objects: (X, r) , $X \in \mathcal{C}$, $r : T(X) \rightarrow X$, st:

$$rT(r) = r\mu_X, \quad r\eta_X = \text{id}_X.$$

Consider the forgetful functor $\mathcal{U} : \mathcal{C}^T \rightarrow \mathcal{C}$.

A monad T is a *bimonad* iff \mathcal{C}^T is a monoidal category st \mathcal{U} is strict monoidal.

Equivalently, T is a *comonoidal endofunctor*:

Hopf monads and their modules

Let \mathcal{C} be a monoidal category. A monad T on \mathcal{C} is an algebra $T \in \text{End}(\mathcal{C})$:

$$\mu : T^2 \rightarrow T, \quad \eta : \text{id}_{\mathcal{C}} \rightarrow T.$$

\mathcal{C}^T : T -modules in \mathcal{C} . Objects: (X, r) , $X \in \mathcal{C}$, $r : T(X) \rightarrow X$, st:

$$rT(r) = r\mu_X, \quad r\eta_X = \text{id}_X.$$

Consider the forgetful functor $\mathcal{U} : \mathcal{C}^T \rightarrow \mathcal{C}$.

A monad T is a *bimonad* iff \mathcal{C}^T is a monoidal category st \mathcal{U} is strict monoidal.

Equivalently, T is a *comonoidal endofunctor*:

$$T_2(X, Y) : T(X \otimes Y) \rightarrow T(X) \otimes T(Y), \quad T_0 : T(\mathbb{K}) \rightarrow \mathbb{K},$$

$\forall X, Y \in \mathcal{C}$, satisfying:

$$(T_2(X, Y) \otimes \text{id}_{T(Z)}) T_2(X \otimes Y, Z) = (\text{id}_{T(X)} \otimes T_2(Y, Z)) T_2(X, Y \otimes Z),$$

$$(\text{id}_{T(X)} \otimes T_0) T_2(X, \mathbb{K}) = \text{id}_{T(X)} = (T_0 \otimes \text{id}_{T(X)}) T_2(\mathbb{K}, X),$$

and st μ, η are monoidal transformations, that is:

$$T_2(X, Y) \mu_{X \otimes Y} = (\mu_X \otimes \mu_Y) T_2(T(X), T(Y)) T(T_2(X, Y)),$$

$$T_0 \mu_{\mathbb{K}} = T_0 T(T_0), \quad T_2(X, Y) \eta_{X \otimes Y} = \eta_X \otimes \eta_Y, \quad T_0 \eta_{\mathbb{K}} = \text{id}_{\mathbb{K}}.$$

$$(T_2(X, Y) \otimes \text{id}_{T(Z)}) T_2(X \otimes Y, Z) = (\text{id}_{T(X)} \otimes T_2(Y, Z)) T_2(X, Y \otimes Z),$$

$$(\text{id}_{T(X)} \otimes T_0) T_2(X, \mathbb{K}) = \text{id}_{T(X)} = (T_0 \otimes \text{id}_{T(X)}) T_2(\mathbb{K}, X),$$

and st μ, η are monoidal transformations, that is:

$$T_2(X, Y) \mu_{X \otimes Y} = (\mu_X \otimes \mu_Y) T_2(T(X), T(Y)) T(T_2(X, Y)),$$

$$T_0 \mu_{\mathbb{K}} = T_0 T(T_0), \quad T_2(X, Y) \eta_{X \otimes Y} = \eta_X \otimes \eta_Y, \quad T_0 \eta_{\mathbb{K}} = \text{id}_{\mathbb{K}}.$$

Suppose \mathcal{C} is rigid.

$$(T_2(X, Y) \otimes \text{id}_{T(Z)}) T_2(X \otimes Y, Z) = (\text{id}_{T(X)} \otimes T_2(Y, Z)) T_2(X, Y \otimes Z),$$

$$(\text{id}_{T(X)} \otimes T_0) T_2(X, \mathbb{K}) = \text{id}_{T(X)} = (T_0 \otimes \text{id}_{T(X)}) T_2(\mathbb{K}, X),$$

and st μ, η are monoidal transformations, that is:

$$T_2(X, Y) \mu_{X \otimes Y} = (\mu_X \otimes \mu_Y) T_2(T(X), T(Y)) T(T_2(X, Y)),$$

$$T_0 \mu_{\mathbb{K}} = T_0 T(T_0), \quad T_2(X, Y) \eta_{X \otimes Y} = \eta_X \otimes \eta_Y, \quad T_0 \eta_{\mathbb{K}} = \text{id}_{\mathbb{K}}.$$

Suppose \mathcal{C} is rigid.

A bimonad T on \mathcal{C} is a *Hopf monad* if \mathcal{C}^T is rigid.

Example

Example

Let $\mathcal{C} \simeq \text{vec}_k$. If H is a finite dimensional Hopf algebra over k , then $H \otimes ? : \mathcal{C} \rightarrow \mathcal{C}$ is a k -linear Hopf monad on \mathcal{C} .

Example

Let $\mathcal{C} \simeq \text{vec}_k$. If H is a finite dimensional Hopf algebra over k , then $H \otimes ? : \mathcal{C} \rightarrow \mathcal{C}$ is a k -linear Hopf monad on \mathcal{C} .

$H \mapsto H \otimes ?$ defines an equivalence of categories between:

Example

Let $\mathcal{C} \simeq \text{vec}_k$. If H is a finite dimensional Hopf algebra over k , then $H \otimes ? : \mathcal{C} \rightarrow \mathcal{C}$ is a k -linear Hopf monad on \mathcal{C} .

$H \mapsto H \otimes ?$ defines an equivalence of categories between:

- Finite dimensional Hopf algebras over k , and

Example

Let $\mathcal{C} \simeq \text{vec}_k$. If H is a finite dimensional Hopf algebra over k , then $H \otimes ? : \mathcal{C} \rightarrow \mathcal{C}$ is a k -linear Hopf monad on \mathcal{C} .

$H \mapsto H \otimes ?$ defines an equivalence of categories between:

- ▶ Finite dimensional Hopf algebras over k , and
- ▶ k -linear Hopf monads on \mathcal{C} .

Example

Let $\mathcal{C} \simeq \text{vec}_k$. If H is a finite dimensional Hopf algebra over k , then $H \otimes ? : \mathcal{C} \rightarrow \mathcal{C}$ is a k -linear Hopf monad on \mathcal{C} .

$H \mapsto H \otimes ?$ defines an equivalence of categories between:

- ▶ Finite dimensional Hopf algebras over k , and
- ▶ k -linear Hopf monads on \mathcal{C} .

Proposition

Let \mathcal{C} be a tensor category over k , T a k -linear right exact Hopf monad on \mathcal{C} . Then:

Example

Let $\mathcal{C} \simeq \text{vec}_k$. If H is a finite dimensional Hopf algebra over k , then $H \otimes ? : \mathcal{C} \rightarrow \mathcal{C}$ is a k -linear Hopf monad on \mathcal{C} .

$H \mapsto H \otimes ?$ defines an equivalence of categories between:

- ▶ Finite dimensional Hopf algebras over k , and
- ▶ k -linear Hopf monads on \mathcal{C} .

Proposition

Let \mathcal{C} be a tensor category over k , T a k -linear right exact Hopf monad on \mathcal{C} . Then:

\mathcal{C}^T is a tensor category over k and the forgetful functor $\mathcal{U} : \mathcal{C}^T \rightarrow \mathcal{C}$ is a tensor functor.

Monadicity

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a tensor functor st F admits a left adjoint G .

Monadicity

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a tensor functor st F admits a left adjoint G .

Then the adjunction $G \vdash F$ is *monadic*: $T = FG$ is a Hopf monad on \mathcal{D} and we have

$$\mathcal{C} \simeq \mathcal{D}^T,$$

as tensor categories.

Monadicity

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a tensor functor st F admits a left adjoint G .

Then the adjunction $G \vdash F$ is *monadic*: $T = FG$ is a Hopf monad on \mathcal{D} and we have

$$\mathcal{C} \simeq \mathcal{D}^T,$$

as tensor categories.

If \mathcal{C}, \mathcal{D} are *finite* tensor categories $\Rightarrow F$ admits a left adjoint.

Monadicity

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a tensor functor st F admits a left adjoint G .

Then the adjunction $G \vdash F$ is *monadic*: $T = FG$ is a Hopf monad on \mathcal{D} and we have

$$\mathcal{C} \simeq \mathcal{D}^T,$$

as tensor categories.

If \mathcal{C}, \mathcal{D} are *finite* tensor categories $\Rightarrow F$ admits a left adjoint.

Therefore in this case F is monadic.

Main theorem

Definition

Let \mathcal{C} be a tensor category.

A k -linear right exact Hopf monad T on \mathcal{C} is *normal* if $T(\mathbb{1})$ is a trivial object of \mathcal{C} .

Main theorem

Definition

Let \mathcal{C} be a tensor category.

A k -linear right exact Hopf monad T on \mathcal{C} is *normal* if $T(\mathbb{1})$ is a trivial object of \mathcal{C} .

The following theorem classifies extensions of tensor categories in terms of Hopf monads.

Main theorem

Definition

Let \mathcal{C} be a tensor category.

A k -linear right exact Hopf monad T on \mathcal{C} is *normal* if $T(\mathbb{1})$ is a trivial object of \mathcal{C} .

The following theorem classifies extensions of tensor categories in terms of Hopf monads.

Theorem

Let \mathcal{C}' , \mathcal{C}'' be tensor categories over a field k st \mathcal{C}' is finite.

Main theorem

Definition

Let \mathcal{C} be a tensor category.

A k -linear right exact Hopf monad T on \mathcal{C} is *normal* if $T(\mathbb{1})$ is a trivial object of \mathcal{C} .

The following theorem classifies extensions of tensor categories in terms of Hopf monads.

Theorem

Let $\mathcal{C}', \mathcal{C}''$ be tensor categories over a field k st \mathcal{C}' is finite.

The following data are equivalent:

Main theorem

Definition

Let \mathcal{C} be a tensor category.

A k -linear right exact Hopf monad T on \mathcal{C} is *normal* if $T(\mathbb{1})$ is a trivial object of \mathcal{C} .

The following theorem classifies extensions of tensor categories in terms of Hopf monads.

Theorem

Let $\mathcal{C}', \mathcal{C}''$ be tensor categories over a field k st \mathcal{C}' is finite.

The following data are equivalent:

- ▶ A normal faithful k -linear right exact Hopf monad T on \mathcal{C}'' , with $T|_{\langle \mathbb{1} \rangle} = H$, st $\mathcal{C}' \simeq \text{comod-}H$;

Main theorem

Definition

Let \mathcal{C} be a tensor category.

A k -linear right exact Hopf monad T on \mathcal{C} is *normal* if $T(\mathbb{1})$ is a trivial object of \mathcal{C} .

The following theorem classifies extensions of tensor categories in terms of Hopf monads.

Theorem

Let $\mathcal{C}', \mathcal{C}''$ be tensor categories over a field k st \mathcal{C}' is finite.

The following data are equivalent:

- ▶ A normal faithful k -linear right exact Hopf monad T on \mathcal{C}'' , with $T|_{\langle \mathbb{1} \rangle} = H$, st $\mathcal{C}' \simeq \text{comod-}H$;
- ▶ An extension $\mathcal{C}' \rightarrow \mathcal{C} \rightarrow \mathcal{C}''$ of \mathcal{C}'' by \mathcal{C}' with induced Hopf algebra H .

Equivariantization revisited

Let \mathcal{C} be a tensor category over k .

Equivariantization revisited

Let \mathcal{C} be a tensor category over k .

Let $\rho : \underline{G} \rightarrow \underline{\text{Aut}}_{\otimes} \mathcal{C}$ be an action of a finite group G on \mathcal{C} . Then:

Equivariantization revisited

Let \mathcal{C} be a tensor category over k .

Let $\rho : \underline{G} \rightarrow \underline{\text{Aut}}_{\otimes} \mathcal{C}$ be an action of a finite group G on \mathcal{C} . Then:

- The k -linear exact endofunctor

$$T^{\rho} = \bigoplus_{g \in G} \rho^g$$

is canonically a Hopf monad on \mathcal{C} ;

Equivariantization revisited

Let \mathcal{C} be a tensor category over k .

Let $\rho : \underline{G} \rightarrow \underline{\text{Aut}}_{\otimes} \mathcal{C}$ be an action of a finite group G on \mathcal{C} . Then:

- ▶ The k -linear exact endofunctor

$$T^{\rho} = \bigoplus_{g \in G} \rho^g$$

is canonically a Hopf monad on \mathcal{C} ;

- ▶ There is a canonical isomorphism:

$$\mathcal{C}^G \simeq \mathcal{C}^{T^{\rho}}.$$

Equivariantization revisited

Let \mathcal{C} be a tensor category over k .

Let $\rho : \underline{G} \rightarrow \underline{\text{Aut}}_{\otimes} \mathcal{C}$ be an action of a finite group G on \mathcal{C} . Then:

- ▶ The k -linear exact endofunctor

$$T^{\rho} = \bigoplus_{g \in G} \rho^g$$

is canonically a Hopf monad on \mathcal{C} ;

- ▶ There is a canonical isomorphism:

$$\mathcal{C}^G \simeq \mathcal{C}^{T^{\rho}}.$$

- ▶ The Hopf monad T^{ρ} is faithful, normal, and *cocommutative*.

Equivariantization revisited

Let \mathcal{C} be a tensor category over k .

Let $\rho : \underline{G} \rightarrow \underline{\text{Aut}}_{\otimes} \mathcal{C}$ be an action of a finite group G on \mathcal{C} . Then:

- ▶ The k -linear exact endofunctor

$$T^\rho = \bigoplus_{g \in G} \rho^g$$

is canonically a Hopf monad on \mathcal{C} ;

- ▶ There is a canonical isomorphism:

$$\mathcal{C}^G \simeq \mathcal{C}^{T^\rho}.$$

- ▶ The Hopf monad T^ρ is faithful, normal, and *cocommutative*.
- ▶ The induced Hopf algebra of T is k^G .

Hopf monad of a group extension

Let $1 \rightarrow G'' \xrightarrow{\iota} G \xrightarrow{\pi} G' \rightarrow 1$ be an exact sequence of finite groups.

Hopf monad of a group extension

Let $1 \rightarrow G'' \xrightarrow{\iota} G \xrightarrow{\pi} G' \rightarrow 1$ be an exact sequence of finite groups.

Then we have an exact sequence:

$$\operatorname{rep} G' \xrightarrow{\pi^*} \operatorname{rep} G \xrightarrow{\iota^*} \operatorname{rep} G''.$$

Hopf monad of a group extension

Let $1 \rightarrow G'' \xrightarrow{\iota} G \xrightarrow{\pi} G' \rightarrow 1$ be an exact sequence of finite groups.

Then we have an exact sequence:

$$\mathrm{rep} G' \xrightarrow{\pi^*} \mathrm{rep} G \xrightarrow{\iota^*} \mathrm{rep} G''.$$

Identify: $G'' \trianglelefteq G$, $G' = G/G''$.

Hopf monad of a group extension

Let $1 \rightarrow G'' \xrightarrow{\iota} G \xrightarrow{\pi} G' \rightarrow 1$ be an exact sequence of finite groups.

Then we have an exact sequence:

$$\text{rep } G' \xrightarrow{\pi^*} \text{rep } G \xrightarrow{\iota^*} \text{rep } G''.$$

Identify: $G'' \trianglelefteq G$, $G' = G/G''$.

Then $i^* = \text{Res}_{G''}^G$ (restriction functor).

Hopf monad of a group extension

Let $1 \rightarrow G'' \xrightarrow{\iota} G \xrightarrow{\pi} G' \rightarrow 1$ be an exact sequence of finite groups.

Then we have an exact sequence:

$$\text{rep } G' \xrightarrow{\pi^*} \text{rep } G \xrightarrow{\iota^*} \text{rep } G''.$$

Identify: $G'' \trianglelefteq G$, $G' = G/G''$.

Then $i^* = \text{Res}_{G''}^G$ (restriction functor).

The induction functor $\text{Ind}_{G''}^G : \text{rep } G'' \rightarrow \text{rep } G$ is left adjoint to $\text{Res}_{G''}^G$.

Hopf monad of a group extension

Let $1 \rightarrow G'' \xrightarrow{\iota} G \xrightarrow{\pi} G' \rightarrow 1$ be an exact sequence of finite groups.

Then we have an exact sequence:

$$\operatorname{rep} G' \xrightarrow{\pi^*} \operatorname{rep} G \xrightarrow{\iota^*} \operatorname{rep} G''.$$

Identify: $G'' \trianglelefteq G$, $G' = G/G''$.

Then $i^* = \operatorname{Res}_{G''}^G$ (restriction functor).

The induction functor $\operatorname{Ind}_{G''}^G : \operatorname{rep} G'' \rightarrow \operatorname{rep} G$ is left adjoint to $\operatorname{Res}_{G''}^G$.

Let Y be a kG'' -module. Then:

Hopf monad of a group extension

Let $1 \rightarrow G'' \xrightarrow{\iota} G \xrightarrow{\pi} G' \rightarrow 1$ be an exact sequence of finite groups.

Then we have an exact sequence:

$$\operatorname{rep} G' \xrightarrow{\pi^*} \operatorname{rep} G \xrightarrow{\iota^*} \operatorname{rep} G''.$$

Identify: $G'' \trianglelefteq G$, $G' = G/G''$.

Then $i^* = \operatorname{Res}_{G''}^G$ (restriction functor).

The induction functor $\operatorname{Ind}_{G''}^G : \operatorname{rep} G'' \rightarrow \operatorname{rep} G$ is left adjoint to $\operatorname{Res}_{G''}^G$.

Let Y be a kG'' -module. Then:

$$T(Y) = \operatorname{Res}_{G''}^G \operatorname{Ind}_{G''}^G(Y) \simeq \bigoplus_{\gamma \in G'} {}^\gamma Y,$$

Hopf monad of a group extension

Let $1 \rightarrow G'' \xrightarrow{\iota} G \xrightarrow{\pi} G' \rightarrow 1$ be an exact sequence of finite groups.

Then we have an exact sequence:

$$\text{rep } G' \xrightarrow{\pi^*} \text{rep } G \xrightarrow{\iota^*} \text{rep } G''.$$

Identify: $G'' \trianglelefteq G$, $G' = G/G''$.

Then $i^* = \text{Res}_{G''}^G$ (restriction functor).

The induction functor $\text{Ind}_{G''}^G : \text{rep } G'' \rightarrow \text{rep } G$ is left adjoint to $\text{Res}_{G''}^G$.

Let Y be a kG'' -module. Then:

$$T(Y) = \text{Res}_{G''}^G \text{Ind}_{G''}^G(Y) \simeq \bigoplus_{\gamma \in G'} {}^\gamma Y,$$

where ${}^\gamma Y$ denotes the kG'' -module conjugated to Y . (Mackey's Subgroup Theorem.)

Hopf monad of a group extension

Let $1 \rightarrow G'' \xrightarrow{\iota} G \xrightarrow{\pi} G' \rightarrow 1$ be an exact sequence of finite groups.

Then we have an exact sequence:

$$\operatorname{rep} G' \xrightarrow{\pi^*} \operatorname{rep} G \xrightarrow{\iota^*} \operatorname{rep} G''.$$

Identify: $G'' \trianglelefteq G$, $G' = G/G''$.

Then $i^* = \operatorname{Res}_{G''}^G$ (restriction functor).

The induction functor $\operatorname{Ind}_{G''}^G : \operatorname{rep} G'' \rightarrow \operatorname{rep} G$ is left adjoint to $\operatorname{Res}_{G''}^G$.

Let Y be a kG'' -module. Then:

$$T(Y) = \operatorname{Res}_{G''}^G \operatorname{Ind}_{G''}^G(Y) \simeq \bigoplus_{\gamma \in G'} {}^\gamma Y,$$

where ${}^\gamma Y$ denotes the kG'' -module conjugated to Y . (Mackey's Subgroup Theorem.)

In fact, conjugation under $\gamma \in G' = G/G''$ defines an action of G' on $\operatorname{rep} G''$ by tensor autoequivalences.

Tensor functors of small index

The *Frobenius-Perron index* of a dominant tensor functor $F : \mathcal{C} \rightarrow \mathcal{D}$ between fusion categories is:

Tensor functors of small index

The *Frobenius-Perron index* of a dominant tensor functor $F : \mathcal{C} \rightarrow \mathcal{D}$ between fusion categories is:

$$\mathrm{FPind}(F) = \mathrm{FPind}(\mathcal{C} : \mathcal{D}) = \frac{\mathrm{FPdim} \mathcal{C}}{\mathrm{FPdim} \mathcal{D}}.$$

Tensor functors of small index

The *Frobenius-Perron index* of a dominant tensor functor $F : \mathcal{C} \rightarrow \mathcal{D}$ between fusion categories is:

$$\mathrm{FPind}(F) = \mathrm{FPind}(\mathcal{C} : \mathcal{D}) = \frac{\mathrm{FPdim} \mathcal{C}}{\mathrm{FPdim} \mathcal{D}}.$$

$\mathrm{FPind}(F)$ is an algebraic integer.

Tensor functors of small index

The *Frobenius-Perron index* of a dominant tensor functor $F : \mathcal{C} \rightarrow \mathcal{D}$ between fusion categories is:

$$\mathrm{FPind}(F) = \mathrm{FPind}(\mathcal{C} : \mathcal{D}) = \frac{\mathrm{FPdim} \mathcal{C}}{\mathrm{FPdim} \mathcal{D}}.$$

$\mathrm{FPind}(F)$ is an algebraic integer.

Theorem

Suppose $\mathrm{char} k = 0$. Let \mathcal{C}, \mathcal{D} be fusion categories over k and let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a dominant tensor functor. If $\mathrm{FPind}(\mathcal{C} : \mathcal{D}) = 2$, then F is an equivariantization.

Tensor functors of small index

The *Frobenius-Perron index* of a dominant tensor functor $F : \mathcal{C} \rightarrow \mathcal{D}$ between fusion categories is:

$$\text{FPind}(F) = \text{FPind}(\mathcal{C} : \mathcal{D}) = \frac{\text{FPdim } \mathcal{C}}{\text{FPdim } \mathcal{D}}.$$

$\text{FPind}(F)$ is an algebraic integer.

Theorem

Suppose $\text{char } k = 0$. Let \mathcal{C}, \mathcal{D} be fusion categories over k and let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a dominant tensor functor. If $\text{FPind}(\mathcal{C} : \mathcal{D}) = 2$, then F is an equivariantization.

Theorem

Suppose $\text{char } k = 0$. Let \mathcal{C}, \mathcal{D} be fusion categories over k st \mathcal{C} is weakly integral. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a dominant tensor functor. If $\text{FPind}(\mathcal{C} : \mathcal{D}) = p$, where p is the smallest prime divisor of $\text{FPdim } \mathcal{C}$, then F is an equivariantization.

References



A. Bruguières, S. Natale.

Exact sequences of tensor categories.

Int. Math. Res. Not. **2011** (24) 5644-5705, 2011.



A. Bruguières, S. Natale.

Central exact sequences of tensor categories,
equivariantizations and applications.

Preprint [arXiv:1112.3135](https://arxiv.org/abs/1112.3135), 2011.



A. Bruguières, S. Lack, A. Virelizier.

Hopf monads on monoidal categories.

Adv. Math. **227** 745-800, 2011.