

On Tensor Products with the Steinberg Character of the Unitary Group

Rod Gow

School of Mathematical Sciences/CSI
University College Dublin, Ireland
rod.gow@ucd.ie

Regina, June 4, 2012

Outline

- 1 The Weil Representation of the Unitary Group
- 2 Reality Properties of the Irreducible Constituents of the Weil Character
- 3 Tensoring with the Steinberg Character
- 4 Some Consequences

The Unitary Group

For the purposes of this talk, we will restrict our attention to the unitary group, $U(n, q)$.

Note here that we consider what is sometimes called the *general* unitary group.

Let $q = p^c$, where p is a prime and c is a positive integer.

The Unitary Group

For the purposes of this talk, we will restrict our attention to the unitary group, $U(n, q)$.

Note here that we consider what is sometimes called the *general* unitary group.

Let $q = p^c$, where p is a prime and c is a positive integer.

Let \mathbb{F}_{q^2} denote the finite field of order q^2 .

The Unitary Group

For the purposes of this talk, we will restrict our attention to the unitary group, $U(n, q)$.

Note here that we consider what is sometimes called the *general* unitary group.

Let $q = p^c$, where p is a prime and c is a positive integer.

Let \mathbb{F}_{q^2} denote the finite field of order q^2 .

The unitary group $U(n, q)$ is the group of isometries of a non-degenerate hermitian form defined on $V \times V$, where V is a vector space of dimension n over \mathbb{F}_{q^2} .

Properties of the Unitary Group

The order of the unitary group is

$$|U(n, q)| = q^{n(n-1)/2} \prod_{i=1}^n (q^i - (-1)^i).$$

Properties of the Unitary Group

The order of the unitary group is

$$|U(n, q)| = q^{n(n-1)/2} \prod_{i=1}^n (q^i - (-1)^i).$$

The centre Z_n of $U(n, q)$ is a cyclic group of order $q + 1$.

Properties of the Unitary Group

The order of the unitary group is

$$|U(n, q)| = q^{n(n-1)/2} \prod_{i=1}^n (q^i - (-1)^i).$$

The centre Z_n of $U(n, q)$ is a cyclic group of order $q + 1$.

We wish to describe two important representations (or characters) of $U(n, q)$.

Properties of the Unitary Group

The order of the unitary group is

$$|U(n, q)| = q^{n(n-1)/2} \prod_{i=1}^n (q^i - (-1)^i).$$

The centre Z_n of $U(n, q)$ is a cyclic group of order $q + 1$.

We wish to describe two important representations (or characters) of $U(n, q)$.

The first is the Steinberg representation, which is irreducible of degree $q^{n(n-1)/2}$. It can be constructed by using the theory of a group with a B, N pair.

Properties of the Unitary Group

The order of the unitary group is

$$|U(n, q)| = q^{n(n-1)/2} \prod_{i=1}^n (q^i - (-1)^i).$$

The centre Z_n of $U(n, q)$ is a cyclic group of order $q + 1$.

We wish to describe two important representations (or characters) of $U(n, q)$.

The first is the Steinberg representation, which is irreducible of degree $q^{n(n-1)/2}$. It can be constructed by using the theory of a group with a B, N pair.

The Steinberg representation is defined over the rational numbers and Z_n is contained in its kernel.

The Steinberg and Weil Characters

Let St_n denote the character of the Steinberg representation. Clearly, St_n is rational-valued.

The Steinberg and Weil Characters

Let St_n denote the character of the Steinberg representation. Clearly, St_n is rational-valued.

We have $St_n(1) = q^{n(n-1)/2}$, which is the p -part of $|U(n, q)|$.

The Steinberg and Weil Characters

Let St_n denote the character of the Steinberg representation. Clearly, St_n is rational-valued.

We have $St_n(1) = q^{n(n-1)/2}$, which is the p -part of $|U(n, q)|$.

Thus St_n has defect 0 modulo p and therefore vanishes on all elements of order divisible by p in $U(n, q)$.

The Steinberg and Weil Characters

Let St_n denote the character of the Steinberg representation. Clearly, St_n is rational-valued.

We have $St_n(1) = q^{n(n-1)/2}$, which is the p -part of $|U(n, q)|$.

Thus St_n has defect 0 modulo p and therefore vanishes on all elements of order divisible by p in $U(n, q)$.

The Weil representation of $U(n, q)$ is a complex representation of degree q^n of this group.

The Steinberg and Weil Characters

Let St_n denote the character of the Steinberg representation. Clearly, St_n is rational-valued.

We have $St_n(1) = q^{n(n-1)/2}$, which is the p -part of $|U(n, q)|$.

Thus St_n has defect 0 modulo p and therefore vanishes on all elements of order divisible by p in $U(n, q)$.

The Weil representation of $U(n, q)$ is a complex representation of degree q^n of this group.

Let ϕ_n denote the character of the Weil representation of $U(n, q)$.

The Steinberg and Weil Characters

Let St_n denote the character of the Steinberg representation. Clearly, St_n is rational-valued.

We have $St_n(1) = q^{n(n-1)/2}$, which is the p -part of $|U(n, q)|$.

Thus St_n has defect 0 modulo p and therefore vanishes on all elements of order divisible by p in $U(n, q)$.

The Weil representation of $U(n, q)$ is a complex representation of degree q^n of this group.

Let ϕ_n denote the character of the Weil representation of $U(n, q)$.

Given an element g of $U(n, q)$, we have

$$\phi_n(g) = (-1)^n (-q)^{c(g)}$$

where $c(g) = \dim \ker (g - I)$.

Irreducible Constituents of the Weil Character

This implies in particular that ϕ_n is rational-valued.

Irreducible Constituents of the Weil Character

This implies in particular that ϕ_n is rational-valued.

If $n \geq 2$, ϕ_n has $q + 1$ irreducible constituents, each occurring with multiplicity 1, which we shall denote by $\phi_n^0, \dots, \phi_n^q$.

Irreducible Constituents of the Weil Character

This implies in particular that ϕ_n is rational-valued.

If $n \geq 2$, ϕ_n has $q + 1$ irreducible constituents, each occurring with multiplicity 1, which we shall denote by $\phi_n^0, \dots, \phi_n^q$.

These irreducible characters may be described in the following way.

Irreducible Constituents of the Weil Character

This implies in particular that ϕ_n is rational-valued.

If $n \geq 2$, ϕ_n has $q + 1$ irreducible constituents, each occurring with multiplicity 1, which we shall denote by $\phi_n^0, \dots, \phi_n^q$.

These irreducible characters may be described in the following way.

Let z denote a generator of Z_n and let ϵ denote a complex root of unity of order $q + 1$.

Irreducible Constituents of the Weil Character

This implies in particular that ϕ_n is rational-valued.

If $n \geq 2$, ϕ_n has $q + 1$ irreducible constituents, each occurring with multiplicity 1, which we shall denote by $\phi_n^0, \dots, \phi_n^q$.

These irreducible characters may be described in the following way.

Let z denote a generator of Z_n and let ϵ denote a complex root of unity of order $q + 1$.

Let λ_i denote the linear character of Z_n given by

$$\lambda_i(z^j) = \epsilon^{ij}$$

for $0 \leq i, j \leq q$.

Degrees of the Constituents

Then we have

$$\phi_n^i|_{Z_n} = \phi_n(1)\lambda_i$$

Degrees of the Constituents

Then we have

$$\phi_n^i|_{Z_n} = \phi_n(1)\lambda_i$$

and we see that the irreducible constituents of ϕ_n are labelled by the irreducible characters of Z_n .

Degrees of the Constituents

Then we have

$$\phi_n^i|_{Z_n} = \phi_n(1)\lambda_i$$

and we see that the irreducible constituents of ϕ_n are labelled by the irreducible characters of Z_n .

In particular, ϕ_n^0 is the unique irreducible constituent of ϕ_n that is trivial on Z_n and it must therefore be rational-valued.

Degrees of the Constituents

Then we have

$$\phi_n^i|_{Z_n} = \phi_n(1)\lambda_i$$

and we see that the irreducible constituents of ϕ_n are labelled by the irreducible characters of Z_n .

In particular, ϕ_n^0 is the unique irreducible constituent of ϕ_n that is trivial on Z_n and it must therefore be rational-valued.

We have the degree formulae

$$\begin{aligned}\phi_n^0(1) &= \frac{q^n - (-1)^n}{q + 1} + (-1)^n \\ \phi_n^i(1) &= \frac{q^n - (-1)^n}{q + 1} \text{ for } 1 \leq i \leq q.\end{aligned}$$

Outline

- 1 The Weil Representation of the Unitary Group
- 2 Reality Properties of the Irreducible Constituents of the Weil Character**
- 3 Tensoring with the Steinberg Character
- 4 Some Consequences

The Real-Valued Irreducible Constituents

Lemma

Suppose that $n \geq 2$.

The Real-Valued Irreducible Constituents

Lemma

Suppose that $n \geq 2$.

The character ϕ_n^0 is the unique real-valued irreducible constituent of ϕ_n when q is a power of 2. It is also rational-valued.

The Real-Valued Irreducible Constituents

Lemma

Suppose that $n \geq 2$.

The character ϕ_n^0 is the unique real-valued irreducible constituent of ϕ_n when q is a power of 2. It is also rational-valued.

When q is a power of an odd prime, there are two real-valued irreducible constituents of ϕ_n , namely,

The Real-Valued Irreducible Constituents

Lemma

Suppose that $n \geq 2$.

The character ϕ_n^0 is the unique real-valued irreducible constituent of ϕ_n when q is a power of 2. It is also rational-valued.

When q is a power of an odd prime, there are two real-valued irreducible constituents of ϕ_n , namely, ϕ_n^0 and ϕ_n^r , where $2r = q + 1$.

The Real-Valued Irreducible Constituents

Lemma

Suppose that $n \geq 2$.

The character ϕ_n^0 is the unique real-valued irreducible constituent of ϕ_n when q is a power of 2. It is also rational-valued.

When q is a power of an odd prime, there are two real-valued irreducible constituents of ϕ_n , namely, ϕ_n^0 and ϕ_n^r , where $2r = q + 1$.

These two characters are in fact rational-valued.

The Frobenius-Schur Indicator

Let G be a finite group and let χ be an irreducible complex character of G .

The Frobenius-Schur Indicator

Let G be a finite group and let χ be an irreducible complex character of G .

We define the Frobenius-Schur indicator $\nu(\chi)$ of χ by

The Frobenius-Schur Indicator

Let G be a finite group and let χ be an irreducible complex character of G .

We define the Frobenius-Schur indicator $\nu(\chi)$ of χ by

$$\nu(\chi) = \frac{1}{|G|} \sum_{g \in G} \chi(g^2).$$

The Frobenius-Schur Indicator

Let G be a finite group and let χ be an irreducible complex character of G .

We define the Frobenius-Schur indicator $\nu(\chi)$ of χ by

$$\nu(\chi) = \frac{1}{|G|} \sum_{g \in G} \chi(g^2).$$

Then $\nu(\chi)$ takes one of the three values, 0, 1 and -1 .

The Frobenius-Schur Indicator

Let G be a finite group and let χ be an irreducible complex character of G .

We define the Frobenius-Schur indicator $\nu(\chi)$ of χ by

$$\nu(\chi) = \frac{1}{|G|} \sum_{g \in G} \chi(g^2).$$

Then $\nu(\chi)$ takes one of the three values, 0, 1 and -1 .

We have $\nu(\chi) = 0$ if χ is not real-valued.

The Frobenius-Schur Indicator

Let G be a finite group and let χ be an irreducible complex character of G .

We define the Frobenius-Schur indicator $\nu(\chi)$ of χ by

$$\nu(\chi) = \frac{1}{|G|} \sum_{g \in G} \chi(g^2).$$

Then $\nu(\chi)$ takes one of the three values, 0, 1 and -1 .

We have $\nu(\chi) = 0$ if χ is not real-valued.

$\nu(\chi) = 1$ if χ is both real-valued and the character of a representation defined over \mathbb{R} .

The Frobenius-Schur Indicator

$\nu(\chi) = -1$ if χ is real-valued and not the character of a representation defined over \mathbb{R} .

The Frobenius-Schur Indicator

$\nu(\chi) = -1$ if χ is real-valued and not the character of a representation defined over \mathbb{R} .

We sometimes say that an irreducible character χ is of *orthogonal* type if $\nu(\chi) = 1$ and of *symplectic* type if $\nu(\chi) = -1$.

The Frobenius-Schur Indicator

$\nu(\chi) = -1$ if χ is real-valued and not the character of a representation defined over \mathbb{R} .

We sometimes say that an irreducible character χ is of *orthogonal* type if $\nu(\chi) = 1$ and of *symplectic* type if $\nu(\chi) = -1$.

It is an interesting problem, not entirely solved, to find the Frobenius-Schur indicator of, say, any irreducible complex character of a finite group of Lie type.

The Frobenius-Schur Indicator

$\nu(\chi) = -1$ if χ is real-valued and not the character of a representation defined over \mathbb{R} .

We sometimes say that an irreducible character χ is of *orthogonal* type if $\nu(\chi) = 1$ and of *symplectic* type if $\nu(\chi) = -1$.

It is an interesting problem, not entirely solved, to find the Frobenius-Schur indicator of, say, any irreducible complex character of a finite group of Lie type.

In this talk, we are looking at a special case in $U(n, q)$.

The Indicators of Constituents of the Weil Character

Theorem

- Suppose that q is odd and n is even. Then $\nu(\phi_n^r) = -1$ (recall $2r = q + 1$).

The Indicators of Constituents of the Weil Character

Theorem

- Suppose that q is odd and n is even. Then $\nu(\phi_n^r) = -1$ (recall $2r = q + 1$).
- Suppose that $n \geq 3$ is odd. Then $\nu(\phi_n^0) = -1$.

The Indicators of Constituents of the Weil Character

Theorem

- Suppose that q is odd and n is even. Then $\nu(\phi_n^r) = -1$ (recall $2r = q + 1$).
- Suppose that $n \geq 3$ is odd. Then $\nu(\phi_n^0) = -1$.
- In all other cases, the real-valued irreducible constituents of the Weil character have indicator equal to 1.

The Indicators of Constituents of the Weil Character

Theorem

- Suppose that q is odd and n is even. Then $\nu(\phi_n^r) = -1$ (recall $2r = q + 1$).
- Suppose that $n \geq 3$ is odd. Then $\nu(\phi_n^0) = -1$.
- In all other cases, the real-valued irreducible constituents of the Weil character have indicator equal to 1.

The proof of this theorem is by induction on n , making use of knowledge of the restriction of ϕ_n to smaller unitary subgroups.

Outline

- 1 The Weil Representation of the Unitary Group
- 2 Reality Properties of the Irreducible Constituents of the Weil Character
- 3 Tensoring with the Steinberg Character**
- 4 Some Consequences

Theorem of Hiss-Zaleski-Brunat

When q is odd, we let ϵ_n denote the linear character of order 2 of $U(n, q)$ given by

Theorem of Hiss-Zaleski-Brunat

When q is odd, we let ϵ_n denote the linear character of order 2 of $U(n, q)$ given by

$$\epsilon(g) = \det(g)^r,$$

where, as usual, $2r = q + 1$.

Theorem of Hiss-Zaleski-Brunat

When q is odd, we let ϵ_n denote the linear character of order 2 of $U(n, q)$ given by

$$\epsilon(g) = \det(g)^r,$$

where, as usual, $2r = q + 1$.

We define the character ω_n of $U(n, q)$ to equal ϕ_n if q is even and to equal $\epsilon_n \phi_n$ if q is odd.

Theorem of Hiss-Zaleski-Brunat

When q is odd, we let ϵ_n denote the linear character of order 2 of $U(n, q)$ given by

$$\epsilon(g) = \det(g)^r,$$

where, as usual, $2r = q + 1$.

We define the character ω_n of $U(n, q)$ to equal ϕ_n if q is even and to equal $\epsilon_n \phi_n$ if q is odd.

We now apply a theorem of Hiss, Zaleski and Brunat, *The Weil-Steinberg character of finite classical groups*. With an appendix by Olivier Brunat. Represent. Theory **13** (2009), 427–459.

Multiplicity-Free Product

Theorem: HZB (2009)

The product $\omega_n St_n$ is a multiplicity-free character of $U(n, q)$.

Multiplicity-Free Product

Theorem: HZB (2009)

The product $\omega_n St_n$ is a multiplicity-free character of $U(n, q)$.

Furthermore,

$$St_{n+1}|_{U(n,q)} = \omega_n St_n.$$

Multiplicity-Free Product

Theorem: HZB (2009)

The product $\omega_n St_n$ is a multiplicity-free character of $U(n, q)$.

Furthermore,

$$St_{n+1}|_{U(n,q)} = \omega_n St_n.$$

It follows that $\phi_n St_n$ is also multiplicity-free and its irreducible constituents have the form $\epsilon_n \chi$, where χ is any irreducible constituent of $\omega_n St_n$.

Multiplicity-Free Product

Theorem: HZB (2009)

The product $\omega_n St_n$ is a multiplicity-free character of $U(n, q)$.

Furthermore,

$$St_{n+1}|_{U(n, q)} = \omega_n St_n.$$

It follows that $\phi_n St_n$ is also multiplicity-free and its irreducible constituents have the form $\epsilon_n \chi$, where χ is any irreducible constituent of $\omega_n St_n$.

We now prove a result about the Schur index of each constituent of $\phi_n St_n$.

Schur Index 1

Corollary

Each irreducible constituent of $\phi_n St_n$ has Schur index 1 over \mathbb{Q} .

Schur Index 1

Corollary

Each irreducible constituent of $\phi_n St_n$ has Schur index 1 over \mathbb{Q} .

Proof

St_{n+1} is the character of a representation of $U(n+1, q)$ defined over \mathbb{Q} .

Schur Index 1

Corollary

Each irreducible constituent of $\phi_n St_n$ has Schur index 1 over \mathbb{Q} .

Proof

St_{n+1} is the character of a representation of $U(n+1, q)$ defined over \mathbb{Q} .

It follows that the restriction of St_{n+1} to $U(n, q)$ is the character of a representation of $U(n, q)$ defined over \mathbb{Q} .

Schur Index 1

Corollary

Each irreducible constituent of $\phi_n St_n$ has Schur index 1 over \mathbb{Q} .

Proof

St_{n+1} is the character of a representation of $U(n+1, q)$ defined over \mathbb{Q} .

It follows that the restriction of St_{n+1} to $U(n, q)$ is the character of a representation of $U(n, q)$ defined over \mathbb{Q} .

Since each irreducible constituent of this restriction occurs with multiplicity 1, the result is clear, as we know now that $\phi_n St_n$ is the character of this restriction.

Outline

- 1 The Weil Representation of the Unitary Group
- 2 Reality Properties of the Irreducible Constituents of the Weil Character
- 3 Tensoring with the Steinberg Character
- 4 Some Consequences**

More on Schur Indices

While we know that $\phi_n St_n$ is rational-valued, we do not necessarily know much about the field of values of its irreducible constituents.

More on Schur Indices

While we know that $\phi_n St_n$ is rational-valued, we do not necessarily know much about the field of values of its irreducible constituents.

However, we know that it must also be the case that each irreducible constituent of the products $\phi_n^i St_n$, for $0 \leq i \leq q$, has Schur index 1 over \mathbb{Q} , since $\phi_n^i St_n$ is a subcharacter of $\phi_n St_n$.

More on Schur Indices

While we know that $\phi_n St_n$ is rational-valued, we do not necessarily know much about the field of values of its irreducible constituents.

However, we know that it must also be the case that each irreducible constituent of the products $\phi_n^i St_n$, for $0 \leq i \leq q$, has Schur index 1 over \mathbb{Q} , since $\phi_n^i St_n$ is a subcharacter of $\phi_n St_n$.

This has interesting consequences when we choose ϕ_n^i to be a character with $\nu(\phi_n^i) = -1$.

Non-Real Constituents

Corollary

Suppose that n is odd and at least 3. Then no irreducible constituent of $\phi_n^0 St_n$ is real-valued.

Non-Real Constituents

Corollary

Suppose that n is odd and at least 3. Then no irreducible constituent of $\phi_n^0 St_n$ is real-valued.

The same conclusion holds for the irreducible constituents of $\phi_n^r St_n$ if n is even and q is odd.

Non-Real Constituents

Corollary

Suppose that n is odd and at least 3. Then no irreducible constituent of $\phi_n^0 St_n$ is real-valued.

The same conclusion holds for the irreducible constituents of $\phi_n^r St_n$ if n is even and q is odd.

Proof

Briefly, this follows since St_n is real-valued with $\nu(St_n) = 1$, and, under the given hypotheses, ϕ_n^i is real-valued with $\nu(\phi_n^i) = -1$.

Non-Real Constituents

Corollary

Suppose that n is odd and at least 3. Then no irreducible constituent of $\phi_n^0 St_n$ is real-valued.

The same conclusion holds for the irreducible constituents of $\phi_n^r St_n$ if n is even and q is odd.

Proof

Briefly, this follows since St_n is real-valued with $\nu(St_n) = 1$, and, under the given hypotheses, ϕ_n^i is real-valued with $\nu(\phi_n^i) = -1$.

Given this, and the fact that all irreducible constituents of $\phi_n^i St_n$ have Schur index 1, these constituents cannot be real-valued.

Corollary

Let χ be any real-valued irreducible character of $U(n, q)$.

Corollary

Let χ be any real-valued irreducible character of $U(n, q)$.

Then ϕ_n^0 is not a constituent of χSt_n if n is odd and at least 3.

Corollary

Let χ be any real-valued irreducible character of $U(n, q)$.

Then ϕ_n^0 is not a constituent of χSt_n if n is odd and at least 3.

Similarly, ϕ_n^r is not a constituent of χSt_n if n is even and q is odd.

Corollary

Let χ be any real-valued irreducible character of $U(n, q)$.

Then ϕ_n^0 is not a constituent of χSt_n if n is odd and at least 3.

Similarly, ϕ_n^r is not a constituent of χSt_n if n is even and q is odd.

Proof

We just consider the case that n is odd.

Corollary

Let χ be any real-valued irreducible character of $U(n, q)$.

Then ϕ_n^0 is not a constituent of χSt_n if n is odd and at least 3.

Similarly, ϕ_n^r is not a constituent of χSt_n if n is even and q is odd.

Proof

We just consider the case that n is odd.

Since we are assuming that χ is real-valued,

$$(\phi_n^0 St_n, \chi) = 0,$$

Corollary

Let χ be any real-valued irreducible character of $U(n, q)$.

Then ϕ_n^0 is not a constituent of χSt_n if n is odd and at least 3.

Similarly, ϕ_n^r is not a constituent of χSt_n if n is even and q is odd.

Proof

We just consider the case that n is odd.

Since we are assuming that χ is real-valued,

$$(\phi_n^0 St_n, \chi) = 0,$$

where the parentheses denote the usual inner product of characters.

Proof Concluded

However, as all characters involved in the inner product are real-valued, we have

$$(\chi St_n, \phi_n^0) = (\phi_n^0 St_n, \chi) = 0.$$

Proof Concluded

However, as all characters involved in the inner product are real-valued, we have

$$(\chi St_n, \phi_n^0) = (\phi_n^0 St_n, \chi) = 0.$$

Thus ϕ_n^0 is not a constituent of χSt_n .

The Steinberg Square

Corollary

When n is odd and at least 3, ϕ_n^0 is not a constituent of St_n^2 .

The Steinberg Square

Corollary

When n is odd and at least 3, ϕ_n^0 is not a constituent of St_n^2 .

This is an immediate consequence of the previous corollary and it may seem strange that we have drawn attention to it.

The Steinberg Square

Corollary

When n is odd and at least 3, ϕ_n^0 is not a constituent of St_n^2 .

This is an immediate consequence of the previous corollary and it may seem strange that we have drawn attention to it.

As far as we know, St^2 does contain every other irreducible character χ of $U(n, q)$ as a constituent, provided Z_n is contained in the kernel of χ .

The Steinberg Square

Corollary

When n is odd and at least 3, ϕ_n^0 is not a constituent of St_n^2 .

This is an immediate consequence of the previous corollary and it may seem strange that we have drawn attention to it.

As far as we know, St^2 does contain every other irreducible character χ of $U(n, q)$ as a constituent, provided Z_n is contained in the kernel of χ .

Indeed, it may be the case that the square of the Steinberg character of a simple group of Lie type contains all irreducible characters of the group, except for this special case.

The Steinberg Square

We note that if G is a finite group of Lie type over \mathbb{F}_q , where q is a power of 2, and St is the Steinberg character of G , then St occurs with *odd* multiplicity in St^2 .

The Steinberg Square

We note that if G is a finite group of Lie type over \mathbb{F}_q , where q is a power of 2, and St is the Steinberg character of G , then St occurs with *odd* multiplicity in St^2 .

Furthermore, St occurs with non-zero multiplicity in the symmetric part of the tensor square of St^2 .

The Steinberg Square

We note that if G is a finite group of Lie type over \mathbb{F}_q , where q is a power of 2, and St is the Steinberg character of G , then St occurs with *odd* multiplicity in St^2 .

Furthermore, St occurs with non-zero multiplicity in the symmetric part of the tensor square of St^2 .

This is not necessarily true when q is odd.